

Ph-507. Midterm Exam, March 10, 2008.

HINT: all three problems may be done with very modest amount of calculation. However, feel free to do the problems the way you like.

PROBLEM 1 (5pts)

A particle of mass m with total energy E , is scattered by central field

$$U(r) = \frac{k}{r^2}.$$

It was found that at certain finite values of impact parameter s there is no apparent scattering. Find those values, and sketch the corresponding trajectories. The potential is *not weak*.

PROBLEM 2 (5 pts.)

A space station has a circular orbit about certain planet. The magnitude of its linear momentum in the reference frame of the planet is p_0 . Find the minimal impulse that the station's engines must generate to change the period of its orbital motion by factor α . Assume the mass of the station to remain constant.

PROBLEM 3 (5 pts.)

The Lagrangian of certain particle is

$$L = \frac{m\dot{x}^2}{2} - [U_0(x) + \delta U(x)]$$

Assume that the solution for the unperturbed problem, $\delta U(x) = 0$, is known. Derive the expression for the period $T(E)$, which takes into account all perturbations, up to the n -th order in δU .

Solution to Problem 1

$$\Delta\theta = 2 \int_{r_{\min}}^{\infty} \frac{1}{\sqrt{1 - (s^2 + k/E)/r^2}} \frac{sdr}{r^2} = 2 \int_{r_{\min}}^{\infty} \frac{1}{\sqrt{1 - S^2/r^2}} \frac{sdr}{r^2}$$

Here $s^2 + k/E = S^2$. Use the fact that $\theta_m = \pi$ for a free particle ($U = 0$):

$$\Delta\theta = \frac{2s}{S} \int_{r_{\min}}^{\infty} \frac{1}{\sqrt{1 - S^2/r^2}} \frac{Sdr}{r^2} = \frac{\pi s}{\sqrt{s^2 + k/E}}$$

We now have to find s at which $\Delta\theta = \pi + 2\pi N$:

$$1 + \frac{k}{s^2 E} = \frac{1}{(1 + 2N)^2}, \text{ therefore } s = \frac{1}{N + 1/2} \sqrt{-\frac{kN(1 + N)}{E}}$$

Note that solution only exists for an attractive potential, $k < 0$. It corresponds to the trajectory which makes N complete turns about the scattering center, and returns to the original direction of the flight.

Solution to Problem 2

The period is the function of the total energy. Its change by the factor α implies that the energy changes by $\alpha^{-2/3}$:

$$K + U = \alpha^{-2/3} (K_0 + U_0)$$

Note that the impulse from engine will change the kinetic but not potential energy ($U = U_0$), and that according to virial theorem $U_0 = -2K_0$.

Therefore,

$$K = K_0 (2 - \alpha^{-2/3})$$

This means that the engine should change the momentum of the station from p_0 to $p_0 \sqrt{2 - \alpha^{-2/3}}$. The minimal impulse needed to do so is,

$$P = p_0 \left| \sqrt{2 - \alpha^{-2/3}} - 1 \right|$$

Solution to Problem 3

$$\begin{aligned} T &= \left(\int_{x_1 + \delta x_1}^{x_2 + \delta x_2'} \frac{\sqrt{2m} dx}{\sqrt{E - U(x) - \delta U(x)}} \right) = \frac{\sqrt{2m}}{\prod_{k=1}^n (k - 1/2)} \frac{\partial^n}{\partial E^n} \int_{x_1}^{x_2} (E - U(x) - \delta U(x))^{n-1/2} dx \\ &= \frac{\sqrt{2m}}{\prod_{k=1}^n (k - 1/2)} \frac{\partial^n}{\partial E^n} \int_{x_1}^{x_2} \sum_{m=0}^n \frac{(-1)^m}{m!} \delta U^m(x) \frac{\partial^m}{\partial E^m} (E - U(x))^{n-1/2} dx = \sum_{m=0}^n \frac{(-1)^m}{m!} \frac{\partial^m}{\partial E^m} \int_{x_1}^{x_2} \frac{\sqrt{2m} \delta U^m(x)}{\sqrt{E - U}} dx \\ &= \sum_{m=0}^n \frac{(-1)^m}{m!} \frac{\partial^m}{\partial E^m} T_0 \langle \delta U^m \rangle \end{aligned}$$