

STABILITY IN THE HOMOLOGY OF UNIPOTENT GROUPS

ANDREW PUTMAN, STEVEN V SAM, AND ANDREW SNOWDEN

ABSTRACT. Let R be a (not necessarily commutative) ring whose additive group is finitely generated and let $U_n(R) \subset \mathbf{GL}_n(R)$ be the group of upper-triangular unipotent matrices over R . We study how the homology groups of $U_n(R)$ vary with n from the point of view of representation stability. Our main theorem asserts that if for each n we have representations M_n of $U_n(R)$ over a ring \mathbf{k} that are appropriately compatible and satisfy suitable finiteness hypotheses, then the rule $[n] \mapsto \mathbf{H}_i(U_n(R), M_n)$ defines a finitely generated \mathbf{OI} -module. As a consequence, if \mathbf{k} is a field then $\dim \mathbf{H}_i(U_n(R), \mathbf{k})$ is eventually equal to a polynomial in n . We also prove similar results for the Iwahori subgroups of $\mathbf{GL}_n(\mathcal{O})$ for number rings \mathcal{O} .

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 2. Representations of categories | 4 |
| 3. The category \mathbf{OI} and variants | 7 |
| 4. The category \mathbf{OVI} and its variants | 10 |
| 5. Noetherianity of \mathbf{OVI} -modules | 12 |
| 6. Homology of \mathbf{OVI} -modules | 26 |
| 7. Application to Iwahori subgroups | 30 |
| References | 31 |

1. INTRODUCTION

Let $U_n \subset \mathbf{GL}_n$ be the group of upper-triangular unipotent matrices and let R be a ring (not necessarily commutative). In this paper, we study how the homology groups of $U_n(R)$ vary as n changes. The following is a special case of one of our main theorems:

Theorem 1.1. *Let R be a ring whose additive group is finitely generated and let \mathbf{k} be a field. For all $i \geq 0$, there exists some $f_i(t) \in \mathbf{Q}[t]$ such that $\dim \mathbf{H}_i(U_n(R), \mathbf{k}) = f_i(n)$ for $n \gg 0$.*

For instance, the ring R in Theorem 1.1 could be a number ring. We prove this using the methods of representation stability: we endow the sequence of groups $\{\mathbf{H}_i(U_n(R), \mathbf{k})\}_{n \geq 0}$ with additional structure (in this case, that of an \mathbf{OI} -module), and deduce concrete results like the above theorem from the existence of this structure. An interesting feature here is that, unlike in similar results proved under the aegis of representation stability, the individual homology groups $\mathbf{H}_i(U_n(R), \mathbf{k})$ themselves are not representations of a sequence of groups.

Date: November 28, 2017.

2010 *Mathematics Subject Classification.* 16P40, 20J05.

AP was partially supported by NSF DMS-1737434.

SS was partially supported by NSF DMS-1500069, DMS-1651327, and a Sloan research fellowship.

AS was partially supported by NSF DMS-1303082, DMS-1453893, and a Sloan research fellowship.

Example 1.2. The case $R = \mathbf{Z}$ and $\mathbf{k} = \mathbf{Q}$ of Theorem 1.1 follows from work of Dwyer [D, Theorem 1.1]. He shows that the dimension of $H_i(U_n(\mathbf{Z}), \mathbf{Q})$ is the number of permutations in S_n with length i , where the length of a permutation σ is the number of pairs $i < j$ such that $\sigma(i) > \sigma(j)$. Denote this number by $I(i, n)$. We claim that $n \mapsto I(i, n)$ is a polynomial of degree i for $n \gg 0$. As an aside, this shows that the degree of the polynomials $f_i(t)$ in Theorem 1.1 cannot be bounded as we let i vary. We prove the claim by induction. For $i = 1$, we have $I(1, n) = n - 1$ for $n > 0$. In general, we have the identity

$$\sum_{i \geq 0} I(i, n)q^i = (1 + q)(1 + q + q^2) \cdots (1 + q + q^2 + \cdots + q^{n-1});$$

see [St, Corollary 1.3.13]. It follows that $I(i, n) - I(i, n - 1) = \sum_{j=0}^{i-1} I(j, n - 1)$ for $n > i$. By induction, the right hand side is a polynomial of degree $i - 1$ for $n \gg 0$. Hence $I(i, n)$ is a polynomial of degree i for $n \gg 0$, as claimed. \square

1.1. Main results. Our main result is a refined version of Theorem 1.1 where we allow systems of non-trivial coefficients and give a stronger conclusion. To formulate this, we must make sense of a “system” of representations of $U_n(R)$. For this, we introduce the category $\mathbf{OVI}(R)$. An object of $\mathbf{OVI}(R)$ is a finite rank free R -module equipped with a totally ordered basis. A morphism of $\mathbf{OVI}(R)$ is a map of R -modules that is upper-triangular with respect to the distinguished ordered bases (see §4.1). An $\mathbf{OVI}(R)$ -module over a commutative ring \mathbf{k} is a functor $\mathbf{OVI}(R) \rightarrow \text{Mod}_{\mathbf{k}}$. Every object in $\mathbf{OVI}(R)$ is isomorphic to R^n equipped with its standard basis for some n , and the automorphism group of this object is the group $U_n(R)$. Thus an $\mathbf{OVI}(R)$ -module M gives rise to a sequence $\{M_n\}_{n \geq 0}$, where $M_n = M(R^n)$ is a representation of $U_n(R)$, and therefore provides a reasonable notion of a system of $U_n(R)$ representations.

Let M be an $\mathbf{OVI}(R)$ -module, and fix $i \geq 0$. For each n , we consider the homology group $H_i(U_n(R), M_n)$. Letting $[n]$ denote the ordered set $\{1, \dots, n\}$, if $[n] \rightarrow [m]$ is an order-preserving injection of finite sets then there is an associated morphism $R^n \rightarrow R^m$ in $\mathbf{OVI}(R)$. This gives a map $M_n \rightarrow M_m$, which induces a map $H_i(U_n(R), M_n) \rightarrow H_i(U_m(R), M_m)$. This suggests that $[n] \mapsto H_i(U_n(R), M_n)$ defines an \mathbf{OI} -module, where \mathbf{OI} is the category whose objects are finite totally ordered sets and whose morphisms are order-preserving injections. We show that this is indeed the case, and denote this \mathbf{OI} -module by $\mathbf{H}_i(\mathbf{U}, M)$.

We can now state our main theorem:

Theorem 1.3. *Let R be a ring whose additive group is finitely generated, let \mathbf{k} be a noetherian commutative ring, and M be a finitely generated $\mathbf{OVI}(R)$ -module over \mathbf{k} . Then $\mathbf{H}_i(\mathbf{U}, M)$ is a finitely generated \mathbf{OI} -module for all $i \geq 0$.*

Theorem 1.1 follows immediately from this theorem by taking M_n to be the trivial representation of $U_n(R)$ for all n and appealing to the fact that a finitely generated \mathbf{OI} -module over a field has eventually polynomial dimension (see Proposition 3.5 below).

A key result used in the proof of Theorem 1.3 is the following noetherianity result:

Theorem 1.4. *Let R be a ring whose additive group is finitely generated and let \mathbf{k} be a noetherian commutative ring. Then the category of $\mathbf{OVI}(R)$ -modules over \mathbf{k} is locally noetherian, that is, any submodule of a finitely generated module is finitely generated.*

A surprising aspect of Theorem 1.4 is that, in contrast to previous work on categories of R -modules in representation stability (for instance, the results of [PuSa] and [SaSn]), the ring

R is allowed to be infinite. The groups of automorphisms studied in this previous work were $\mathbf{GL}_n(R)$ and its relatives, and R was required to be finite since the group ring of $\mathbf{GL}_n(R)$ is not noetherian when R is infinite. In our situation, the automorphism groups are $U_n(R)$. When the additive group of R is finitely generated, these groups are virtually polycyclic, and a classical result of Philip Hall [Ha] says that group rings of virtually polycyclic groups are noetherian. Our proof of Theorem 1.4 is inspired in part by Hall's proof of this fact.

Remark 1.5. It is easy to see that this result is false if the additive group of R is not finitely generated (see §5.4). \square

Remark 1.6. When the ring R is finite, we in fact show that the category of $\mathbf{OVI}(R)$ -modules is quasi-Gröbner in the sense of [SaSn, §4], which implies local noetherianity (but is stronger). In the general case, we do not show that the category of $\mathbf{OVI}(R)$ -modules is quasi-Gröbner (and expect that it is not), and the proof of local noetherianity is far more difficult. \square

Remark 1.7. \mathbf{OI} -modules are closely related to the \mathbf{FI} -modules of Church–Ellenberg–Farb [CEF]. The second two authors used \mathbf{OI} -modules in [SaSn, §7] to study \mathbf{FI} -modules. However, as far as we are aware, Theorem 1.3 is the first time \mathbf{OI} -modules have been used as the primary structure in representation stability. \square

Remark 1.8. An \mathbf{OI} -module is the same as a semi-simplicial object of $\text{Mod}_{\mathbf{k}}^{\text{op}}$. \square

1.2. Application to Iwahori groups. Let \mathcal{O} be a number ring and let \mathbf{k} be a commutative noetherian ring. A classical result of van der Kallen [VdK] says that the homology of the group $\mathbf{GL}_n(\mathcal{O})$ stabilizes: for any fixed i the canonical map

$$H_i(\mathbf{GL}_n(\mathcal{O}), \mathbf{k}) \rightarrow H_i(\mathbf{GL}_{n+1}(\mathcal{O}), \mathbf{k})$$

is an isomorphism for $n \gg 0$. In particular, if \mathbf{k} is a field then the dimension of $H_i(\mathbf{GL}_n(\mathcal{O}), \mathbf{k})$ is eventually constant.

Now let \mathfrak{a} be a nonzero proper ideal in \mathcal{O} and let $\mathbf{GL}_n(\mathcal{O}, \mathfrak{a})$ be the principal congruence subgroup of level \mathfrak{a} , i.e., the subgroup of $\mathbf{GL}_n(\mathcal{O})$ consisting of matrices that are congruent to the identity modulo \mathfrak{a} . The homology of these groups does not stabilize; for instance, for $\ell \geq 2$ and $n \geq 3$ the abelianization of $\mathbf{GL}_n(\mathbf{Z}, \ell\mathbf{Z})$ is $(\mathbf{Z}/\ell)^{n^2-1}$ (see [LSz]). Building on work of the first author [Pu], Church–Ellenberg–Farb–Nagpal [CEFN] proved instead that the homology of $\mathbf{GL}_n(\mathcal{O}, \mathfrak{a})$ satisfies a version of representation stability: the rule $[n] \mapsto H_i(\mathbf{GL}_n(\mathcal{O}, \mathfrak{a}), \mathbf{k})$ defines a finitely generated \mathbf{FI} -module. Consequently, when \mathbf{k} is a field, the dimension is eventually polynomial.

The Iwahori subgroup $\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a})$ is the subgroup of $\mathbf{GL}_n(\mathcal{O})$ consisting of matrices that are upper-triangular modulo \mathfrak{a} . Using Theorem 1.3, we prove an analog of Church–Ellenberg–Farb–Nagpal's result for $\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a})$:

Theorem 1.9. *Let \mathcal{O} be a number ring, let $\mathfrak{a} \subset \mathcal{O}$ be a nonzero proper ideal, and let \mathbf{k} be a commutative noetherian ring. Then the following hold for all $i \geq 0$.*

- *The rule $[n] \mapsto H_i(\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}), \mathbf{k})$ defines a finitely generated \mathbf{OI} -module over \mathbf{k} .*
- *If \mathbf{k} is a field then there is a polynomial $f \in \mathbf{Q}[t]$ such that $\dim H_i(\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}), \mathbf{k}) = f(n)$ for $n \gg 0$.*

1.3. Outline. In §2 we review generalities on modules over categories. In §3 we introduce the category \mathbf{OI} and its variants $\mathbf{OI}(d)$ and establish basic results about them. In §4 we

introduce the category $\mathbf{OVI}(R)$ and its variants $\mathbf{OVI}(R, d)$ and establish basic results about them. In §5, we prove the main noetherianity result for $\mathbf{OVI}(R)$ (Theorem 1.4). In §6 we prove the main result of the paper (Theorem 1.3). Finally, in §7 we prove Theorem 1.9.

1.4. Notation. Throughout, \mathbf{k} denotes a commutative ring, typically noetherian. Unless otherwise specified, $1 \neq 0$ in all of our rings. For a fixed category \mathcal{C} , we write $\underline{\mathbf{k}}$ for the constant functor $\mathcal{C} \rightarrow \text{Mod}_{\mathbf{k}}$ taking everything to \mathbf{k} and all morphisms to the identity. We let $B_n \subset \mathbf{GL}_n$ be the group of upper-triangular matrices, and $U_n \subset B_n$ the subgroup where the diagonal entries are equal to 1. We use R to denote the ring appearing in the definition of $\mathbf{OVI}(R)$, and that is typically plugged in to U_n or B_n . We generally do not require it to be commutative. We set $[0] = \emptyset$, and if n is a positive integer, then $[n]$ denotes the set $\{1, \dots, n\}$.

Acknowledgments. We thank Benjamin Steinberg for pointing out a significant simplification to the proof of Lemma 5.4.

2. REPRESENTATIONS OF CATEGORIES

2.1. Generalities. Let \mathcal{C} be a category and let \mathbf{k} be a noetherian commutative ring. A \mathcal{C} -**module** over \mathbf{k} is a functor $M: \mathcal{C} \rightarrow \text{Mod}_{\mathbf{k}}$. For an object $x \in \mathcal{C}$, we denote by M_x the image of x under M . Denote the category of \mathcal{C} -modules by $\text{Rep}_{\mathbf{k}}(\mathcal{C})$. It is an abelian category. For each $x \in \mathcal{C}$, we define a \mathcal{C} -module P_x via the formula $(P_x)_y = \mathbf{k}[\text{Hom}(x, y)]$. One easily sees that for any \mathcal{C} -module M one has a natural identification $\text{Hom}(P_x, M) = M_x$. It follows that P_x is a projective \mathcal{C} -module; we call it the **principal projective** at x . A general \mathcal{C} -module M is finitely generated if and only if there exists a surjection $\bigoplus_{i=1}^k P_{x_i} \rightarrow M$ for some $x_1, \dots, x_k \in \mathcal{C}$. A \mathcal{C} -module is said to be **noetherian** if all of its submodules are finitely generated, and the category $\text{Rep}_{\mathbf{k}}(\mathcal{C})$ is said to be **locally noetherian** if all finitely generated objects are noetherian.

If $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ is a functor and M is a \mathcal{D} -module then the **pullback** of M along Φ , denoted $\Phi^*(M)$, is the \mathcal{C} -module defined via the formula $\Phi^*(M) = M \circ \Phi$, so that $\Phi^*(M)_x = M_{\Phi(x)}$. We now review how the pullback operation interacts with finite generation. The following definition is [SaSn, Def. 3.2.1]:

Definition 2.1. We say that a functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ satisfies **property (F)** if the following condition holds for all $y \in \mathcal{D}$. There exist finitely many objects $x_1, \dots, x_n \in \mathcal{C}$ together with morphisms $f_i: y \rightarrow \Phi(x_i)$ in \mathcal{D} with the following property: for any $x \in \mathcal{C}$ and any morphism $f: y \rightarrow \Phi(x)$ in \mathcal{D} , there exists an i , and a morphism $g: x_i \rightarrow x$ in \mathcal{C} , such that $f = \Phi(g) \circ f_i$. \square

Definition 2.2. A category \mathcal{C} satisfies property (F) if the diagonal $\mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$ satisfies property (F). \square

The importance of these definitions is due to the following results.

Proposition 2.3. *A functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ satisfies property (F) if and only if $\Phi^*(M)$ is a finitely generated \mathcal{C} -module for all finitely generated \mathcal{D} -modules M .*

Proof. See [SaSn, Prop. 3.2.3]. \square

Recall that a functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ is **essentially surjective** if for all $y \in \mathcal{D}$, there exists some $x \in \mathcal{C}$ such that $\Phi(x)$ is isomorphic to y .

Proposition 2.4. *Let \mathcal{C} be a category such that $\text{Rep}_{\mathbf{k}}(\mathcal{C})$ is locally noetherian and let $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ be an essentially surjective functor satisfying property (F). Then $\text{Rep}_{\mathbf{k}}(\mathcal{D})$ is locally noetherian.*

Proof. See [SaSn, Cor. 3.2.5]. □

If \mathcal{C} is a category and M_1 and M_2 are \mathcal{C} -modules, then we define $M_1 \otimes M_2$ to be the \mathcal{C} -module defined by the formula $(M_1 \otimes M_2)_x = (M_1)_x \otimes (M_2)_x$ for all $x \in \mathcal{C}$.

Proposition 2.5. *Let \mathcal{C} be a category that satisfies property (F) and let M and N be finitely generated \mathcal{C} -modules. Then $M \otimes N$ is finitely generated.*

Proof. See [SaSn, Prop. 3.3.2]. □

We require a slight variant of the above proposition. We say that a \mathcal{C} -module M is **generated in finite degrees** if there exist $x_1, \dots, x_k \in \mathcal{C}$ such that M is generated by the M_{x_i} , that is, the canonical map $\bigoplus_{i=1}^k M_{x_i} \otimes P_{x_i} \rightarrow M$ is surjective. Note that if M is generated in finite degrees and M_x is a finitely generated \mathbf{k} -module for all $x \in \mathcal{C}$ then M is finitely generated.

Proposition 2.6. *Let \mathcal{C} be a category that satisfies property (F) and let M and N be \mathcal{C} -modules generated in finite degrees. Then $M \otimes N$ is generated in finite degrees.*

Proof. Observe that (a) a finite sum of \mathcal{C} -modules generated in finite degrees is generated in finite degrees; (b) if K is a \mathcal{C} -module generated in finite degrees and U is any \mathbf{k} -module then $U \otimes K$ is generated in finite degrees; (c) any quotient of a \mathcal{C} -module generated in finite degrees is generated in finite degrees. Now, choose surjections $\bigoplus_{i=1}^k V_i \otimes P_{x_i} \rightarrow M$ and $\bigoplus_{j=1}^{\ell} W_j \otimes P_{y_j} \rightarrow N$, where the x_i and y_j are objects of \mathcal{C} and the V_i and W_j are \mathbf{k} -modules (one can take $V_i = M_{x_i}$ and $W_j = N_{y_j}$). We thus have a surjection

$$\bigoplus_{i,j} V_i \otimes W_j \otimes P_{x_i} \otimes P_{y_j} \rightarrow M \otimes N.$$

Since \mathcal{C} satisfies property (F), each $P_{x_i} \otimes P_{y_j}$ is finitely generated (Proposition 2.5). Thus each term in the sum is generated in finite degrees by (b); since the sum is finite, it is generated in finite degree by (a); and so we conclude $M \otimes N$ is generated in finite degrees by (c). □

Now we recall the notion of a Gröbner category. See [SaSn, §4.3] for more details.

Definition 2.7. Let \mathcal{C} be an essentially small category, i.e., there exists a set I containing a unique representative of each isomorphism class in \mathcal{C} . For $x \in \mathcal{C}$, define $|S_x| = \coprod_{y \in I} \text{Hom}(x, y)$. Partially order $|S_x|$ by defining $f \preceq g$ if there exists a morphism h such that $g = hf$. We say that \mathcal{C} is **Gröbner** if the following holds for all $x \in \mathcal{C}$.

- The poset $(|S_x|, \preceq)$ is noetherian.
- $|S_x|$ admits a total ordering \leq with the following two properties.
 - The ordering \leq is compatible with left composition, i.e., $f \leq g$ implies $hf \leq hg$.
 - The restriction of \leq to each $\text{Hom}(x, y)$ is a well-ordering.

We say that \mathcal{C} is **quasi-Gröbner** if there exists a Gröbner category \mathcal{C}' and an essentially surjective functor $\mathcal{C}' \rightarrow \mathcal{C}$ satisfying property (F). □

The key result about quasi-Gröbner categories is the following [SaSn, Theorem 4.3.2]:

Theorem 2.8. *Let \mathcal{C} be a quasi-Gröbner category. Then for any noetherian commutative ring \mathbf{k} , the category $\text{Rep}_{\mathbf{k}}(\mathcal{C})$ is locally noetherian.*

2.2. Kan extension. Let $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. The pullback functor Φ^* on modules admits a left adjoint $\Phi_!$ called the **left Kan extension**. It also admits a right adjoint Φ_* called the **right Kan extension**, but we will not need this.

The left Kan extension can be described explicitly as follows. Let y be an object of \mathcal{D} . Define a category $\mathcal{C}_{/y}$ as follows. An object of $\mathcal{C}_{/y}$ is a pair (x, f) , where x is an object of \mathcal{C} and $f: \Phi(x) \rightarrow y$ is a morphism in \mathcal{D} . A morphism $(x', f') \rightarrow (x, f)$ in $\mathcal{C}_{/y}$ is a morphism $g: x' \rightarrow x$ in \mathcal{C} such that $f' = f \circ \Phi(g)$. Suppose now that M is a \mathcal{C} -module over \mathbf{k} . For $y \in \mathcal{D}$, define $M|_{\mathcal{C}_{/y}}$ to be the $\mathcal{C}_{/y}$ -module defined via the formula $(M|_{\mathcal{C}_{/y}})_{(x,f)} = M_x$. We then have

$$\Phi_!(M)_y = \text{colim}(M|_{\mathcal{C}_{/y}}).$$

That is, the value of $\Phi_!(M)$ on y is the colimit of the functor $M|_{\mathcal{C}_{/y}}: \mathcal{C}_{/y} \rightarrow \text{Mod}_{\mathbf{k}}$. In certain cases, there is an even nicer description:

Proposition 2.9. *Let $\Phi: \mathcal{C} \rightarrow \mathcal{D}$ be a faithful functor. Assume that for all $x', x \in \mathcal{C}$, the $\text{Aut}(\Phi(x))$ -orbit of any element of $\text{Hom}_{\mathcal{D}}(\Phi(x'), \Phi(x))$ contains an element of the form $\Phi(f)$ for some $f \in \text{Hom}_{\mathcal{C}}(x', x)$ that is unique up to the action of $\text{Aut}(x)$. Let M be a \mathcal{C} -module. Then for all $x \in \mathcal{C}$ we have a canonical isomorphism*

$$\Psi_!(M)_{\Phi(x)} = \text{Ind}_{\text{Aut}(x)}^{\text{Aut}(\Phi(x))}(M_x).$$

Proof. Let $\{h_i\}_{i \in I}$ be a set of coset representatives for $\text{Aut}(\Phi(x))/\text{Aut}(x)$. For each $i \in I$, we thus have an object (x, h_i) of $\mathcal{C}_{/\Phi(x)}$. Consider an object (x', g) of $\mathcal{C}_{/\Phi(x)}$. To prove the proposition, it is enough to prove that there is a unique $i \in I$ and a unique morphism $(x', g) \rightarrow (x, h_i)$ of $\mathcal{C}_{/\Phi(x)}$.

By definition, g is a morphism $\Phi(x') \rightarrow \Phi(x)$ in \mathcal{D} . By assumption, we can factor g as $h\Phi(f)$ for some $h \in \text{Aut}(\Phi(x'))$ and some $f \in \text{Hom}_{\mathcal{C}}(x', x)$. Moreover, this factorization is unique up to the action of $\text{Aut}(x)$. It follows that there is a unique factorization of the form $h_i\Phi(f)$. The morphism f now furnishes a map $(x', g) \rightarrow (x, h_i)$ in $\mathcal{C}_{/\Phi(x)}$. It is clear from the discussion that this is the unique i for which there is such a morphism, and that f is the unique such morphism. \square

Left Kan extensions can be used to construct principal projectives, as follows. Let $x \in \mathcal{C}$, let pt be the point category (one object, one morphism), and let $i_x: \text{pt} \rightarrow \mathcal{C}$ be the functor taking the object of pt to x . Regarding \mathbf{k} as a pt -module, we have $(i_x)_!(\mathbf{k}) = P_x$. Indeed, if M is a \mathcal{C} -module, then by definition

$$\text{Hom}_{\text{Rep}_{\mathbf{k}}(\mathcal{C})}((i_x)_!(\mathbf{k}), M) = \text{Hom}_{\text{Rep}_{\mathbf{k}}(\text{pt})}(\mathbf{k}, i_x^*(M)) = M_x,$$

and thus $(i_x)_!(\mathbf{k})$ represents the same functor as P_x .

Return now to the setting of a functor $\Phi: \mathcal{C} \rightarrow \mathcal{D}$. Put $y = \Phi(x)$. Then $\Phi \circ i_x = i_y$, so

$$(2.10) \quad P_y = (i_y)_!(\mathbf{k}) = \Phi_!((i_x)_!(\mathbf{k})) = \Phi_!(P_x).$$

We thus see that the left Kan extension takes principal projectives to principal projectives. Since $\Phi_!$ is right exact, it follows from this that $\Phi_!$ takes finitely generated \mathcal{C} -modules to finitely generated \mathcal{D} -modules.

2.3. \mathcal{C} -groups and their representations. Let \mathcal{C} be a category. A \mathcal{C} -group is a functor from \mathcal{C} to the category of groups. Fix a \mathcal{C} -group \mathbf{G} . A \mathbf{G} -module over \mathbf{k} is a \mathcal{C} -module M equipped with a \mathbf{k} -linear action of \mathbf{G}_x on M_x for all $x \in \mathcal{C}$, such that for all morphisms $f: x \rightarrow y$ in \mathcal{C} the induced morphism $f_*: M_x \rightarrow M_y$ is compatible with the actions via the induced homomorphism $f_*: \mathbf{G}_x \rightarrow \mathbf{G}_y$. In other words, for $m \in M_x$ and $g \in \mathbf{G}_x$ we have $f_*(gm) = f_*(g)f_*(m)$. The category $\text{Rep}_{\mathbf{k}}(\mathbf{G})$ of \mathbf{G} -modules is a Grothendieck abelian category.

Let M be a \mathbf{G} -module. For $x \in \mathcal{C}$, let $\mathbf{H}_i(\mathbf{G}, M)_x$ be the group homology $\mathbf{H}_i(\mathbf{G}_x, M_x)$. If $f: x \rightarrow y$ is a morphism in \mathcal{C} , then the induced morphisms $f_*: \mathbf{G}_x \rightarrow \mathbf{G}_y$ and $f_*: M_x \rightarrow M_y$ together induce a morphism $f_*: \mathbf{H}_i(\mathbf{G}, M)_x \rightarrow \mathbf{H}_i(\mathbf{G}, M)_y$. This yields a \mathcal{C} -module structure on $\mathbf{H}_i(\mathbf{G}, M)$. If \mathbf{k} is a commutative ring, then we will denote by $\underline{\mathbf{k}}$ the constant \mathcal{C} -module defined via the formula $\underline{\mathbf{k}}_x = \mathbf{k}$. We then have $\mathbf{H}_i(\mathbf{G}, \underline{\mathbf{k}})_x = \mathbf{H}_i(\mathbf{G}_x, \mathbf{k})$.

The following proposition concerns the homology of a semi-direct product of \mathcal{C} -groups.

Proposition 2.11. *Let \mathbf{G} and \mathbf{E} be \mathcal{C} -groups, and let $\pi: \mathbf{G} \rightarrow \mathbf{E}$ and $\iota: \mathbf{E} \rightarrow \mathbf{G}$ be morphisms of \mathcal{C} -groups such that $\pi \circ \iota = \text{id}$. Let $\mathbf{K} = \ker(\pi)$, which is also a \mathcal{C} -group. Then we have the following:*

- (1) $\mathbf{H}_i(\mathbf{K}, \underline{\mathbf{k}})$ is naturally an \mathbf{E} -module.
- (2) As a \mathcal{C} -module, $\mathbf{H}_i(\mathbf{E}, \underline{\mathbf{k}})$ is a direct summand of $\mathbf{H}_i(\mathbf{G}, \underline{\mathbf{k}})$ via ι_* and π_* .
- (3) Write $\mathbf{H}_r(\mathbf{G}, \underline{\mathbf{k}}) = \mathbf{H}_r(\mathbf{E}, \underline{\mathbf{k}}) \oplus M$ as in (2). Then M admits a \mathcal{C} -module filtration where the graded pieces are subquotients of $\mathbf{H}_i(\mathbf{E}, \mathbf{H}_{r-i}(\mathbf{K}, \underline{\mathbf{k}}))$ with $0 \leq i \leq r - 1$.

Proof. (1) The conjugation action of \mathbf{G} on \mathbf{K} is \mathcal{C} -linear. On homology, \mathbf{K} acts trivially, and hence this action descends to give an \mathbf{E} -module structure on $\mathbf{H}_i(\mathbf{K}, \underline{\mathbf{k}})$.

(2) This is clear.

(3) For $x \in \mathcal{C}$ we have a short exact sequence of groups $1 \rightarrow \mathbf{K}_x \rightarrow \mathbf{G}_x \rightarrow \mathbf{E}_x \rightarrow 1$, which gives a Hochschild–Serre spectral sequence

$$E_{p,q}^2 = \mathbf{H}_p(\mathbf{E}_x, \mathbf{H}_q(\mathbf{K}_x, \mathbf{k})) \implies \mathbf{H}_{p+q}(\mathbf{G}_x, \mathbf{k}).$$

The spectral sequence is functorial in x , and so we get a spectral sequence of \mathcal{C} -modules

$$E_{p,q}^2 = \mathbf{H}_p(\mathbf{E}, \mathbf{H}_q(\mathbf{K}, \underline{\mathbf{k}})) \implies \mathbf{H}_{p+q}(\mathbf{G}, \underline{\mathbf{k}}).$$

In particular, $\mathbf{H}_r(\mathbf{G}, \underline{\mathbf{k}})$ has a filtration by subquotients of the terms $E_{i,r-i}^2$. The edge map $\mathbf{H}_r(\mathbf{G}, \underline{\mathbf{k}}) \rightarrow \mathbf{H}_r(\mathbf{E}, \mathbf{H}_0(\mathbf{K}, \underline{\mathbf{k}}))$ coincides with the map on \mathbf{H}_r induced by π (see [W, §6.8.2]) which we know is a split surjection, so the kernel M has a filtration by subquotients of $E_{i,r-i}^2$ for $0 \leq i \leq r - 1$. \square

3. THE CATEGORY \mathbf{OI} AND VARIANTS

3.1. Definitions and first results. Let \mathbf{OI} be the category whose objects are finite totally ordered sets and whose morphisms are order-preserving injections. For a non-negative integer d , we define a variant $\mathbf{OI}(d)$ as follows. An object of $\mathbf{OI}(d)$ is a pair (S, λ) where S is a totally ordered set and $\lambda = (\lambda_1 < \dots < \lambda_d)$ is an increasing d -tuple in S . A morphism $(S, \lambda) \rightarrow (T, \mu)$ in $\mathbf{OI}(d)$ is an order-preserving injection $f: S \rightarrow T$ satisfying $f(\lambda) = \mu$. Note that $\mathbf{OI} = \mathbf{OI}(0)$. There is a functor $\Phi: \mathbf{OI}(d) \rightarrow \mathbf{OI}$ given by $\Phi(S, \lambda) = S$. We will continue to use the notation Φ for this functor throughout the paper (and use it for all values of d).

Remark 3.1. We introduce $\mathbf{OI}(d)$ to help us study an analogous category $\mathbf{OVI}(R, d)$, the motivation for which is discussed in Remark 4.1 below. \square

Recall that $[n]$ denotes the ordered set $\{1, \dots, n\}$. Given an \mathbf{OI} -module M , we will write M_n for $M_{[n]}$. The category \mathbf{OI} is equivalent to its full subcategory spanned by the $[n]$, so the data of an \mathbf{OI} -module M is equivalent to the data of the M_n together with the maps $f_*: M_n \rightarrow M_m$ induced by the order preserving maps $f: [n] \rightarrow [m]$. Similarly, if M is an $\mathbf{OI}(d)$ -module and λ is an increasing d -tuple in $[n]$, then we will write $M_{n,\lambda}$ for $M_{([n],\lambda)}$.

Proposition 3.2. *There is an equivalence of categories $\mathbf{OI}(d) = \mathbf{OI}^{d+1}$.*

Proof. Let (S, λ) be an object of $\mathbf{OI}(d)$. For $1 \leq i \leq d+1$, let S_i be the set of elements $x \in S$ such that $\lambda_{i-1} < x < \lambda_i$ where, by convention, $\lambda_0 < x < \lambda_{d+1}$ for all x . One easily verifies that $(S, \lambda) \mapsto (S_1, \dots, S_{d+1})$ is an equivalence. \square

Corollary 3.3. *The category $\mathbf{OI}(d)$ is Gröbner. In particular, the category of $\mathbf{OI}(d)$ -modules is locally noetherian.*

Proof. By [SaSn, Theorem 7.1.2] the category \mathbf{OI} is Gröbner, and by [SaSn, Proposition 4.3.5] a finite product of Gröbner categories is Gröbner, so by Proposition 3.2 the category $\mathbf{OI}(d)$ is Gröbner. The assertion about finitely generated $\mathbf{OI}(d)$ -modules now follows from Theorem 2.8. \square

Corollary 3.4. *The category $\mathbf{OI}(d)$ satisfies Property (F). In particular, the tensor product of finitely generated $\mathbf{OI}(d)$ -modules is a finitely generated $\mathbf{OI}(d)$ -module and the tensor product of \mathbf{OI} -modules that are generated in finite degree is also generated in finite degree.*

Proof. The category \mathbf{OI} satisfies Property (F): this can be proved similarly to [SaSn, Proposition 7.3.1]. One easily sees that a finite product of categories satisfying Property (F) again satisfies Property (F), which combined with Proposition 3.2 yields the fact that $\mathbf{OI}(d)$ satisfies Property (F). The assertion about tensor products of finitely generated $\mathbf{OI}(d)$ -modules now follows from Proposition 2.5, and the assertion about tensor products of \mathbf{OI} -modules that are generated in finite degree follows from Proposition 2.6. \square

Finally, we state a result about the growth of finitely generated \mathbf{OI} -modules over fields.

Proposition 3.5. *Let M be a finitely generated \mathbf{OI} -module over a field \mathbf{k} . Then the function $n \mapsto \dim_{\mathbf{k}} M_n$ is a polynomial function for $n \gg 0$.*

Proof. By [SaSn, Theorem 7.1.2], \mathbf{OI} is an ‘‘O-lingual category’’, and by [SaSn, Theorem 6.3.2], this implies the polynomiality statement. \square

3.2. Kan extension. We now study left Kan extensions along the functor $\Phi: \mathbf{OI}(d) \rightarrow \mathbf{OI}$.

Proposition 3.6. *Let M be an $\mathbf{OI}(d)$ -module. Then $\Phi_!(M)_n = \bigoplus_{\lambda} M_{n,\lambda}$, where the sum is taken over all increasing d -tuples λ in $[n]$.*

Proof. By §2.2, we see that $\Phi_!(M)_n$ is $\text{colim}(M|_{\mathbf{OI}(d)_{/[n]}})$. The category $\mathbf{OI}(d)_{/[n]}$ can be viewed as consisting of triples (S, μ, f) , where $(S, \mu) \in \mathbf{OI}(d)$ and $f: S \rightarrow [n]$ is a morphism in \mathbf{OI} . For an increasing d -tuple λ in $[n]$, let $\mathbf{OI}(d)_{/[n],\lambda}$ be the full subcategory of $\mathbf{OI}(d)_{/[n]}$ spanned by triples (S, μ, f) such that f takes μ to λ . Then $\mathbf{OI}(d)_{/[n]}$ is the disjoint union of its subcategories $\mathbf{OI}(d)_{/[n],\lambda}$. Furthermore, $([n], \lambda, \text{id})$ is the final object of $\mathbf{OI}(d)_{/[n],\lambda}$. The result now follows. \square

Corollary 3.7. *The functor $\Phi_!$ is exact.*

3.3. Shift functors. Fix a functorial coproduct \amalg on the category of finite sets. For finite sets S and T , we view $S \amalg T$ as the disjoint union of S and T ; of course, this requires care when S and T share elements. Consider the functor $\Sigma_0: \mathbf{OI}(d) \rightarrow \mathbf{OI}(d)$ given by $\Sigma_0(S, \lambda) = (S \amalg \{\infty\}, \lambda)$, where $S \amalg \{\infty\}$ is given a total order by setting $x < \infty$ for all $x \in S$. Given an $\mathbf{OI}(d)$ -module M , we define the **shift** of M , denoted $\Sigma(M)$, to be $\Sigma_0^*(M)$. There is a map $(S, \lambda) \rightarrow (S \amalg \{\infty\}, \lambda)$ in $\mathbf{OI}(d)$ induced by the inclusion $S \hookrightarrow S \amalg \{\infty\}$. This map induces a map $M \rightarrow \Sigma(M)$ of $\mathbf{OI}(d)$ -modules. We let $\overline{\Sigma}(M)$ denote the cokernel of this map. We call it the **reduced shift** of M . This has the following nice property:

Proposition 3.8. *Suppose that M is an \mathbf{OI} -module such that M_0 is a finitely generated \mathbf{k} -module and $\overline{\Sigma}(M)$ is a finitely generated \mathbf{OI} -module. Then M is a finitely generated \mathbf{OI} -module.*

Proof. By assumption, we can find x_1, \dots, x_m with $x_i \in M_{n_i}$ such that the following holds. Let $\bar{x}_i \in \Sigma(M)_{n_i-1} \cong M_{n_i}$ be the associated element. Then the images of $\{\bar{x}_1, \dots, \bar{x}_m\}$ in $\overline{\Sigma}(M)$ generate $\overline{\Sigma}(M)$. We claim that $\{x_1, \dots, x_m\}$ together with a spanning set of M_0 is a generating set for M . Consider $y \in M_n$ for some $n \geq 0$. We must show that y is in the span of the indicated elements. We will do this by induction on n . The base case $n = 0$ being trivial, we can assume that $n \geq 1$. Let $\bar{y} \in \Sigma(M)_{n-1} \cong M_n$ be the associated element. The image of \bar{y} in $\overline{\Sigma}(M)_n$ is in the span of the images of $\{\bar{x}_1, \dots, \bar{x}_m\}$. It follows that we can write $y = y' + y''$, where y' is in the span of $\{x_1, \dots, x_m\}$ and y'' is in the image of the composition $M_{n-1} \rightarrow \Sigma(M)_{n-1} \cong M_n$. By induction, y'' is in the span of $\{x_1, \dots, x_m\}$ together with a spanning set of M_0 , so y is as well. \square

There is a similar functor $\Delta_0: \mathbf{OI}(d-1) \rightarrow \mathbf{OI}(d)$ defined by $\Delta_0(S, \lambda) = (S \amalg \{\infty\}, \lambda')$, where λ' is obtained by appending ∞ to the end of λ . For an $\mathbf{OI}(d)$ -module M , we let $\Delta(M) = \Delta_0^*(M)$, which is an $\mathbf{OI}(d-1)$ -module. For $d = 0$, we put $\Delta(M) = 0$ by convention.

The following result shows how the shift functor interacts with the Kan extension along the functor $\Phi: \mathbf{OI}(d) \rightarrow \mathbf{OI}$.

Proposition 3.9. *Let M be an $\mathbf{OI}(d)$ -module. Then there is a natural isomorphism*

$$\Sigma(\Phi_!(M)) \cong \Phi_!(\Sigma(M)) \oplus \Phi_!(\Delta(M)).$$

Moreover, if $\alpha: \Phi_!(M) \rightarrow \Sigma(\Phi_!(M))$ and $\beta: M \rightarrow \Sigma(M)$ denote the natural maps, then the diagram

$$\begin{array}{ccc} & \Phi_!(M) & \\ \alpha \swarrow & & \searrow \Phi_!(\beta) \oplus 0 \\ \Sigma(\Phi_!(M)) & \xrightarrow{\cong} & \Phi_!(\Sigma(M)) \oplus \Phi_!(\Delta(M)) \end{array}$$

commutes. In particular, we have a natural isomorphism

$$\overline{\Sigma}(\Phi_!(M)) = \Phi_!(\overline{\Sigma}(M)) \oplus \Phi_!(\Delta(M)).$$

Proof. Using Proposition 3.6, we have

$$\Sigma(\Phi_!(M))_n \cong \bigoplus_{\lambda} M_{n+1, \lambda},$$

where the sum is over all increasing d -tuples λ in $[n+1]$. Similarly, we have

$$\Phi_!(\Sigma(M))_n \cong \bigoplus_{\lambda} M_{n+1, \lambda},$$

where the sum is over all increasing d -tuples λ in $[n]$. Finally, using the obvious analogue of Proposition 3.6 for Δ we have

$$\Phi_!(\Delta(M))_n \cong \bigoplus_{\lambda} M_{n+1,\lambda},$$

where the sum is over all increasing d -tuples λ in $[n]$ that end in $n+1$. Combining these isomorphisms, we obtain an identification

$$\Sigma(\Phi_!(M))_n \cong \Phi_!(\Sigma(M))_n \oplus \Phi_!(\Delta(M))_n.$$

It is clear that this identification comes from an isomorphism of **OI**-modules. The rest of the proposition follows easily. \square

4. THE CATEGORY **OVI** AND ITS VARIANTS

4.1. Definitions. Fix a ring R (always assumed to be associative and unital, though not necessarily commutative). Define **OVI**(R) to be the following category. The objects are **ordered free R -modules**, that is, pairs $(V, \{v_i\}_{i \in I})$ where V is a finite rank free left R -module and $\{v_i\}$ is a basis indexed by a totally ordered set I . The morphisms $(V, \{v_i\}_{i \in I}) \rightarrow (W, \{w_j\}_{j \in J})$ are pairs (f, f_0) , where $f: V \rightarrow W$ is a linear map and $f_0: I \rightarrow J$ is an order-preserving injection, such that $f(v_i) = w_{f_0(i)} + \sum_{j < f_0(i)} a_{i,j} w_j$ for scalars $a_{i,j}$. In words, f takes the i th basis vector of V to the $f_0(i)$ th basis vector of W up to “lower order” terms. We note that f_0 can be recovered from f , so it is often omitted. Furthermore, f is necessarily a split injection. If the ring R is clear, we will just write **OVI**.

For a non-negative integer n , we regard R^n as an ordered free module by endowing it with the standard basis. Every object of **OVI** is isomorphic to R^n for a unique n . For an **OVI**-module M , we write M_n for its value on R^n . The automorphism group of R^n in **OVI** is $U_n(R)$, which we denote simply by U_n in this section. It is the subgroup of $\mathbf{GL}_n(R)$ consisting of upper unitriangular matrices.

Let d be a non-negative integer. We define a variant **OVI**(R, d) = **OVI**(d) as follows. An object is a tuple $(V, \{v_i\}_{i \in I}, \lambda)$ where $(V, \{v_i\}_{i \in I})$ is an ordered free module and λ is an increasing d -tuple in I . A morphism $(V, \{v_i\}_{i \in I}, \lambda) \rightarrow (W, \{w_j\}_{j \in J}, \mu)$ is a morphism $(f, f_0): (V, \{v_i\}) \rightarrow (W, \{w_j\})$ in **OVI** such that $f_0(\lambda) = \mu$ and such that $f(v_i) = w_{f_0(i)}$ for all i appearing in λ (i.e., no lower terms are allowed on marked basis vectors).

For a tuple $\lambda = (1 \leq \lambda_1 < \dots < \lambda_d \leq n)$ we have an object (R^n, λ) of **OVI**(d). Every object of **OVI**(d) is isomorphic to a unique (R^n, λ) . For an **OVI**(d)-module M , we write $M_{n,\lambda}$ for its value on (R^n, λ) . We let $U_{n,\lambda}$ be the automorphism group of (R^n, λ) in **OVI**(d). It is the subgroup of U_n fixing the basis vectors e_{λ_i} for $1 \leq i \leq d$.

Remark 4.1. We introduce **OVI**(d) as a technical device for proving Theorem 1.3, which concerns the homology groups $\mathbf{H}_i(\mathbf{U}, M)$ for **OVI**(R)-modules M . We will see in Corollary 6.5 that the homology of the principal projective **OVI** module at d can be understood in terms of the homology of the trivial **OVI**(d)-module, a helpful simplification. \square

There are several functors to mention:

- There is a functor **OI** \rightarrow **OVI** taking a totally ordered set S to the ordered free module $R[S]$ with basis S . There is a similar functor **OI**(d) \rightarrow **OVI**(d).
- There is a functor **OVI** \rightarrow **OI** taking an ordered free module $(V, \{v_i\}_{i \in I})$ to the totally ordered set I and a morphism (f, f_0) to f_0 . There is a similar functor **OVI**(d) \rightarrow **OI**(d).

- There is a functor $\Psi: \mathbf{OVI}(d) \rightarrow \mathbf{OVI}$ given by forgetting λ . We continue to use the notation Ψ for this functor throughout the paper.

We have the following basic fact that follows from interpreting left multiplication by a matrix as a sequence of row operations.

Proposition 4.2. *Every morphism $\varphi: (R^n, \lambda) \rightarrow (R^m, \mu)$ in $\mathbf{OI}(d)$ has a unique factorization $\varphi = \psi f$ where $\psi \in \text{Aut}(R^m, \mu)$ and f is in the image of the functor $\mathbf{OI}(d) \rightarrow \mathbf{OVI}(d)$.*

4.2. The case where R is finite. The purpose of this section is to prove the following fundamental result:

Theorem 4.3. *If $|R| < \infty$, then the category \mathbf{OVI} is quasi-Gröbner. In particular, by Theorem 2.8 the category $\text{Rep}_{\mathbf{k}}(\mathbf{OVI})$ is locally noetherian when \mathbf{k} is noetherian.*

Proof. An **ordered surjection** $f: S \rightarrow T$ of totally ordered finite sets is a surjection such that for all $i < j$ in T we have $\min f^{-1}(i) < \min f^{-1}(j)$. We let \mathbf{OS} be the category whose objects are finite totally ordered sets and whose morphisms are ordered surjections. This category is known to be Gröbner [SaSn, Theorem 8.1.1]. Given a totally ordered set S , we will regard the dual $R[S]^* = \text{Hom}_R(R[S], R)$ as an element of \mathbf{OVI} as follows. Let $S^* \subset R[S]^*$ be the dual basis to the basis S , and for $s \in S$, write $s^* \in S^*$ for the dual element. Then we order S^* via the rule

$$(4.3.a) \quad s_1^* < s_2^* \quad \text{when} \quad s_2 < s_1.$$

Using this convention, there is a functor $\mathbf{OS}^{\text{op}} \rightarrow \mathbf{OVI}$ taking a totally ordered set S to $R[S]^*$ and an ordered surjection $T \rightarrow S$ to the dual of the induced surjective linear map $R[T] \rightarrow R[S]$. We will show that this functor satisfies property (F), which will complete the proof.

Let V be an object of \mathbf{OVI} . Let $T_1, \dots, T_n \in \mathbf{OS}$ be objects and $f_i: V \rightarrow R[T_i]^*$ be \mathbf{OVI} -morphisms such that the f_i are an enumeration of all possible morphisms satisfying the following condition:

- The set T_i is a total ordering of a finite subset of V^* that spans V^* and $f_i: V \rightarrow R[T_i]^*$ is an \mathbf{OVI} -morphism that is dual to the natural surjection $R[T_i] \rightarrow V$.

Since V is finite, there are only finitely many such f_i . Now consider some $S \in \mathbf{OS}$ and an \mathbf{OVI} -morphism $f: V \rightarrow R[S]^*$. To prove that our functor satisfies property (F), it is enough to prove that for some $1 \leq i \leq n$ we can write $f = g \circ f_i$, where $g: R[T_i]^* \rightarrow R[S]^*$ is dual to an \mathbf{OS} -morphism $S \rightarrow T_i$. Let $T \subset V^*$ be the image of S under the dual surjection $f^*: R[S] \rightarrow V^*$. Let $h: S \rightarrow T$ be the resulting surjection. Order T via the rule

$$(4.3.b) \quad t_1 < t_2 \quad \text{when} \quad \min h^{-1}(t_1) < \min h^{-1}(t_2),$$

which makes h an \mathbf{OS} -morphism. Combining (4.3.b) with (4.3.a) (applied to order both S^* and T^*), we see that T^* has the ordering

$$(4.3.c) \quad t_1^* < t_2^* \quad \text{when} \quad \max\{s^* \mid s \in h^{-1}(t_1)\} < \max\{s^* \mid s \in h^{-1}(t_2)\};$$

Let $g: R[T]^* \rightarrow R[S]^*$ be the \mathbf{OVI} -morphism dual to h , so

$$(4.3.d) \quad g(t^*) = \sum_{s \in h^{-1}(t)} s^* \quad (t \in T).$$

Finally, let $F: V \rightarrow R[T]^*$ be the injection dual to the surjection $R[T] \rightarrow V^*$ induced by the inclusion $T \hookrightarrow V^*$, so $f = g \circ F$. The fact that f is an \mathbf{OVI} -morphism together with (4.3.c)

and (4.3.d) implies that F is an **OVI**-morphism. This implies that for some $1 \leq i \leq n$ we have $T = T_i$ and $F = f_i$, and we are done. \square

Remark 4.4. By making use of a variant **OS**(d) of **OS**, one can prove a version of the above theorem for **OVI**(d). Since we do not need this, we omit the details. \square

4.3. Kan extension. We now study left Kan extensions along the functor $\Psi: \mathbf{OVI}(d) \rightarrow \mathbf{OVI}$.

Proposition 4.5. *Let M be an **OVI**(d)-module. Then*

$$\Psi_!(M)_n = \bigoplus_{\lambda} \text{Ind}_{U_{n,\lambda}}^{U_n}(M_{n,\lambda}),$$

the sum taken over all increasing sequences $1 \leq \lambda_1 < \dots < \lambda_d \leq n$.

Proof. Let $\mathbf{OVI}(d)'$ be the category whose objects are those of $\mathbf{OVI}(d)$ and where a morphism $(V, \{v_i\}_{i \in I}, \lambda) \rightarrow (W, \{w_j\}_{j \in J}, \mu)$ is a morphism (f, f_0) as in **OVI** (ignoring the λ and μ) such that f_0 is a morphism in **OI**(d). The automorphism groups in $\mathbf{OVI}(d)'$ are the U_n 's. The functor Ψ factors as $\Psi_2 \circ \Psi_1$, where $\Psi_1: \mathbf{OVI}(d) \rightarrow \mathbf{OVI}(d)'$ and $\Psi_2: \mathbf{OVI}(d)' \rightarrow \mathbf{OVI}$ are the natural functors. Proposition 2.9 applies to the functor Ψ_1 , and so we find

$$(\Psi_1)_!(M)_{n,\lambda} = \text{Ind}_{U_{n,\lambda}}^{U_n}(M_{n,\lambda}).$$

Arguing exactly as in the proof of Proposition 3.6, we find

$$(\Psi_2)_!(N)_n = \bigoplus_{\lambda} N_{n,\lambda}$$

for any $\mathbf{OVI}(d)'$ -module N . The result follows. \square

4.4. OVI-modules and representations of \mathbf{U} . Define an **OI**(d)-group \mathbf{U}_d by $(\mathbf{U}_d)_{n,\lambda} = U_{n,\lambda}$. If M is an **OVI**(d)-module then we can regard it as an **OI**(d)-module via the functor $\mathbf{OI}(d) \rightarrow \mathbf{OVI}(d)$, and as such it has the structure of a \mathbf{U}_d -module. We thus have a functor

$$\{\mathbf{OVI}(d)\text{-modules}\} \rightarrow \{\mathbf{U}_d\text{-modules}\}.$$

One can show that the above functor is fully faithful. We do not need this result, so we do not include a proof. We write \mathbf{U} in place of \mathbf{U}_0 .

5. NOETHERIANITY OF **OVI**-MODULES

The goal of this section is to prove Theorem 1.4, which we recall says that if R is a ring whose underlying additive group is finitely generated and \mathbf{k} is a commutative noetherian ring, then the category of **OVI**(R)-modules over \mathbf{k} is locally noetherian, that is, any submodule of a finitely generated module is finitely generated. The ring R here is not required to be commutative. When R is finite, this follows from the much easier Theorem 4.3. We will also prove a converse to this result that says that (ignoring degenerate cases) the category $\text{Rep}_{\mathbf{k}}(\mathbf{OVI}(R))$ is locally noetherian only if \mathbf{k} is noetherian and the additive group of R is finitely generated. We thus have a complete characterization of when $\text{Rep}_{\mathbf{k}}(\mathbf{OVI}(R))$ is locally noetherian.

This section has four subsections. We begin in §5.1 by describing a toy version of our proof. We then prove a technical ring-theoretic result in §5.2. The proof of Theorem 1.4 is in the long §5.3. Finally, in §5.4 we prove the aforementioned converse to Theorem 1.4.

5.1. **A toy version of Theorem 1.4.** In the next sections, we prove Theorem 1.4. The proof is a bit lengthy and heavy on notation, but the idea behind it is not too complicated. In this section we sketch the proof of a simpler result that illustrates the main ideas.

Theorem 1.4 (with $R = \mathbf{Z}$) implies that the group algebra $\mathbf{k}[U_n(\mathbf{Z})]$ is left-noetherian, provided \mathbf{k} is noetherian. Let us try to prove this for $n = 3$. The group algebra can be identified, as a \mathbf{k} -module, with

$$Q = x_2 y_3 \mathbf{k}[x_1^{\pm 1}, y_1^{\pm 1}, y_2^{\pm 1}],$$

which we treat as a \mathbf{k} -submodule of the Laurent polynomial ring in the five variables. The monomials in this module correspond to the group elements in $\mathbf{k}[U_3(\mathbf{Z})]$; the exponents of the x 's give the second column, while the exponents of the y 's gives the third.

We must show that any $U_3(\mathbf{Z})$ -submodule of Q is finitely generated. Let M be a given submodule. Let Q_+ be the \mathbf{k} -submodule of Q where only positive powers of the variables appear. We would like to associate to M a monomial ideal in Q_+ , and then use the noetherianity of monomial ideals to conclude that M is finitely generated. By "ideal" here we really mean $\mathbf{k}[x_1, y_1, y_2]$ -submodule. The obvious attempt at this is to first form $M_+ = M \cap Q_+$ and then take its initial module $\text{in}(M_+)$, the \mathbf{k} -span of the initial terms of its elements under some monomial order. The problem with this is that $\text{in}(M_+)$ need not be an ideal. For example, suppose that M_+ contains the element $f = x_2 y_3 (y_2 + 1)$, with initial term $\text{in}(f) = x_2 y_2 y_3$. Let's try to find $x_1 \text{in}(f)$ in $\text{in}(M_+)$. If we apply the matrix

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

to f , we get the element $f' = x_1 x_2 y_3 (y_1 y_2 + 1)$, with initial term $x_1 x_2 y_1 y_2 y_3$. This is equal to $x_1 y_1 \text{in}(f)$, so we now need to get rid of the y_1 . We therefore apply the matrix

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

to f' , to get the element $f'' = x_1 x_2 y_1^{-1} y_3 (y_1 y_2 + 1)$. This has the correct leading term. However, it no longer belongs to M_+ : the power of y in the non-leading term is negative. Thus $\text{in}(f'')$ does not give an element of $\text{in}(M_+)$. There does not seem to be a way to produce $x_1 \text{in}(f)$ in $\text{in}(M_+)$.

Remark 5.1. This approach is really attempting to show that the monoid algebra $Q_+ = \mathbf{k}[U_3(\mathbf{Z}_{\geq 0})]$ is noetherian. In fact, it is not noetherian. For example, the left ideal generated

by the matrices $\begin{pmatrix} 1 & n & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ for $n \geq 0$ in Q_+ is not finitely generated. \square

To overcome this problem, we take a more subtle approach. Let Q_* be the submodule of Q where the exponent of y_2 is positive, but we still allow negative powers of x_1 and y_1 . Given $M \subset Q$, let $M_* = M \cap Q_*$. We can then form the initial module with respect to y_2 (that is, we treat the other variables as constants); call this $\text{in}_2(M_*)$. Since we allow negative powers of y_1 , the issue in the previous paragraph does not arise, and $\text{in}_2(M_*)$ is closed under multiplication by $x_1^{\pm 1}$, $y_1^{\pm 1}$, and y_2 . We now intersect $\text{in}_2(M_*)$ with M_+ and then take initial terms with respect to x_1 and y_1 . The result is a monomial ideal of Q_+ . Call this monomial

ideal $I(M)$. One can show that if $M \subset M'$ and $I(M) = I(M')$ then $M = M'$. Since Q_+ is noetherian as a $\mathbf{k}[x_1, y_1, y_2]$ -module, this proves that Q is noetherian as a $\mathbf{k}[U_3(\mathbf{Z})]$ -module.

The same approach works for $\mathbf{k}[U_n(\mathbf{Z})]$, but the process is more involved. Let Q be the group algebra, which we identify with a \mathbf{k} -submodule of the Laurent polynomial ring in variables $x_{i,j}$ with $i \leq j$. We let $Q^{(k)}$ be the \mathbf{k} -submodule where the exponents of $x_{i,j}$ with $i \geq k$ are positive. Thus $Q^{(n)} = Q$ and $Q^{(0)}$ is what we would call Q_+ . Let M be a $U_n(\mathbf{Z})$ -submodule of Q . We obtain a monomial ideal in Q_+ as follows: intersect with $Q^{(n-1)}$ and take the initial submodule with respect to $x_{\bullet, n}$; then intersect with $Q^{(n-2)}$ and take the initial submodule with respect to $x_{\bullet, n-1}$; and so on. After n steps we obtain a monomial ideal in Q_+ . The argument then proceeds as in the previous case.

Remark 5.2. The strategy employed here has some parallels with Hall's proof [Ha, Lemma 3] that the group ring $\mathbf{k}[\Gamma]$ of a polycyclic group Γ is noetherian. There the key point is to take a normal subgroup Γ' such that $\Gamma/\Gamma' \cong \mathbf{Z}$ and treat each element of $\mathbf{k}[\Gamma]$ as a Laurent polynomial in x with coefficients in $\mathbf{k}[\Gamma']$ (where x is some generator for \mathbf{Z}) and argue by passing to initial terms. \square

The proof for $\mathbf{OVI}(R)$ differs from the above in only two respects. First, there is a great deal of additional bookkeeping. Second, we need a noetherianity result for the kind of \mathbf{OI} -monomial ideals that appear in the reduction. This follows easily from Higman's lemma, and is closely related to the theorem (of Cohen [Co] and Aschenbrenner–Hillar–Sullivant [AHi, HilSu]) that $\mathbf{k}[x_i]_{i \in \mathbf{N}}$ is $\text{Inc}(\mathbf{N})$ -noetherian, where $\text{Inc}(\mathbf{N})$ is the monoid of increasing functions $\mathbf{N} \rightarrow \mathbf{N}$.

5.2. Eliminating additive torsion. For technical reasons, Theorem 1.4 is easier to prove when R is a ring whose additive group is a finitely generated free abelian group. In this section, we show how to reduce to that case. Our main tool is the following lemma.

Lemma 5.3. *Let S be a ring and let \mathbf{k} be a commutative ring such that the category of $\mathbf{OVI}(S)$ -modules over \mathbf{k} is locally noetherian. Assume that S surjects onto a ring R . Then the category of $\mathbf{OVI}(R)$ -modules over \mathbf{k} is locally noetherian.*

Proof. The surjection $S \rightarrow R$ induces a functor $\Phi: \mathbf{OVI}(S) \rightarrow \mathbf{OVI}(R)$. By Proposition 2.4, it is enough to show that Φ satisfies property (F). For some $d \geq 1$, let P_d be the principal projective $\mathbf{OVI}(R)$ -module associated to R^d , so

$$(P_d)_n = \mathbf{k}[\text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n)] \quad (n \geq 1).$$

By Proposition 2.3, to prove that Φ satisfies property (F) it is enough to prove that $\Phi^*(P_d)$ is finitely generated. Since the map $S \rightarrow R$ of rings is surjective, the induced map

$$\text{Hom}_{\mathbf{OVI}(S)}(S^d, S^n) \rightarrow \text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n)$$

is also surjective for all $n \geq 1$. This implies that there is a surjective map from the principal projective $\mathbf{OVI}(S)$ -module associated to S^d to $\Phi^*(P_d)$, and thus that $\Phi^*(P_d)$ is finitely generated, as desired. \square

Lemma 5.4. *Let R be a ring whose additive group is finitely generated. Then there exists a ring S and a surjection $S \rightarrow R$ such that the additive group of S is free and finitely generated.*

Proof. Let R_{tor} be the torsion subgroup of the additive group of R and let $N \geq 1$ be the exponent of R_{tor} , i.e., the minimal number such that $NR_{\text{tor}} = 0$. The proof is by induction on N . In the base case where $N = 1$, the group R_{tor} is trivial and there is nothing to prove.

Assume, therefore, that $N > 1$ and that the lemma is true for all smaller exponents. Let p be a prime dividing N . The ring R/pR is a finite ring. Let $\mathbf{Z}[R/pR]$ be the monoid ring of the multiplicative monoid underlying R/pR , so $\mathbf{Z}[R/pR]$ consists of finite sums of formal symbols $\{[x] \mid x \in R/pR\}$ with the ring structure defined by $[x][y] = [xy]$. The additive group of the ring $\mathbf{Z}[R/pR]$ is free abelian with basis in bijection with the elements of R/pR , and there exists a ring surjection $\mathbf{Z}[R/pR] \rightarrow R/pR$ taking $[x] \in \mathbf{Z}[R/pR]$ to $x \in R/pR$. Let R' be the fiber product of the surjections $\mathbf{Z}[R/pR] \rightarrow R/pR$ and $R \rightarrow R/pR$, so we have a cartesian square

$$\begin{array}{ccc} R' & \longrightarrow & \mathbf{Z}[R/pR] \\ \downarrow & & \downarrow \\ R & \longrightarrow & R/pR. \end{array}$$

Concretely,

$$R' = \{(x, r) \in \mathbf{Z}[R/pR] \times R \mid x \text{ and } r \text{ map to same element of } R/pR\}.$$

Since the maps $R \rightarrow R/pR$ and $\mathbf{Z}[R/pR] \rightarrow R/pR$ are surjective, so is the map $R' \rightarrow R$. Since the additive group underlying $\mathbf{Z}[R/pR]$ is torsion-free, the torsion subgroup $(R')_{\text{tor}}$ consists of pairs $(0, r) \in \mathbf{Z}[R/pR] \times R_{\text{tor}}$ such that $r \in R_{\text{tor}}$ maps to 0 in R/pR . It follows that

$$(R')_{\text{tor}} \cong R_{\text{tor}} \cap pR = pR_{\text{tor}}.$$

The exponent of $(R')_{\text{tor}}$ is thus N/p , so by induction there exists a ring S whose additive group is finitely generated and free together with a surjection $S \rightarrow R'$. The desired surjection to R is then the composition $S \rightarrow R' \rightarrow R$. \square

5.3. The proof of Theorem 1.4. We now commence with the proof of Theorem 1.4, which we recall says that if R is a ring whose underlying additive group is finitely generated and \mathbf{k} is a commutative noetherian ring, then the category of $\mathbf{OVI}(R)$ -modules over \mathbf{k} is locally noetherian. By Lemmas 5.3 and 5.4, we can assume that the additive group of R is a finitely generated free abelian group (this assumption will first be used in Substep 2a below). Fix some $d \geq 0$ and let P_d be the principal projective of $\mathbf{OVI}(R)$ defined by the formula

$$(P_d)_n = \mathbf{k}[\text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n)] \quad (n \geq 1).$$

To prove the theorem, it is enough to prove that the poset of $\mathbf{OVI}(R)$ -submodules of P_d is noetherian, i.e., has no infinite strictly increasing sequences. This is trivial for $d = 0$, so we can assume that $d \geq 1$.

Say that a map $f: I \rightarrow J$ of posets is **conservative** if for all $i, i' \in I$ satisfying $i \leq i'$ and $f(i) = f(i')$, we have $i = i'$. If J is a noetherian poset and $f: I \rightarrow J$ is a conservative map, then I is also noetherian. Our strategy will be to use a sequence of conservative poset maps to reduce proving that the poset of $\mathbf{OVI}(R)$ -submodules of P_d is noetherian to proving that another easier poset $\mathfrak{M}^{(0)}$ is noetherian. To help the reader understand its structure, we divide our proof into three steps (each of which is divided into a number of substeps).

Since we will introduce a lot of notation, to help the reader recall the meanings of symbols we will list the notation that is defined in each substep.

Step 1. *We construct a poset \mathfrak{M} and reduce the theorem to showing that \mathfrak{M} is noetherian.*

As in the toy version of our proof, the first step will be to relate the poset of $\mathbf{OVI}(R)$ -submodules of P_d to a poset \mathfrak{M} constructed using certain “generalized polynomial rings”. In fact, \mathfrak{M} will be a poset of certain special $\mathbf{OI}(d)$ -submodules of an $\mathbf{OI}(d)$ -module Q . There are three substeps: in Substep 1a we construct the $\mathbf{OI}(d)$ -module Q , in Substep 1b we construct the poset \mathfrak{M} of special $\mathbf{OI}(d)$ -submodules of Q , and then finally in Substep 1c we construct a conservative poset map from the poset of $\mathbf{OVI}(R)$ -submodules of P_d to \mathfrak{M} .

Substep 1a. *We construct the $\mathbf{OI}(d)$ -module Q .*

Notation defined: $\Lambda_n, T_{i,j}^r, \mathbf{T}_n, \Lambda_n(\mathbf{S}), \Lambda_{n,\alpha}, \mathbf{T}_{n,\alpha}, Q, Q_{n,\alpha}$

We will want to view matrices with entries in R as certain kinds of “monomials”. Since we will be focusing on P_d , the relevant matrices will have d columns and some number $n \geq 1$ of rows. To that end, we make the following definition:

- Define Λ_n to be the commutative monoid generated by the set of formal symbols $T_{i,j}^r$ with $1 \leq i \leq n$ and $1 \leq j \leq d$ and $r \in R$ subject to the relations $T_{i,j}^{r_1} T_{i,j}^{r_2} = T_{i,j}^{r_1+r_2}$, where $1 \leq i \leq n$ and $1 \leq j \leq d$ and $r_1, r_2 \in R$.

Elements of Λ_n are thus “monomials” in the $T_{i,j}^r$, and are naturally in bijection with $n \times d$ matrices with entries in R : given such a matrix $(r_{i,j})$, the associated element of Λ_n is the product of the $T_{i,j}^{r_{i,j}}$, where i ranges over $1 \leq i \leq n$ and j ranges over $1 \leq j \leq d$. The monoid product in Λ_n corresponds to matrix addition. For later use, setting $\mathbf{T}_n = \{T_{i,j} \mid 1 \leq i \leq n, 1 \leq j \leq d\}$, for $\mathbf{S} \subset \mathbf{T}_n$ we define $\Lambda_n(\mathbf{S})$ to be the submonoid of Λ_n generated by $\{T_{i,j}^r \mid T_{i,j} \in \mathbf{S}, r \in R\}$.

Now consider an element $f \in \text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n)$. By definition, f is a linear map $R^d \rightarrow R^n$ such that there exists a strictly increasing sequence $\alpha = (\alpha_1, \dots, \alpha_d)$ of d elements of $[n] = \{1, \dots, n\}$ with the following property:

- For $1 \leq i \leq d$, the map f takes the i th basis element of R^d to the sum of the α_i th basis element of R^n and an R -linear combination of the basis elements of R^n that occur before α_i .

Define $\Lambda_{n,\alpha}$ to be the subset of Λ_n consisting of elements associated to $n \times d$ matrices of this form. Defining

$$\mathbf{T}_{n,\alpha} = \{T_{i,j} \mid 1 \leq j \leq d, 1 \leq i < \alpha_j\},$$

an element $\tau \in \Lambda_{n,\alpha}$ can be written as

$$(5.5) \quad \tau = T_{\alpha_1,1}^1 T_{\alpha_2,2}^1 \cdots T_{\alpha_d,d}^1 \tau' \quad \text{with } \tau' \in \Lambda_n(\mathbf{T}_{n,\alpha}).$$

We thus have a bijection of sets

$$\text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n) \cong \bigsqcup_{\alpha} \Lambda_{n,\alpha},$$

where the disjoint union ranges over the strictly increasing sequences α of d elements of $[n]$. It follows that

$$(5.6) \quad (P_d)_n = \mathbf{k}[\text{Hom}_{\mathbf{OVI}(R)}(R^d, R^n)] = \bigoplus_{\alpha} \mathbf{k}[\Lambda_{n,\alpha}].$$

The various $\mathbf{k}[\Lambda_{n,\alpha}]$ fit together into an $\mathbf{OI}(d)$ -module Q with

$$Q_{n,\alpha} = \mathbf{k}[\Lambda_{n,\alpha}] \quad ((n, \alpha) \in \mathbf{OI}(d)).$$

Substep 1b. *We construct a poset \mathfrak{M} of $\mathbf{OI}(d)$ -submodules of Q .*

Notation defined: $\mathfrak{M}, E_{i,\alpha_j}^r$

Consider an $\mathbf{OVI}(R)$ -submodule M of P_d . We say that M is a **homogeneous $\mathbf{OVI}(R)$ -submodule of P_d** if for all $n \geq 1$, the \mathbf{k} -submodule M_n of $(P_d)_n$ splits according to the decomposition (5.6), i.e., for all $(n, \alpha) \in \mathbf{OI}(d)$ there exists some \mathbf{k} -submodule $M_{n,\alpha}$ of $\mathbf{k}[\Lambda_{n,\alpha}]$ such that

$$M_n = \bigoplus_{\alpha} M_{n,\alpha}.$$

In this case, the various $M_{n,\alpha}$ fit together into an $\mathbf{OI}(d)$ -submodule of Q . We thus get a poset injection

$$\{\text{homogeneous } \mathbf{OVI}(R)\text{-submodules of } P_d\} \hookrightarrow \{\mathbf{OI}(d)\text{-submodules of } Q\}.$$

The image of this injection consists of all $\mathbf{OI}(d)$ -submodules M of Q such that each $M_{n,\alpha} \subset Q_{n,\alpha}$ is preserved by the action of $U_n(R)$, which acts on $Q_{n,\alpha}$ via the identification of $Q_{n,\alpha}$ with the set of formal \mathbf{k} -linear combinations of appropriate $n \times d$ matrices.

For the sake of our later arguments, we will actually consider a larger collection of submodules. Define \mathfrak{M} to be the poset of all $\mathbf{OI}(d)$ -submodules M of Q such that the following hold. Consider $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$. Let $\{\vec{e}_1, \dots, \vec{e}_n\}$ be the standard basis for R^n . For $1 \leq j \leq d$ and $1 \leq i < \alpha_j$ and $r \in R$, define $E_{i,\alpha_j}^r \in U_n(R)$ to be the element that takes \vec{e}_{α_j} to $r\vec{e}_i + \vec{e}_{\alpha_j}$ and fixes all of the other basis vectors. We then require that $M_{n,\alpha}$ be preserved by all of the E_{i,α_j}^r for $1 \leq j \leq d$ and $1 \leq i < \alpha_j$ and $r \in R$. The construction in the previous paragraph gives a poset injection

$$(5.7) \quad \{\text{homogeneous } \mathbf{OVI}(R)\text{-submodules of } P_d\} \hookrightarrow \mathfrak{M}.$$

Substep 1c. We construct a conservative poset map $\{\mathbf{OVI}(R)\text{-submodules of } P_d\} \longrightarrow \mathfrak{M}$.

Notation defined: none

By (5.7), it is enough to construct a conservative poset map

$$(5.8) \quad \{\mathbf{OVI}(R)\text{-submodules of } P_d\} \rightarrow \{\text{homogeneous } \mathbf{OVI}(R)\text{-submodules of } P_d\}.$$

For each $n \geq 1$, put a total ordering on the set of all strictly increasing sequences α of d elements of $[n]$ using the lexicographic ordering: $\alpha < \alpha'$ if the first nonzero entry $\alpha' - \alpha$ is positive. Given a nonzero element $f \in (P_d)_n$, use the identification (5.6) to write $f = \sum_{\alpha} f_{n,\alpha}$ with $f_{n,\alpha} \in \mathbf{k}[\Lambda_{n,\alpha}]$. Define $\text{in}(f) = f_{n,\alpha_0}$, where α_0 is the largest index such that $f_{n,\alpha_0} \neq 0$.

Given an $\mathbf{OVI}(R)$ -submodule M of P_d and some $n \geq 1$, define $\text{in}(M)_n$ to be the \mathbf{k} -span of $\{\text{in}(f) \mid f \in M_n\}$. It is easy to see that $\text{in}(M)$ is also an $\mathbf{OVI}(R)$ -submodule of P_d . Moreover, by construction $\text{in}(M)$ is homogeneous. The map $M \mapsto \text{in}(M)$ is thus a poset map as in (5.8). We must prove that it is conservative. Assume otherwise, and let M and M' be $\mathbf{OVI}(R)$ -submodules of P_d such that $M \subsetneq M'$ and $\text{in}(M) = \text{in}(M')$. Let $n \geq 1$ be such that $M_n \subsetneq M'_n$. Let $x \in M'_n \setminus M_n$ be such that $\text{in}(x)$ lies in $\mathbf{k}[\Lambda_{n,\alpha}]$ with α as small as possible. Since $\text{in}(M) = \text{in}(M')$, we can find some $x' \in M_n$ with $\text{in}(x) = \text{in}(x')$. But then $x - x' \in M'_n \setminus M_n$, while $\text{in}(x - x')$ lies in $\mathbf{k}[\Lambda_{n,\alpha'}]$ with $\alpha' < \alpha$, a contradiction.

Step 2. We construct a poset $\mathfrak{M}^{(0)}$ and reduce the theorem to showing that $\mathfrak{M}^{(0)}$ is noetherian.

In Step 1, we reduced the theorem to showing that the poset \mathfrak{M} constructed in Substep 1b is noetherian. The goal of this step is to construct a conservative poset map from \mathfrak{M} to a simpler poset $\mathfrak{M}^{(0)}$. This will be done in a sequence of steps. Recall that \mathfrak{M} is a subposet

of the poset of $\mathbf{OI}(d)$ -submodules of an $\mathbf{OI}(d)$ -module Q . In Substep 2a we will construct an $\mathbf{OI}(d)$ -module filtration

$$Q^{(0)} \subset Q^{(1)} \subset \dots \subset Q^{(d)} = Q.$$

Next, in Substeps 2b and 2c we will construct two posets $\mathfrak{M}^{(k)}$ and $\mathfrak{N}^{(k)}$ of special $\mathbf{OI}(d)$ -submodules of $Q^{(k)}$ such that $\mathfrak{M}^{(d)} = \mathfrak{M}$. Finally, in Substeps 2d and 2e we will construct a sequence of conservative poset maps

$$\mathfrak{M} = \mathfrak{M}^{(d)} \rightarrow \mathfrak{N}^{(d-1)} \rightarrow \mathfrak{M}^{(d-1)} \rightarrow \mathfrak{N}^{(d-2)} \rightarrow \dots \rightarrow \mathfrak{N}^{(0)} \rightarrow \mathfrak{M}^{(0)}.$$

This reduces the theorem to showing that the poset $\mathfrak{M}^{(0)}$ is noetherian.

Substep 2a. *We construct an $\mathbf{OI}(d)$ -module filtration*

$$Q^{(0)} \subset Q^{(1)} \subset \dots \subset Q^{(d)} = Q.$$

Notation defined: $(R, +) = (\mathbf{Z}^\lambda, +)$, $R_{\geq 0}$, $\Lambda_{n,\alpha,k+}$, $\Lambda_{n,\alpha,+}$, $\Lambda_{n,+}$, $\Lambda_{n,+}(\mathbf{S})$, $Q^{(k)}$, $Q_{n,\alpha}^{(k)}$

This step is where we use the fact that the additive group of R is a finitely generated free abelian group. Fix an identification of this additive group with \mathbf{Z}^λ for some $\lambda \geq 1$ such that the multiplicative identity $1 \in R$ is identified with an element of $(\mathbf{Z}_{\geq 0})^\lambda$. Let $R_{\geq 0}$ be the submonoid of the additive group of R corresponding to $(\mathbf{Z}_{\geq 0})^\lambda$. The monoid $R_{\geq 0}$ contains $1 \in R$, but is not necessarily closed under multiplication.

Consider $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$. For $0 \leq k \leq d$, define $\Lambda_{n,\alpha,k+}$ to be the set of all $\tau \in \Lambda_{n,\alpha}$ such that if $T_{i,j}^r$ appears in τ with $i \geq \alpha_k$, then $r \in R_{\geq 0}$. For $k = 0$, we use the convention $\alpha_0 = 0$, and we will also frequently omit the k , so $\Lambda_{n,\alpha,+}$ is the set of all $\tau \in \Lambda_{n,\alpha}$ such that if $T_{i,j}^r$ appears in τ , then $r \in R_{\geq 0}$. We will similarly define $\Lambda_{n,+}$ and $\Lambda_{n,+}(\mathbf{S})$ for $\mathbf{S} \subset \mathbf{T}_n$. We then define $Q^{(k)}$ to be the $\mathbf{OI}(d)$ -submodule of Q where for all $(n, \alpha) \in \mathbf{OI}(d)$, we have

$$Q_{n,\alpha}^{(k)} = \mathbf{k}[\Lambda_{n,\alpha,k+}].$$

We thus have $Q^{(d)} = Q$. Moreover,

$$Q_{n,\alpha}^{(0)} = \mathbf{k}[\Lambda_{n,\alpha,+}].$$

Substep 2b. *For $0 \leq k \leq d$, we construct a subposet $\mathfrak{M}^{(k)}$ of the poset of $\mathbf{OI}(d)$ -submodules of $Q^{(k)}$ such that $\mathfrak{M}^{(d)} = \mathfrak{M}$.*

Notation defined: $\mathfrak{M}^{(k)}$, $(a.i_k)$, $(a.ii_k)$, (b_k) , (c_k)

We begin with some terminology. A \mathbf{k} -submodule X of $\mathbf{k}[\Lambda_n]$ is **homogeneous** with respect to $\mathbf{S} \subset \mathbf{T}_n$ if the following holds for all $x \in X$. Write

$$x = \sum_{q=1}^m \tau_q y_q,$$

where for all $1 \leq q \leq m$ we have the following:

- $\tau_q \in \Lambda_n(\mathbf{S})$, and the different τ_q are all distinct.
- $y_q \in \mathbf{k}[\Lambda_n(\mathbf{T}_n \setminus \mathbf{S})]$.

We then require that $\tau_q y_q \in X$ for all $1 \leq q \leq m$.

Now consider some $0 \leq k \leq d$. Define $\mathfrak{M}^{(k)}$ to be the set of all $\mathbf{OI}(d)$ -submodules M of $Q^{(k)}$ such that for all $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$, the following conditions $(a.i_k)$, $(a.ii_k)$, (b_k) , and (c_k) hold. To simplify our notation, we will set $\alpha_0 = 0$.

- (a) The \mathbf{k} -module $M_{n,\alpha} \subset \mathbf{k}[\Lambda_{n,\alpha,k+}]$ is closed under multiplication by the following elements:
- (i_k) $T_{i,j}^r$ with $k \leq j \leq d$ and $1 \leq i < \alpha_k$ and $r \in R$.
 - (ii_k) $T_{i,j}^r$ with $k+1 \leq j \leq d$ and $\alpha_k \leq i < \alpha_j$ and $r \in R_{\geq 0}$.
- (b_k) The \mathbf{k} -module $M_{n,\alpha}$ is closed under the operators E_{i,α_j}^r with $1 \leq j \leq k$ and $1 \leq i < \alpha_j$ and $r \in R$.
- (c_k) The \mathbf{k} -module $M_{n,\alpha} \subset \mathbf{k}[\Lambda_{n,\alpha,k+}]$ is homogeneous with respect to
- $$\{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+1 \leq j \leq d \text{ and } \max(\alpha_k, 1) \leq i < \alpha_j\}.$$

We claim that $\mathfrak{M}^{(d)} = \mathfrak{M}$. Condition (b_d) implies that $\mathfrak{M}^{(d)} \subset \mathfrak{M}$, so we must only prove that $\mathfrak{M} \subset \mathfrak{M}^{(d)}$. Consider $M \in \mathfrak{M}$ and $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$. We must verify that $M_{n,\alpha}$ satisfies the properties above:

- For (a.i_d), we must show that $M_{n,\alpha}$ is closed under multiplication by $T_{i,d}^r$ for $1 \leq i < \alpha_d$ and $r \in R$. But this can be achieved using the operator E_{i,α_d}^r , and by the definition of \mathfrak{M} the \mathbf{k} -module $M_{n,\alpha}$ is closed under this operator, so (a.i_d) follows.
- No pairs (i, j) satisfy the conditions of (a.ii_d), so that condition is trivial.
- Condition (b_d) is a special case of the condition defining \mathfrak{M} , so it follows.
- The set referred to in condition (c_d) consists only of

$$\{T_{\alpha_j,j} \mid 1 \leq j \leq d\},$$

and by definition every element of $\mathbf{k}[\Lambda_{n,\alpha}]$ is homogeneous with respect to these variables (see (5.5)), so that condition follows.

Substep 2c. For $0 \leq k < d$, we construct a subposet $\mathfrak{N}^{(k)}$ of the poset of $\mathbf{OI}(d)$ -submodules of $Q^{(k)}$.

Notation defined: $\mathfrak{N}^{(k)}$, $(a'.i'_k)$, $(a'.ii'_k)$, (b'_k) , (c'_k)

Our definition of $\mathfrak{N}^{(k)}$ will be a slight modification of our definition of $\mathfrak{M}^{(k)}$. Define $\mathfrak{N}^{(k)}$ to be the set of all $\mathbf{OI}(d)$ -submodules N of $Q^{(k)}$ such that for all $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$, the following conditions $(a'.i'_k)$, $(a'.ii'_k)$, (b'_k) , and (c'_k) hold. To simplify our notation, we will set $\alpha_0 = 0$.

- (a') The \mathbf{k} -module $N_{n,\alpha} \subset \mathbf{k}[\Lambda_{n,\alpha,k+}]$ is closed under multiplication by the following elements:
- (i'_k) $T_{i,j}^r$ with $k+1 \leq j \leq d$ and $1 \leq i < \alpha_k$ and $r \in R$.
 - (ii'_k) $T_{i,j}^r$ with $k+1 \leq j \leq d$ and $\alpha_k \leq i < \alpha_j$ and $r \in R_{\geq 0}$.
- (b'_k) The \mathbf{k} -module $N_{n,\alpha}$ is closed under the operators E_{i,α_j}^r with $1 \leq j \leq k$ and $1 \leq i < \alpha_j$ and $r \in R$.
- (c'_k) The \mathbf{k} -module $N_{n,\alpha} \subset \mathbf{k}[\Lambda_{n,\alpha,k+}]$ is homogeneous with respect to
- $$\{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+2 \leq j \leq d \text{ and } \alpha_{k+1} \leq i < \alpha_j\}.$$

Substep 2d. For $1 \leq k \leq d$, we construct a conservative poset map $\mathfrak{M}^{(k)} \rightarrow \mathfrak{N}^{(k-1)}$.

Notation defined: none.

Consider $M \in \mathfrak{M}^{(k)}$, so M is an $\mathbf{OI}(d)$ -submodule of $Q^{(k)}$. Define $N = M \cap Q^{(k-1)}$. We claim that $N \in \mathfrak{N}^{(k-1)}$. This requires checking the conditions $(a'.i'_{k-1})$, $(a'.ii'_{k-1})$, (b'_{k-1}) , and (c'_{k-1}) . Consider some $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$.

- Condition $(a'.i'_{k-1})$ asserts that $N_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k \leq j \leq d$ and $1 \leq i < \alpha_{k-1}$ and $r \in R$. This follows from the fact that both $M_{n,\alpha}$ and

$Q_{n,\alpha}^{(k-1)}$ are closed under multiplication by these elements. This is immediate for $Q_{n,\alpha}^{(k-1)}$. For $M_{n,\alpha}$, it follows from $(a.i_k)$, which says that $M_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k \leq j \leq d$ and $1 \leq i < \alpha_k$ and $r \in R$.

- Condition $(a'.ii'_{k-1})$ asserts that $N_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k \leq j \leq d$ and $\alpha_{k-1} \leq i < \alpha_j$ and $r \in R_{\geq 0}$. This follows from the fact that both $M_{n,\alpha}$ and $Q_{n,\alpha}^{(k-1)}$ are closed under multiplication by these elements. This is immediate for $Q_{n,\alpha}^{(k-1)}$. For $M_{n,\alpha}$, it follows from a combination of $(a.i_k)$, which handles the cases where $\alpha_{k-1} \leq i < \alpha_k$ and gives the stronger conclusion that we can use $r \in R$ instead of just $r \in R_{>0}$, and $(a.ii_k)$, which handles the cases where $\alpha_k \leq i < \alpha_j$. Here one might worry that $(a.ii_k)$ requires $k+1 \leq j \leq d$ instead of $k \leq j \leq d$; however, the case $j = k$ is not needed since no i satisfies $\alpha_k \leq i < \alpha_k$.
- Condition (b'_{k-1}) asserts that $N_{n,\alpha}$ is closed under the operators E_{i,α_j}^r with $1 \leq j \leq k-1$ and $1 \leq i < \alpha_j$ and $r \in R$. This follows from the fact that both $M_{n,\alpha}$ and $Q_{n,\alpha}^{(k-1)}$ are closed under these operators. This is immediate for $Q_{n,\alpha}^{(k-1)}$. For $M_{n,\alpha}$, it follows from (b_k) , which says that $M_{n,\alpha}$ is closed under the operators E_{i,α_j}^r with $1 \leq j \leq k$ and $1 \leq i < \alpha_j$ and $r \in R$.
- Condition (c'_{k-1}) asserts that $N_{n,\alpha}$ is homogeneous with respect to

$$\{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+1 \leq j \leq d \text{ and } \alpha_k \leq i < \alpha_j\}.$$

Condition (c_k) says that $M_{n,\alpha}$ is homogeneous with respect to this same set, and this homogeneity is preserved when we intersect $M_{n,\alpha}$ with $Q_{n,\alpha}^{(k-1)}$.

We thus can define a poset map $\mathfrak{M}^{(k)} \rightarrow \mathfrak{N}^{(k-1)}$ taking $M \in \mathfrak{M}^{(k)}$ to $M \cap Q_{n,\alpha}^{(k-1)}$. We claim that this poset map is conservative. In fact, it is even injective. Indeed, consider $M, M' \in \mathfrak{M}^{(k)}$. Let $N = M \cap Q_{n,\alpha}^{(k-1)}$ and $N' = M' \cap Q_{n,\alpha}^{(k-1)}$, and assume that $N = N'$. We claim that $M = M'$. By symmetry, it is enough to prove that $M \subset M'$. Consider $(n, \alpha) \in \mathbf{OI}(d)$ and $x \in M_{n,\alpha}$. We must prove that $x \in M'_{n,\alpha}$. We have $x \in Q_{n,\alpha}^{(k)}$. Setting

$$\begin{aligned} \mathbf{S} &= \{T_{i,j} \mid 1 \leq j \leq d, 1 \leq i < \alpha_j, \alpha_{k-1} \leq i < \alpha_k\} \\ &= \{T_{i,j} \mid k \leq j \leq d, \alpha_{k-1} \leq i < \alpha_k\}, \end{aligned}$$

there exists some $\tau \in \Lambda_{n,\alpha}(\mathbf{S})$ such that $\tau x \in Q_{n,\alpha}^{(k-1)}$. By $(a.i_k)$, we have $\tau x \in M_{n,\alpha}$, and thus $\tau x \in N_{n,\alpha}$. Since $N = N' \subset M'$, we deduce that $\tau x \in M'_{n,\alpha}$. Define $\tau^{-1} \in \Lambda_{n,\alpha}(\mathbf{S})$ to be the result of replacing all the $T_{i,j}^r$ terms in τ with $T_{i,j}^{-r}$. Another application of $(a.i_k)$ shows that $\tau^{-1}\tau x = x \in M'_{n,\alpha}$, as desired.

Substep 2e. For $0 \leq k \leq d-1$, we construct a conservative poset map $\mathfrak{N}^{(k)} \rightarrow \mathfrak{M}^{(k)}$.

Notation defined: none.

Fix some $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$. The most important difference between $\mathfrak{M}^{(k)}$ and $\mathfrak{N}^{(k)}$ is that by (c_k) the \mathbf{k} -modules making up $\mathfrak{M}^{(k)}$ must be homogeneous with respect to

$$\mathbf{S}_{n,\alpha,k} = \{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+1 \leq j \leq d \text{ and } \alpha_k \leq i < \alpha_j \text{ and } i \geq 1\},$$

while by (c'_k) the \mathbf{k} -modules making up $\mathfrak{N}^{(k)}$ must only be homogeneous with respect to

$$\mathbf{S}_{n,\alpha,k+1} = \{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+2 \leq j \leq d \text{ and } \alpha_{k+1} \leq i < \alpha_j\}.$$

The main function of our poset map $\mathfrak{N}^{(k)} \rightarrow \mathfrak{M}^{(k)}$ will be to achieve the needed increase in homogeneity.

For $x \in Q_{n,\alpha}^{(k)}$, we will define an ‘‘initial term’’ $\text{in}(x) \in Q_{n,\alpha}^{(k)}$ as follows. Define

$$\mathbf{S}'_{n,\alpha,k} = \mathbf{S}_{n,\alpha,k} \setminus \mathbf{S}_{n,\alpha,k+1} = \{T_{i,j} \mid k+1 \leq j \leq d \text{ and } \max(\alpha_k, 1) \leq i < \alpha_{k+1}\}.$$

Recall that R is identified as an additive group with \mathbf{Z}^λ and that $R_{\geq 0} = (\mathbf{Z}_{\geq 0})^\lambda \subset R$. Using the identification $R = \mathbf{Z}^\lambda$, we will frequently speak of the coordinates of elements of R . We define a total order on $\Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$ in two steps:

- We first order $\mathbf{S}'_{n,\alpha,k}$ by letting $T_{i,j} < T_{i',j'}$ if either $i < i'$ or if $i = i'$ and $j < j'$.
- We then order $\Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$ as follows. Consider distinct $\tau, \tau' \in \Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$. Enumerating the elements of $\mathbf{S}'_{n,\alpha,k}$ in increasing order as $T_{i_1,j_1}, \dots, T_{i_p,j_p}$, we can uniquely write

$$\tau = T_{i_1,j_1}^{r_1} \cdots T_{i_p,j_p}^{r_p} \quad \text{and} \quad \tau' = T_{i_1,j_1}^{r'_1} \cdots T_{i_p,j_p}^{r'_p}.$$

for some $r_i, r'_j \in R_{\geq 0}$. Let $1 \leq q \leq p$ be the minimal number such that $r_q \neq r'_q$. We then say that $\tau < \tau'$ if the first nonzero coordinate of $r'_q - r_q \in R = \mathbf{Z}^\lambda$ is positive.

For nonzero $x \in Q_{n,\alpha}^{(k)}$, we can uniquely write

$$x = \sum_{q=1}^m \tau_q y_q,$$

where for all $1 \leq q \leq m$ we have the following:

- $\tau_q y_q \neq 0$ for all q .
- $\tau_q \in \Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$, and the τ_q are enumerated in increasing order $\tau_1 < \tau_2 < \cdots < \tau_m$.
- $y_q \in \mathbf{k}[\Lambda_n(\mathbf{T}_n \setminus \mathbf{S}'_{n,\alpha,k})]$.

We then define $\text{in}(x) = \tau_m y_m \in Q_{n,\alpha}^{(k)}$. We also set $\text{in}(0) = 0$. We will call τ_m the **initial variable** of x , though we remark that this terminology will not be used again until the final paragraph of this substep.

We now construct the poset map $\mathfrak{N}^{(k)} \rightarrow \mathfrak{M}^{(k)}$ as follows. Consider $N \in \mathfrak{N}^{(k)}$. For $(n, \alpha) \in \mathbf{OI}(d)$, define $\text{in}(N)_{n,\alpha} \subset Q_{n,\alpha}^{(k)}$ to be the \mathbf{k} -span of $\{\text{in}(x) \mid x \in N_{n,\alpha}\}$. It is easy to see that $\text{in}(N)$ is an $\mathbf{OI}(d)$ -submodule of $Q^{(k)}$. We claim that $\text{in}(N) \in \mathfrak{M}^{(k)}$. To see this, we must check the conditions $(a.i_k)$, $(a.ii_k)$, (b_k) , and (c_k) . Consider some $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$.

- We delay $(a.i_k)$ until the end, so we start by verifying condition $(a.ii_k)$, which asserts that $\text{in}(N)_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k+1 \leq j \leq d$ and $\alpha_k \leq i < \alpha_j$ and $r \in R_{\geq 0}$. This is immediate from $(a'.ii'_k)$, which asserts that N is closed under multiplication by these same elements.
- Condition (b_k) asserts that $\text{in}(N)_{n,\alpha}$ is closed under the operators E_{i,α_j}^r with $1 \leq j \leq k$ and $1 \leq i < \alpha_j$ and $r \in R$. Condition (b'_k) says that $N_{n,\alpha}$ is closed under these operators. To prove that this implies that $\text{in}(N)_{n,\alpha}$ is also closed under these operators, it is enough to prove that for $x \in Q_{n,\alpha}^{(k)}$, we have

$$\text{in}(E_{i,\alpha_j}^r(x)) = E_{i,\alpha_j}^r(\text{in}(x)).$$

To help the reader understand the argument below, we recommend reviewing the correspondence between elements of Λ_n and $n \times d$ matrices from Substep 1a. For

nonzero x , write

$$x = \sum_{q=1}^m \tau_q y_q,$$

where for all $1 \leq q \leq m$ we have the following:

- $\tau_q y_q \neq 0$ for all q .
- $\tau_q \in \Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$, and the τ_q are enumerated in increasing order $\tau_1 < \tau_2 < \cdots < \tau_m$.
- $y_q \in \mathbf{k}[\Lambda_n(\mathbf{T}_n \setminus \mathbf{S}'_{n,\alpha,k})]$.

Since $i < \alpha_j \leq \alpha_k$, for all $1 \leq q \leq m$ we have

$$E_{i,\alpha_j}^r(\tau_q) = \tau_q \tau'_q \quad \text{and} \quad E_{i,\alpha_j}^r(y_q) = y_q y'_q$$

for some $\tau'_q \in \Lambda_n(\mathbf{T}_n \setminus \mathbf{S}'_{n,\alpha,k})$ and $y'_q \in \mathbf{k}[\Lambda_n(\mathbf{T}_n \setminus \mathbf{S}'_{n,\alpha,k})]$. We thus have

$$E_{i,\alpha_j}^r(x) = \sum_{q=1}^m E_{i,\alpha_j}^r(\tau_q) \cdot E_{i,\alpha_j}^r(y_q) = \sum_{q=1}^m \tau_q (\tau'_q y_q y'_q)$$

and

$$\text{in}(E_{i,\alpha_j}^r(x)) = \tau_m (\tau'_m y_m y'_m) = E_{i,\alpha_j}^r(\text{in}(x)),$$

as desired.

- Condition (c_k) asserts that $\text{in}(N)_{n,\alpha}$ is homogeneous with respect to

$$\mathbf{S}_{n,\alpha,k} = \{T_{\alpha_j,j} \mid 1 \leq j \leq d\} \cup \{T_{i,j} \mid k+1 \leq j \leq d \text{ and } \alpha_k \leq i < \alpha_j \text{ and } i \geq 1\}.$$

By (c'_k) , the \mathbf{k} -module $N_{n,\alpha}$ is homogeneous with respect to $\mathbf{S}_{n,\alpha,k+1}$, and the very definition of $\text{in}(N)_{n,\alpha}$ is designed to improve this to $\mathbf{S}_{n,\alpha,k}$.

- We now finally verify $(a.i_k)$, which asserts that $\text{in}(N)_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k \leq j \leq d$ and $1 \leq i < \alpha_k$ and $r \in R$. Condition $(a'.i'_k)$ says that $N_{n,\alpha}$ is closed under multiplication by $T_{i,j}^r$ with $k+1 \leq j \leq d$ and $1 \leq i \leq \alpha_j$ and $r \in R$, and this is preserved when we pass to $\text{in}(N)_{n,\alpha}$. We thus must only verify that $\text{in}(N)_{n,\alpha}$ is closed under multiplication by $T_{i,k}^r$ with $1 \leq i < \alpha_k$ and $r \in R$. Consider some $x \in \text{in}(N)_{n,\alpha}$. We must show that $T_{i,k}^r x \in \text{in}(N)_{n,\alpha}$. Using the already verified condition (c_k) , we can assume that $x = \tau y$ with

$$\tau \in \Lambda_{n,\alpha}(\mathbf{S}_{n,\alpha,k}) \quad \text{and} \quad y \in \mathbf{k}[\Lambda_{n,\alpha}(\mathbf{T}_n \setminus \mathbf{S}_{n,\alpha,k})].$$

Using the already verified condition (b_k) , we know that $E_{i,\alpha_k}^r(x) \in \text{in}(N)_{n,\alpha}$. We then calculate that

$$E_{i,\alpha_k}^r(x) = E_{i,\alpha_k}^r(\tau y) = E_{i,\alpha_k}^r(\tau) E_{i,\alpha_k}^r(y) = (\tau T_{i,k}^r \tau') \cdot y,$$

where τ' is a product of elements of $\{T_{i,j'}^{r'} \mid k+1 \leq j' \leq d, r' \in R\}$ that depends on τ and r and i and k . Letting $(\tau')^{-1}$ be the result of replacing each $T_{i,j'}^{r'}$ in τ' with $T_{i,j'}^{-r'}$, our already verified cases of $(a.i_k)$ imply that $\text{in}(N)_{n,\alpha}$ is closed under multiplication by $(\tau')^{-1}$. In particular,

$$(\tau')^{-1} \cdot E_{i,\alpha_k}^r(x) = (\tau')^{-1} \cdot (\tau T_{i,k}^r \tau') \cdot y = T_{i,k}^r \tau y = T_{i,k}^r x \in \text{in}(N)_{n,\alpha},$$

as desired.

The map $N \mapsto \text{in}(N)$ is thus a poset map from $\mathfrak{N}^{(k)}$ to $\mathfrak{M}^{(k)}$.

We claim that this is a conservative poset map. Indeed, consider $N_1, N_2 \in \mathfrak{N}^{(k)}$ such that $N_1 \subset N_2$ and $\text{in}(N_1) = \text{in}(N_2)$. We must prove that $N_1 = N_2$. Assume otherwise. Let $(n, \alpha) \in \mathbf{OI}(d)$ be such that $(N_1)_{n,\alpha} \subsetneq (N_2)_{n,\alpha}$. Pick $x \in (N_2)_{n,\alpha}$ such that $x \notin (N_1)_{n,\alpha}$ and such that the initial variable (see the second paragraph of this substep for the definition of this) of x is as small as possible among elements with these properties (this is possible since with the above ordering $\Lambda_{n,+}(\mathbf{S}'_{n,\alpha,k})$ does not have any infinite strictly decreasing chains). Since $\text{in}(N_1) = \text{in}(N_2)$, we can find some $x' \in (N_1)_{n,\alpha}$ such that $\text{in}(x') = \text{in}(x)$. But then $x - x' \in (N_2)_{n,\alpha}$ and $x - x' \notin (N_1)_{n,\alpha}$, while the initial variable of $x - x'$ is strictly smaller than the initial variable of x , a contradiction.

Step 3. *We prove that $\mathfrak{M}^{(0)}$ is noetherian.*

In Step 2, we reduced the theorem to showing that $\mathfrak{M}^{(0)}$ is noetherian. In this step, we will prove this. Defining

$$\Lambda_+ = \bigsqcup_{(n,\alpha) \in \mathbf{OI}(d)} \Lambda_{n,\alpha,+},$$

in Substep 3a we first construct a useful partial ordering on Λ_+ and prove that it is a well partial ordering (see below for the definition of this). In Substep 3b, we use this partial ordering to prove that $\mathfrak{M}^{(0)}$ is noetherian.

Substep 3a. *We construct a partial ordering on Λ_+ and prove that it is a well partial ordering.*

Notation defined: *none.*

We define a partial ordering on Λ_+ as follows. Consider $\tau, \tau' \in \Lambda_+$. We say that $\tau \preceq \tau'$ if the following condition is satisfied:

- Let $(n, \alpha), (n', \alpha') \in \mathbf{OI}(d)$ be such that $\tau \in \Lambda_{n,\alpha,+}$ and $\tau' \in \Lambda_{n',\alpha',+}$. We then require that there exists an $\mathbf{OI}(d)$ -morphism $\iota: (n, \alpha) \rightarrow (n', \alpha')$ and some $\tau'' \in \Lambda_{n',\alpha',+}$ such that $\tau' = \tau'' \cdot \iota_*(\tau)$.

It is clear that this is a partial ordering.

The main goal of this substep (which we will accomplish at the end after a number of preliminaries) is to prove that this partial ordering on Λ_+ is a well partial ordering, whose definition is as follows. A poset (\mathfrak{P}, \prec) is **well partially ordered** if every infinite sequence of elements of \mathfrak{P} contains an infinite weakly increasing subsequence. See [K] for a survey about well partial orderings. If \mathfrak{P} and \mathfrak{P}' are posets, then we will endow $\mathfrak{P} \times \mathfrak{P}'$ with the ordering where $(p_1, p'_1) \preceq (p_2, p'_2)$ if and only if $p_1 \preceq p_2$ and $p'_1 \preceq p'_2$. If \mathfrak{P} and \mathfrak{P}' are both well partially ordered, then so is $\mathfrak{P} \times \mathfrak{P}'$ (quick proof: given an infinite sequence in $\mathfrak{P} \times \mathfrak{P}'$, first pass to a subsequence to make the first coordinate weakly increasing, then pass to a further subsequence to make the second coordinate also weakly increasing).

Recall that we have identified the additive group of R with \mathbf{Z}^λ and that $R_{\geq 0} = (\mathbf{Z}_{\geq 0})^\lambda$. Using these identifications, we will speak of the coordinates of elements of R and $R_{\geq 0}$. Endow the set $R_{\geq 0} \cup \{\spadesuit\}$ with the following partial ordering:

- \spadesuit is not comparable to any element of $R_{\geq 0}$.
- For $r_1, r_2 \in R_{\geq 0}$, let $r_1 \preceq r_2$ if all the coordinates of $r_2 - r_1$ are nonnegative.

Since the usual ordering on $\mathbf{Z}_{\geq 0}$ is a well partial ordering, the restriction of our partial ordering to $R_{\geq 0} = (\mathbf{Z}_{\geq 0})^\lambda$ is also a well partial ordering. From this, it is easy to see that our partial ordering on $R_{\geq 0} \cup \{\spadesuit\}$ is also a well partial ordering. The product ordering on $(R_{\geq 0} \cup \{\spadesuit\})^d$ is thus also a well partial ordering.

Let \mathcal{W} denote the set of finite words in the alphabet $(R_{\geq 0} \cup \{\spadesuit\})^d$. Endow \mathcal{W} with the partial ordering where $w_1, w_2 \in \mathcal{W}$ satisfy $w_1 \preceq w_2$ if and only if the following condition is satisfied. Write $w_1 = \ell_1 \cdots \ell_n$ and $w_2 = \ell'_1 \cdots \ell'_{n'}$ with each ℓ_i and ℓ'_i an element of $(R_{\geq 0} \cup \{\spadesuit\})^d$. We then require that there exists a strictly increasing function $\iota: [n] \hookrightarrow [n']$ such that $\ell_i \preceq \ell'_{\iota(i)}$ for all $1 \leq i \leq n$. This partial ordering on \mathcal{W} is a well partial ordering by Higman's lemma [Hig, Theorem 4.3].

As promised, we now prove that the partial ordering on Λ_+ defined above is a well partial ordering. Let $\Psi: \Lambda_+ \rightarrow \mathcal{W}$ be the following set function. Consider $\tau \in \Lambda_{n,\alpha,+} \subset \Lambda_+$. Write $\alpha = (\alpha_1, \dots, \alpha_d)$, and expand out τ as

$$\tau = \prod_{\substack{1 \leq j \leq d \\ 1 \leq i \leq \alpha_j}} T_{i,j}^{r_{i,j}} \quad (r_{i,j} \in R_{\geq 0}).$$

For $1 \leq j \leq d$ and $1 \leq i \leq \alpha_j$, define $\bar{r}_{i,j} \in R_{\geq 0} \cup \{\spadesuit\}$ via the formula

$$\bar{r}_{i,j} = \begin{cases} r_{i,j} & \text{if } 1 \leq i < \alpha_j, \\ \spadesuit & \text{if } i = \alpha_j. \end{cases}$$

We remark that by definition we have $r_{\alpha_j,j} = 1$ for all $1 \leq j \leq d$. For $1 \leq i \leq n$, we define

$$\ell_i = (\bar{r}_{i,1}, \bar{r}_{i,2}, \dots, \bar{r}_{i,d}) \in (R_{\geq 0} \cup \{\spadesuit\})^d.$$

Finally, we define

$$\Psi(\tau) = \ell_1 \ell_2 \cdots \ell_n.$$

It is clear that Ψ is injective. What is more, it is immediate from the definitions that for all $\tau, \tau' \in \Lambda_+$ we have

$$\tau \preceq \tau' \quad \text{if and only if } \Psi(\tau) \preceq \Psi(\tau').$$

The key point here is that if we interpret elements of Λ_+ as matrices with d columns and entries in R_+ , the effect of an $\mathbf{OI}(d)$ -morphism on these matrices is to insert extra rows of zeros. Since Ψ is injective and \mathcal{W} is well partially ordered, so is Λ_+ , as claimed.

Substep 3b. *We prove that the poset $\mathfrak{M}^{(0)}$ is noetherian.*

Notation defined: *none.*

Let $(\Lambda_+, <)$ be the partially ordered set constructed in Substep 3a. By definition, $\mathfrak{M}^{(0)}$ is the poset of all $\mathbf{OI}(R)$ -modules $M \subset Q^{(0)}$ such that for all $(n, \alpha) \in \mathbf{OI}(d)$ with $\alpha = (\alpha_1, \dots, \alpha_d)$, the \mathbf{k} -module $M_{n,\alpha} \subset \mathbf{k}[\Lambda_{n,\alpha,+}]$ satisfies the following two properties:

- (†) It is closed under multiplication by $T_{i,j}^r$ for all $1 \leq j \leq d$ and $1 \leq i < \alpha_j$ and $r \in R_{\geq 0}$.
- (††) It is homogeneous with respect to all the possible $T_{i,j}$, i.e., with respect to

$$\{T_{i,j} \mid 1 \leq j \leq d \text{ and } 1 \leq i \leq \alpha_j\}.$$

Property (††) implies that $M_{n,\alpha}$ is spanned as a \mathbf{k} -module by elements of the form $c \cdot \tau$ with $c \in \mathbf{k}$ and $\tau \in \Lambda_{n,\alpha,+}$. Property (†) implies the following:

- (†††) Let $\tau_1 \in \Lambda_{n_1,\alpha_1,+} \subset \Lambda_+$ and $\tau_2 \in \Lambda_{n_2,\alpha_2,+} \subset \Lambda_+$ and $c \in \mathbf{k}$ be such that $c \cdot \tau_1 \in M_{n_1,\alpha_1}$ and $\tau_1 \leq \tau_2$. Then $c \cdot \tau_2 \in M_{n_2,\alpha_2}$.

Now assume for the sake of contradiction that $\mathfrak{M}^{(0)}$ is not noetherian. Let

$$M_1 \subsetneq M_2 \subsetneq M_3 \subsetneq \cdots$$

be an infinite strictly ascending chain in it. By $(\dagger\dagger)$, for all $i \geq 1$ there exists some $(n_i, \alpha_i) \in \mathbf{OI}(d)$ and some $\tau_i \in \Lambda_{n_i, \alpha_i, +}$ and some $c_i \in \mathbf{k}$ such that

$$(5.9) \quad c_i \cdot \tau_i \in (M_i)_{n_i, \alpha_i} \setminus (M_{i-1})_{n_i, \alpha_i}.$$

Since our partial ordering on Λ_+ is a well partial ordering, we can replace our sequence $\{M_i\}_{i=1}^\infty$ with a subsequence and assume that

$$\tau_1 \leq \tau_2 \leq \tau_3 \leq \cdots.$$

For $i \leq i'$, condition $(\dagger \dagger \dagger)$ implies that

$$c_i \cdot \tau_{i'} \in (M_i)_{n_{i'}, \alpha_{i'}}.$$

For all $q \geq 1$, applying this repeatedly with $i' = q + 1$ we see that for all $1 \leq q' \leq q$ we have

$$c_{q'} \cdot \tau_{q+1} \in (M_{q'})_{n_{q+1}, \alpha_{q+1}} \subset (M_q)_{n_{q+1}, \alpha_{q+1}}.$$

Defining I_q to be the ideal of \mathbf{k} generated by $\{c_1, \dots, c_q\}$, this implies that for all $d \in I_q$ we have

$$d \cdot \tau_{q+1} \in (M_q)_{n_{q+1}, \alpha_{q+1}}.$$

Since \mathbf{k} is noetherian, we can pick $q \gg 0$ such that $I_q = I_{q+1}$; in particular, $c_{q+1} \in I_q$. But this implies that

$$c_{q+1} \cdot \tau_{q+1} \in (M_q)_{n_{q+1}, \alpha_{q+1}},$$

contradicting (5.9).

5.4. A converse to Theorem 1.4. We now prove a converse to Theorem 1.4:

Proposition 5.10. *Let R be a ring and \mathbf{k} be a commutative ring such that the category of $\mathbf{OVI}(R)$ -modules over \mathbf{k} is locally noetherian. Then \mathbf{k} is noetherian and the additive group of R is finitely generated.*

Proof. Let P be the principal projective $\mathbf{OVI}(R)$ -module associated to R^2 and let P^+ be the submodule of P generated by all elements lying in P_n with $n > 2$. Then P/P^+ is a finitely generated $\mathbf{OVI}(R)$ -module with

$$(P/P^+)_n = \begin{cases} \mathbf{k}[U_2(R)] & \text{if } n = 2, \\ 0 & \text{otherwise.} \end{cases}$$

It follows that an $\mathbf{OVI}(R)$ -submodule of P/P^+ is exactly the same thing as a left ideal in $\mathbf{k}[U_2(R)]$, so $\mathbf{k}[U_2(R)]$ is a left-Noetherian ring. The group $U_2(R)$ is simply the additive group underlying R , so the proposition follows from the following lemma. \square

Lemma 5.11. *Let \mathbf{k} be a commutative ring and let A be an abelian group such that $\mathbf{k}[A]$ is noetherian. Then \mathbf{k} is noetherian and A is finitely generated.*

Proof. Since \mathbf{k} is a quotient of the noetherian ring $\mathbf{k}[A]$ via the augmentation homomorphism, it is noetherian. For a subgroup B of A , let I_B be the ideal of $\mathbf{k}[A]$ generated by $[b] - [0]$ with $b \in B$. Then $\mathbf{k}[A]/I_B = \mathbf{k}[A/B]$, and so B can be recovered from I_B as the elements $b \in A$ such that $[b] - [0] \in I_B$. Suppose that B_\bullet is an ascending chain of subgroups of A . Then I_{B_\bullet} is an ascending chain of ideals in $\mathbf{k}[A]$ and thus stabilizes. Thus the chain B_\bullet stabilizes as well, and so A is noetherian (and thus finitely generated) as an abelian group. \square

6. HOMOLOGY OF **OVI**-MODULES

In this section, R denotes a (not necessarily commutative) ring whose additive group is a finitely generated abelian group and \mathbf{k} denotes a commutative noetherian ring. Our goal is to prove Theorem 1.3 from the introduction, which says that if M is a finitely generated **OVI**-module then $\mathbf{H}_i(\mathbf{U}, M)$ is a finitely generated **OI**-module for all $i \geq 0$. This theorem is proved in §6.3 below after some preliminaries. We then prove in §6.4 an analogue of Theorem 1.3 where we allow upper triangular matrices that are not necessarily unipotent.

6.1. Homology of some **OI-groups.** Recall that a group Γ is of type FP over \mathbf{k} if the trivial $\mathbf{k}[\Gamma]$ -module \mathbf{k} admits a projective resolution \mathbf{P}_\bullet such that each \mathbf{P}_i is a finitely generated $\mathbf{k}[\Gamma]$ -module. In fact, it is equivalent to ask that each \mathbf{P}_i be a finitely generated free module; see [B, Theorem VIII.4.3]. Many natural classes of groups are of type FP including finite groups, finitely generated abelian groups, and lattices in semisimple Lie groups. See [B, Chapter VIII] for more information.

Proposition 6.1. *Let A be a group of type FP over \mathbf{k} and let \mathbf{E} be the **OI**-group $[n] \mapsto A^n$. Let M be an \mathbf{E} -module which is finitely generated as an **OI**-module. The following then hold.*

- (a) *The **OI**-module $\mathbf{H}_i(\mathbf{E}, M)$ is finitely generated for all $i \geq 0$.*
- (b) *Suppose A is abelian. Let $C \subset A$ be a finite index subgroup, let A_C^n denote the subgroup $\{(a_1, \dots, a_n) \in A^n \mid a_1 + \dots + a_n \in C\}$, and let \mathbf{E}_C be the **OI**-group $[n] \mapsto A_C^n$. Then the **OI**-module $\mathbf{H}_i(\mathbf{E}_C, M)$ is finitely generated for all $i \geq 0$.*

Proof. Pick a free resolution \mathbf{F}_\bullet of the $\mathbf{k}[A]$ -module \mathbf{k} such that each \mathbf{F}_i is a finitely generated $\mathbf{k}[A]$ -module and such that $\mathbf{F}_0 = \mathbf{k}[A]$. For each $n \geq 0$, the complex $(\mathbf{F}^{\otimes n})_\bullet$ is a free resolution of the $\mathbf{k}[A^n]$ -module \mathbf{k} .

For each $i \geq 0$, we assemble the i^{th} terms of $(\mathbf{F}^{\otimes n})_\bullet$ into an **OI**-module $X(i)$ as follows. First, define

$$X(i)_n = (\mathbf{F}^{\otimes n})_i = \bigoplus_{i_1 + \dots + i_n = i} \mathbf{F}_{i_1} \otimes \dots \otimes \mathbf{F}_{i_n}.$$

Next, given an **OI**-morphism $f: [n] \rightarrow [m]$, define $f_*: X(i)_n \rightarrow X(i)_m$ in the following way. Consider a summand $\mathbf{F}_{i_1} \otimes \dots \otimes \mathbf{F}_{i_n}$ of $X(i)_n$. For $1 \leq a' \leq m$, define

$$i'_{a'} = \begin{cases} i_a & \text{if } a' = f(a) \text{ for some } a \in [n], \\ 0 & \text{otherwise.} \end{cases}$$

We thus obtain a summand $\mathbf{F}_{i'_1} \otimes \dots \otimes \mathbf{F}_{i'_m}$ of $X(i)_m$. Define $f_*: X(i)_n \rightarrow X(i)_m$ to be the map that takes $\mathbf{F}_{i_1} \otimes \dots \otimes \mathbf{F}_{i_n}$ to $\mathbf{F}_{i'_1} \otimes \dots \otimes \mathbf{F}_{i'_m}$ by inserting terms that equal 1 in $\mathbf{k}[A] = \mathbf{F}_0$ into the needed places.

For each $i \geq 0$, define $Y(i)$ to be the **OI**-module $[n] \mapsto (X(i)_n \otimes M_n)_{A^n}$, where the subscript indicates that we are taking the A^n -coinvariants. The $Y(i)$ form a complex

$$\dots \longrightarrow Y(3) \longrightarrow Y(2) \longrightarrow Y(1) \longrightarrow Y(0) \longrightarrow 0$$

of **OI**-modules, and the **OI**-module $\mathbf{H}_i(\mathbf{E}, M)$ is the i^{th} homology group of this complex. By the local noetherianity of **OI** (Corollary 3.3), to prove that $\mathbf{H}_i(\mathbf{E}, M)$ is a finitely generated **OI**-module for all $i \geq 0$, it is enough to prove that each $Y(i)$ is a finitely generated **OI**-module, which we now do.

For each $i \geq 0$, the **OI**-module $X(i)$ is generated in finite degree (in fact, only terms of degree at most i are needed). Since M is finitely generated as an **OI**-module, it is in particular

generated in finite degree, so by Corollary 3.4 the \mathbf{OI} -module $X(i) \otimes M$ is also generated in finite degree. This implies that $Y(i)$ is also generated in finite degree. Since \mathbf{F}_i is a finitely generated $\mathbf{k}[A]$ -module for each $i \geq 0$ and M_n is a $\mathbf{k}[A^n]$ -module that is finitely generated as a \mathbf{k} -module for each $n \geq 0$, it follows that the \mathbf{k} -module $Y(i)_n = ((\mathbf{F}^{\otimes n})_i \otimes M_n)_{A^n}$ is a finitely generated \mathbf{k} -module for all $i, n \geq 0$. Combining this with the fact that each $Y(i)$ is generated in finite degree, we deduce that the \mathbf{OI} -module $Y(i)$ is finitely generated for all $i \geq 0$, as desired.

For the second statement, the restriction of $\mathbf{F}^{\otimes n}$ to A_C^n is still finitely generated since A_C^n is a finite index subgroup in A^n , and we can proceed as before. \square

Proposition 6.2. *Let A be a group of type FP over \mathbf{k} and let \mathbf{E}' be the $\mathbf{OI}(d)$ -group given by $\mathbf{E}'_{n,\lambda} = A^n$. Then $\mathbf{H}_i(\mathbf{E}', \mathbf{k})$ is a finitely generated $\mathbf{OI}(d)$ -module for all $i \geq 0$.*

Proof. The $\mathbf{OI}(d)$ -group \mathbf{E}' is the pullback of the \mathbf{OI} -group \mathbf{E} from Proposition 6.1 through the forgetful functor $\Phi: \mathbf{OI}(d) \rightarrow \mathbf{OI}$. Thus $\mathbf{H}_i(\mathbf{E}', \mathbf{k})$ is the pullback to $\mathbf{OI}(d)$ of the \mathbf{OI} -module $\mathbf{H}_i(\mathbf{E}, \mathbf{k})$, which is finitely generated by that proposition. The result now follows from the fact that Φ satisfies Property (F), which follows easily from Proposition 3.2. \square

6.2. A filtration. Our goal in this section is to prove the following result. Recall that $\bar{\Sigma}$ is the reduced shift functor on \mathbf{OI} -modules, i.e., the cokernel of the canonical map $M \rightarrow \Sigma(M)$. Also, P_d is the principal projective \mathbf{OVI} -module associated to the object R^d of \mathbf{OVI} .

Proposition 6.3. *The \mathbf{OI} -module $\bar{\Sigma}(\mathbf{H}_i(\mathbf{U}, P_d))$ has a filtration where the graded pieces are subquotients of \mathbf{OI} -modules of the form $\mathbf{H}_i(\mathbf{U}, P_e)$ with $e < d$ or $\mathbf{H}_j(\mathbf{U}, M)$ with $j < i$ and M a finitely generated \mathbf{OVI} -module.*

We begin with a number of lemmas. Recall that $\Phi: \mathbf{OI}(d) \rightarrow \mathbf{OI}$ and $\Psi: \mathbf{OVI}(d) \rightarrow \mathbf{OVI}$ are the forgetful functors. Also, \mathbf{U}_d is the $\mathbf{OI}(d)$ -group $(\mathbf{U}_d)_{n,\lambda} = U_{n,\lambda}$, where $U_{n,\lambda}$ is the group discussed in §4.1. Finally, the subscript ! is used to denote the left Kan extension discussed in §2.2.

Lemma 6.4. *Let M be an $\mathbf{OVI}(d)$ -module. We have an isomorphism of \mathbf{OI} -modules $\Phi_!(\mathbf{H}_i(\mathbf{U}_d, M)) \cong \mathbf{H}_i(\mathbf{U}, \Psi_!(M))$.*

Proof. Recall from Proposition 4.5 that

$$\Psi_!(M)_n = \bigoplus_{\lambda} \text{Ind}_{U_{n,\lambda}}^{U_n} (M_{n,\lambda}).$$

Thus, by Shapiro's lemma we have

$$\mathbf{H}_i(\mathbf{U}, \Psi_!(M))_n = \mathbf{H}_i(U_n, \Psi_!(M)_n) = \bigoplus_{\lambda} \mathbf{H}_i(U_{n,\lambda}, M_{n,\lambda}),$$

and this is exactly $\Phi_!(\mathbf{H}_i(\mathbf{U}_d, M))$ by Proposition 3.6. This shows that $\Phi_!(\mathbf{H}_i(\mathbf{U}_d, M))$ and $\mathbf{H}_i(\mathbf{U}, \Psi_!(M))$ agree on objects, and a moment's reflection shows that they also agree on morphisms. \square

Corollary 6.5. *We have $\mathbf{H}_i(\mathbf{U}, P_d) = \Phi_!(\mathbf{H}_i(\mathbf{U}_d, \mathbf{k}))$.*

Proof. Let $x = (R^d, \{e_i\}, \lambda) \in \mathbf{OVI}(d)$ where e_i is the standard basis and $\lambda = (1 < 2 < \dots < d)$. Set $y = (R^d, \{e_i\}) \in \mathbf{OVI}$. Then $\Psi_!(P_x) = P_y$ by (2.10). Since x is the initial object of $\mathbf{OVI}(d)$, we have $P_x(y) = \mathbf{k}[\text{Hom}(x, y)] = \mathbf{k}$ for all y , so $P_x = \mathbf{k}$. We thus have $\Psi_!(\mathbf{k}) = P_d$. Using the fact that P_y is just another name for P_d , the result now follows from Lemma 6.4 with $M = \mathbf{k}$. \square

Let $\mathbf{U}'_d = \Sigma(\mathbf{U}_d)$. This is the $\mathbf{OI}(d)$ -group given by $(\mathbf{U}'_d)_{n,\lambda} = U_{n+1,\lambda}$. The group $U_{n+1,\lambda}$ is the semi-direct product $U_{n,\lambda} \rtimes R^n$, and this description is functorial. More precisely, let \mathbf{E}_d be the $\mathbf{OI}(d)$ -group given by $(\mathbf{E}_d)_{n,\lambda} = R^n$. We then have homomorphisms of $\mathbf{OI}(d)$ -groups $i: \mathbf{U}_d \rightarrow \mathbf{U}'_d$ and $p: \mathbf{U}'_d \rightarrow \mathbf{U}_d$ with $pi = \text{id}$ and $\ker(p) = \mathbf{E}_d$. We observe that \mathbf{E}_d is in fact naturally an $\mathbf{OVI}(d)$ -group, and thus $\mathbf{H}_i(\mathbf{E}_d, \underline{\mathbf{k}})$ is naturally an $\mathbf{OVI}(d)$ -module. Proposition 6.2 says that $\mathbf{H}_i(\mathbf{E}_d, \underline{\mathbf{k}})$ is finitely generated as an $\mathbf{OI}(d)$ -module, so it is also finitely generated as an $\mathbf{OVI}(d)$ -module.

Lemma 6.6. *The $\mathbf{OI}(d)$ -module $\overline{\Sigma}(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))$ admits a filtration where the graded pieces are subquotients of $\mathbf{H}_i(\mathbf{U}_d, \mathbf{H}_{r-i}(\mathbf{E}_d, \underline{\mathbf{k}}))$ with $0 \leq i \leq r - 1$.*

Proof. The module $\overline{\Sigma}(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))$ is the cokernel of the map

$$\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}) \rightarrow \mathbf{H}_r(\mathbf{U}'_d, \underline{\mathbf{k}})$$

induced by the homomorphism $i: \mathbf{U}_d \rightarrow \mathbf{U}'_d$. The result therefore follows from Proposition 2.11, taking $\mathbf{G} = \mathbf{U}'_d$ and $\mathbf{K} = \mathbf{U}_d$ and $\mathbf{E} = \mathbf{E}_d$. \square

Recall that if M is an $\mathbf{OI}(d)$ -module, then right before Proposition 3.9 we defined an $\mathbf{OI}(d - 1)$ -module $\Delta(M)$.

Lemma 6.7. *We have $\Phi_!(\Delta(\mathbf{H}_i(\mathbf{U}_d, \underline{\mathbf{k}}))) = \Phi_!(\mathbf{H}_i(\mathbf{U}_{d-1}, \underline{\mathbf{k}}))$.*

Proof. By definition,

$$\Delta(\mathbf{H}_i(\mathbf{U}_d, \underline{\mathbf{k}}))_{n,\lambda} = \mathbf{H}_i(\mathbf{U}_d, \underline{\mathbf{k}})_{[n] \amalg \{\infty\}, \lambda \amalg \{\infty\}} = \mathbf{H}_i(U_{[n] \amalg \{\infty\}, \lambda \amalg \{\infty\}}, \underline{\mathbf{k}}).$$

Since $\{\infty\}$ is the maximal element of $[n] \amalg \{\infty\}$, we have

$$U_{[n] \amalg \{\infty\}, \lambda \amalg \{\infty\}} \cong U_{n,\lambda}.$$

Thus by Proposition 3.6 we have

$$\Phi_!(\Delta(\mathbf{H}_i(\mathbf{U}_d, \underline{\mathbf{k}}))) = \bigoplus_{\lambda} \mathbf{H}_i(U_{n,\lambda}, \underline{\mathbf{k}}),$$

the sum taken over appropriate $d - 1$ tuples λ . Again using Proposition 3.6, this is exactly $\Phi_!(\mathbf{H}_i(\mathbf{U}_{d-1}, \underline{\mathbf{k}}))$. \square

Proof of Proposition 6.3. We have $\mathbf{H}_r(\mathbf{U}, P_d) = \Phi_!(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))$ by Corollary 6.5. Thus by Proposition 3.9, we have

$$\overline{\Sigma}(\mathbf{H}_r(\mathbf{U}, P_d)) = \overline{\Sigma}(\Phi_!(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))) = \Phi_!(\overline{\Sigma}(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))) \oplus \Phi_!(\Delta(\mathbf{H}_r(\mathbf{U}_d, \underline{\mathbf{k}}))).$$

By Lemma 6.7, the second term on the right is $\Phi_!(\mathbf{H}_r(\mathbf{U}_{d-1}, \underline{\mathbf{k}}))$. By Corollary 6.5, this equals $\mathbf{H}_r(\mathbf{U}, P_{d-1})$. By Lemma 6.6, the first term admits a filtration where the graded pieces are subquotients of $\Phi_!(\mathbf{H}_i(\mathbf{U}_d, \mathbf{H}_{r-i}(\mathbf{E}_d, \underline{\mathbf{k}})))$ with $0 \leq i \leq r - 1$. Setting $N_i = \mathbf{H}_{r-i}(\mathbf{E}_d, \underline{\mathbf{k}})$, Proposition 6.2 implies that N_i is a finitely generated $\mathbf{OVI}(d)$ -module. Set $M_i = \Psi_!(N_i)$, so M_i is a finitely generated \mathbf{OVI} -module. By Lemma 6.4, we have

$$\Phi_!(\mathbf{H}_i(\mathbf{U}_d, N_i)) = \mathbf{H}_i(\mathbf{U}, M_i).$$

Combining all of the above, $\overline{\Sigma}(\mathbf{H}_r(\mathbf{U}, P_d))$ admits a filtration where one graded piece is $\mathbf{H}_r(\mathbf{U}, P_{d-1})$ and the other graded pieces are subquotients of $\mathbf{H}_i(\mathbf{U}, M_i)$ for $0 \leq i \leq r - 1$. The result follows. \square

6.3. Proof of Theorem 1.3. We now prove Theorem 1.3. Recall the statement: if R is a ring whose additive group is a finitely generated abelian group, \mathbf{k} is a commutative noetherian ring, and M is a finitely generated **OVI**-module, then $\mathbf{H}_i(\mathbf{U}, M)$ is a finitely generated **OI**-module for all $i \geq 0$. Fix such \mathbf{k} and R for the rest of this section. Consider the following statement:

(S_i) For a finitely generated **OVI**-module M , the **OI**-module $\mathbf{H}_i(\mathbf{U}, M)$ is finitely generated.

Let i be given and suppose that (S_j) is true for all $j < i$ (a vacuous condition if $i = 0$). We will prove (S_i), and this will establish the theorem.

We first show by induction on d that $\mathbf{H}_i(\mathbf{U}, P_d)$ is a finitely generated **OI**-module for all d . Suppose therefore that $\mathbf{H}_i(\mathbf{U}, P_e)$ is a finitely generated **OI**-module for $e < d$ (a vacuous condition for $d = 0$), and let us prove that $\mathbf{H}_i(\mathbf{U}, P_d)$ is a finitely generated **OI**-module. By Proposition 6.3, the **OI**-module $\overline{\Sigma}(\mathbf{H}_i(\mathbf{U}, P_d))$ has a filtration where each graded piece is a subquotient of an **OI**-module of the form $\mathbf{H}_i(\mathbf{U}, P_e)$ with $e < d$ or $\mathbf{H}_j(\mathbf{U}, M)$ with $j < i$ and M finitely generated. By the two inductive hypotheses in force, both of these kinds of **OI**-modules are finitely generated. Using the local noetherianity of **OI**-modules (Corollary 3.3), it follows that $\overline{\Sigma}(\mathbf{H}_i(\mathbf{U}, P_d))$ is a finitely generated **OI**-module. By Proposition 3.8, this implies that the **OI**-module $\mathbf{H}_i(\mathbf{U}, P_d)$ is finitely generated, as desired.

Let M be a finitely generated **OVI**-module. Consider an exact sequence

$$0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$$

where P is a finite direct sum of principal projective **OVI**-modules. Since the category of **OVI**-modules is locally noetherian (Theorem 1.4), the **OVI**-module K is finitely generated. We obtain an exact sequence

$$\mathbf{H}_i(\mathbf{U}, P) \rightarrow \mathbf{H}_i(\mathbf{U}, M) \rightarrow \mathbf{H}_{i-1}(\mathbf{U}, K).$$

By the previous paragraph, the **OI**-module $\mathbf{H}_i(\mathbf{U}, P)$ is finitely generated. By our inductive hypothesis (S_{i-1}), the **OI**-module $\mathbf{H}_{i-1}(\mathbf{U}, K)$ is finitely generated. Using the local noetherianity of **OI** (Corollary 3.3), it follows that the **OI**-module $\mathbf{H}_i(\mathbf{U}, M)$ is finitely generated. We have thus established (S_i), and the proof is complete.

Remark 6.8. The dimension shifting step in the third paragraph above is the only place in the proof of the theorem where the noetherianity of **OVI** is used. We never need noetherianity of **OVI**(d). \square

Remark 6.9. Suppose the additive group of R is a finite rank free abelian group. We outline an alternative way to get finite generation of the **OI**-module $[n] \mapsto \mathbf{H}_i(U_n(R); \mathbf{k})$. Let $\mathfrak{u}_n(R)$ be the Lie algebra of strictly upper-triangular $n \times n$ matrices over R . By [G, Theorem 4.3], there is a spectral sequence beginning with the Lie algebra homology of $\mathfrak{u}_n(R)$ which converges to $\mathbf{H}_i(U_n(R); \mathbf{k})$. The Lie algebra homology of $\mathfrak{u}_n(R)$ can be computed from the Koszul complex, whose terms are exterior powers of $\mathfrak{u}_n(R)$, and hence are finitely generated **OI**-modules (this is similar to the **OI**-structure on $\mathbf{F}^{\otimes n}$ in the proof of Proposition 6.1). By noetherianity, $\mathbf{H}_i(U_n(R); \mathbf{k})$ is a finitely generated **OI**-module. \square

6.4. A variant: relaxing unipotence. For each n , we let $B_n(R)$ denote the group of upper-triangular invertible $n \times n$ matrices with entries in R . We denote the **OI**-group $[n] \mapsto B_n(R)$ by \mathbf{B} . Also, if R is commutative and $C \subset R^\times$ is a subgroup, then let $B_n^C(R) \subset B_n(R)$

be the subgroup whose determinant lies in C . We denote the **OI**-subgroup $[n] \mapsto B_n^C(R)$ by \mathbf{B}^C .

The goal of this section is to prove Theorem 6.11 below, which is an analogue of Theorem 1.3 for \mathbf{B}^C . This requires the following lemma.

Lemma 6.10. *If R is commutative and the additive group of R is finitely generated, then the group of units R^\times is also finitely generated.*

Proof. If R is a domain, then it is either a subring of the ring of integers of a number field, in which case the statement follows from the Dirichlet unit theorem, or it is a finite field, in which case there is nothing to prove.

If R is reduced, then we have an injection $R \rightarrow \prod_P R/P$ where the product is over the finitely many associated primes of R . Thus we have an injection $R^\times \rightarrow \prod_P (R/P)^\times$, and hence R^\times is finitely generated.

Finally, in general we have an exact sequence of groups

$$0 \rightarrow \mathfrak{N}(R) \rightarrow R^\times \rightarrow (R/\mathfrak{N}(R))^\times \rightarrow 0,$$

where $\mathfrak{N}(R)$ is the nilradical of R equipped with the group structure $x * y = x + y + xy$, and the first map takes x to $1 + x$. (We note that the right map is surjective since any lift of a unit in $R/\mathfrak{N}(R)$ to R is automatically a unit.) By the previous cases, the abelian group $(R/\mathfrak{N}(R))^\times$ is finitely generated. The fact that the additive group of R is finitely generated implies that R is noetherian, so $\mathfrak{N}(R)^n = 0$ for some n . For each k , the $*$ operation on $\mathfrak{N}(R)$ descends to ordinary addition on $\mathfrak{N}(R)^k/\mathfrak{N}(R)^{k+1}$. Since the additive group $\mathfrak{N}(R)^k/\mathfrak{N}(R)^{k+1}$ is a subquotient of the finitely generated additive group of R , the additive group $\mathfrak{N}(R)^k/\mathfrak{N}(R)^{k+1}$ is finitely generated. Lifting additive generators for $\mathfrak{N}(R)/\mathfrak{N}(R)^2, \mathfrak{N}(R)^2/\mathfrak{N}(R)^3, \dots, \mathfrak{N}(R)^{n-1}/\mathfrak{N}(R)^n = \mathfrak{N}(R)^{n-1}$ to $\mathfrak{N}(R)$ gives generators for $\mathfrak{N}(R)$ with respect to the operation $*$. We conclude that R^\times is a finitely generated group. \square

Theorem 6.11. *Suppose that R is commutative and $C \subset R^\times$ is a subgroup. If M is a **B**-module which is finitely generated as an **OI**-module, then $\mathbf{H}_i(\mathbf{B}^C, M)$ is a finitely generated **OI**-module for any $i \geq 0$.*

Proof. Let $(R^\times)_C^n$ denote the subgroup of $(R^\times)^n$ consisting of sequences whose product lies in C . We have a short exact sequence of groups

$$1 \rightarrow U_n(R) \rightarrow B_n(R) \rightarrow (R^\times)_C^n \rightarrow 1.$$

The group R^\times is finitely generated by Lemma 6.10, and thus so is $(R^\times)_C^n$. The corollary now follows from the the Hochschild–Serre spectral sequence together with Theorem 1.3 and Proposition 6.1. \square

7. APPLICATION TO IWAHORI SUBGROUPS

The goal of this section is to prove Theorem 1.9, whose statement we now recall. Let \mathcal{O} be a number ring, let $\mathfrak{a} \subset \mathcal{O}$ be a nonzero proper ideal, and let \mathbf{k} be a commutative noetherian ring. For $i \geq 0$, let $X(i)$ be the **OI**-module defined by the rule $[n] \mapsto \mathbf{H}_i(\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}), \mathbf{k})$. We must prove that $X(i)$ is a finitely generated **OI**-module and that if \mathbf{k} is a field then $\dim X(i)_n$ equals a polynomial in n for $n \gg 0$. The polynomiality assertion follows from the finite generation assertion together with Proposition 3.5, so we must only prove that each $X(i)$ is a finitely generated **OI**-module.

Define $R = \mathcal{O}/\mathfrak{a}$ and let $C \subset R^\times$ be the image of \mathcal{O}^\times under the quotient map $\mathcal{O} \rightarrow R$. Let $\mathbf{GL}_n^C(R)$ be the subgroup of $\mathbf{GL}_n(R)$ consisting of matrices whose determinant lies in C . Strong approximation (see, e.g., [PIR, Chapter 7]) implies that the map $\mathbf{SL}_n(\mathcal{O}) \rightarrow \mathbf{SL}_n(R)$ is surjective. This implies that the map $\mathbf{GL}_n(\mathcal{O}) \rightarrow \mathbf{GL}_n^C(R)$ is surjective, which implies that the map $\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}) \rightarrow B_n^C(R)$ is surjective.

We thus have a short exact sequence

$$1 \longrightarrow \mathbf{GL}_n(\mathcal{O}, \mathfrak{a}) \longrightarrow \mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}) \longrightarrow B_n^C(R) \longrightarrow 1.$$

The associated Hochschild–Serre spectral sequence is of the form

$$H_i(B_n^C(R), H_j(\mathbf{GL}_n(\mathcal{O}, \mathfrak{a}), \mathbf{k})) \implies H_{i+j}(\mathbf{GL}_{n,0}(\mathcal{O}, \mathfrak{a}), \mathbf{k}) = X(i+j)_n.$$

Let $M(j)$ be the $\mathbf{OVI}(R)$ -module defined by $M(j)_n = H_j(\mathbf{GL}_n(\mathcal{O}, \mathfrak{a}), \mathbf{k})$. Naturality of the above spectral sequence induces a spectral sequence

$$(7.1) \quad H_i(B_n^C, M(j)) \implies X(i+j)$$

of \mathbf{OI} -modules.

Letting \mathbf{FI} be the category of finite sets and injections, the rule defining $M(j)$ also endows it with an \mathbf{FI} -module structure, which is finitely generated by [CEFN, Theorem D]. The inclusion $\mathbf{OI} \rightarrow \mathbf{FI}$ satisfies Property (F) (see [SaSn, Theorem 7.1.4]), so by Proposition 2.3 the induced \mathbf{OI} -module structure on $M(j)$ is also finitely generated. This implies in particular that $M(j)$ is a finitely generated $\mathbf{OVI}(R)$ -module. Theorem 6.11 now implies that $H_i(B_n^C(R), M(j))$ is a finitely generated \mathbf{OI} -module. Since the category of \mathbf{OI} -modules is locally noetherian (see Corollary 3.3), we can now deduce from (7.1) that each $X(i)$ is a finitely generated \mathbf{OI} -module, as desired.

REFERENCES

- [AHi] M. Aschenbrenner, C. J. Hillar, Finite generation of symmetric ideals, *Trans. Amer. Math. Soc.* 359 (2007), no. 11, 5171–5192, [arXiv:math/0411514v3](https://arxiv.org/abs/math/0411514v3).
Erratum: *Trans. Amer. Math. Soc.* 361 (2009), no. 10, 5627.
- [B] K. Brown, *Cohomology of Groups*, corrected reprint of the 1982 original, Graduate Texts in Mathematics 87, Springer-Verlag, New York, 1994.
- [CEF] T. Church, J. Ellenberg, B. Farb, FI-modules and stability for representations of symmetric groups, *Duke Math. J.* 164 (2015), no. 9, 1833–1910, [arXiv:1204.4533v4](https://arxiv.org/abs/1204.4533v4).
- [CEFN] T. Church, J. Ellenberg, B. Farb, Rohit Nagpal, FI-modules over Noetherian rings, *Geom. Top.* 18 (2014), 2951–2984, [arXiv:1210.1854v2](https://arxiv.org/abs/1210.1854v2).
- [Co] D. E. Cohen, On the laws of a metabelian variety, *J. Algebra* 5 (1967), 267–273.
- [D] W. G. Dwyer, Homology of integral upper-triangular matrices, *Proc. Amer. Math. Soc.* 94 (1985), no. 3, 523–528.
- [G] L. Grünenfelder, On the homology of filtered and graded rings, *J. Pure Appl. Algebra* 14 (1979), 21–37.
- [Ha] P. Hall, Finiteness conditions for soluble groups, *Proc. London Math. Soc.* (3) 4 (1954), 419–436.
- [Hig] G. Higman, Ordering by divisibility in abstract algebras, *Proc. London Math. Soc.* (3) 2 (1952), 326–336.
- [HilSu] C. Hillar, S. Sullivant, Finite Gröbner bases in infinite dimensional polynomial rings and applications, *Adv. Math.* 221 (2012), 1–25, [arXiv:0908.1777v2](https://arxiv.org/abs/0908.1777v2).
- [K] J. B. Kruskal, The theory of well-quasi-ordering: A frequently discovered concept, *J. Combinatorial Theory Ser. A* 13 (1972), 297–305.
- [LSz] R. Lee, R. H. Szczarba, On the homology and cohomology of congruence subgroups, *Invent. Math.* 33 (1976), no. 1, 15–53.

- [PIR] V. Platonov, A. Rapinchuk, *Algebraic Groups and Number Theory*, translated from the 1991 Russian original by R. Rowen, Pure and Applied Mathematics, 139, Academic Press, Boston, MA, 1994.
- [Pu] A. Putman, Stability in the homology of congruence subgroups, *Invent. Math.* 202 (2015), no. 3, 987–1027, [arXiv:1201.4876v5](https://arxiv.org/abs/1201.4876v5).
- [PuSa] A. Putman, S. Sam, Representation stability and finite linear groups, *Duke Math. J.* 166 (2017), no. 13, 2521–2598, [arXiv:1408.3694v3](https://arxiv.org/abs/1408.3694v3).
- [SaSn] S. Sam, A. Snowden, Gröbner methods for representations of combinatorial categories, *J. Amer. Math. Soc.* 30 (2017), 159–203, [arXiv:1409.1670v3](https://arxiv.org/abs/1409.1670v3).
- [St] R. Stanley, *Enumerative Combinatorics. Volume 1*. Second edition, Cambridge Studies in Advanced Mathematics 49, Cambridge University Press, Cambridge, 2012.
- [VdK] W. van der Kallen, Homology stability for linear groups, *Invent. Math.* 60 (1980), no. 3, 269–295.
- [W] C. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press, Cambridge, 1994.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, SOUTH BEND, IN

E-mail address: andyp@nd.edu

URL: <http://www.nd.edu/~andyp/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI

E-mail address: svs@math.wisc.edu

URL: <http://math.wisc.edu/~svs/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI

E-mail address: asnowden@umich.edu

URL: <http://www-personal.umich.edu/~asnowden/>