

TORSION IN THE CRYSTALLINE COHOMOLOGY OF SINGULAR VARIETIES

BHARGAV BHATT

ABSTRACT. This note discusses some examples showing that the crystalline cohomology of even very mildly singular projective varieties tends to be quite large. In particular, any singular projective variety with at worst ordinary double points has infinitely generated crystalline cohomology in at least two cohomological degrees. These calculations rely critically on comparisons between crystalline and derived de Rham cohomology.

Fix an algebraically closed field k of characteristic $p > 0$ with ring of Witt vectors W . Crystalline cohomology is a W -valued cohomology theory for varieties over k (see [Gro68, Ber74]). It is exceptionally well behaved on proper smooth k -varieties: the W -valued theory is finite dimensional [BO78], and the corresponding $W[1/p]$ -valued theory is a Weil cohomology theory [KM74] robust enough to support a p -adic proof of the Weil conjectures [Ked06].

Somewhat unfortunately, crystalline cohomology is often large and somewhat unwieldy outside the world of proper smooth varieties. For example, the crystalline cohomology of a smooth affine variety of dimension > 0 is always infinitely generated as a W -module by the Cartier isomorphism (see Remark 2.4). Even worse, it is not a topological invariant: Berthelot and Ogus showed [BO83, Appendix (A.2)] that the 0th crystalline cohomology group of a fat point in \mathbf{A}^2 has torsion (see also Example 3.4). In this note, we give more examples of such unexpected behaviour:

Theorem. *Let X be a proper lci k -variety. Then the crystalline cohomology of X is infinitely generated if any of the following conditions is satisfied:*

- (1) X has at least one isolated toric singularity, such as a node on a curve.
- (2) X has at least one conical singularity of low degree, such as an ordinary double point of any dimension.

The statement above is informal, and we refer the reader to the body of the note — see examples 3.5, 3.6, 3.12, and 3.13 — for precise formulations. In contrast to the Berthelot-Ogus example, our examples are reduced and lci. We do not know whether to regard these calculations as pathological or indicative of deeper structure; see Question 3.15.

Perhaps more interesting than the theorem above is our method, which relies on Illusie’s derived de Rham cohomology [Ill72]. This theory, which in hindsight belongs to derived algebraic geometry, is a refinement of classical de Rham cohomology that works better for singular varieties; the difference, roughly, is the replacement of the cotangent sheaf with the cotangent complex. Theorems from [Bha12] show: (a) derived de Rham cohomology agrees with crystalline cohomology for lci varieties, and (b) derived de Rham cohomology is computed by a “conjugate” spectral sequence whose E_2 -terms come from *coherent* cohomology on the Frobenius twist. These results transfer calculations from crystalline cohomology to coherent cohomology, where it is much easier to localise calculations at the singularities (see the proof of Proposition 3.1). As a bonus, this method yields a natural (infinite) increasing bounded below exhaustive filtration with finite-dimensional graded pieces on the crystalline cohomology of *any* lci proper variety.

Organisation of this note. In §1, we review the relevant results from derived de Rham cohomology together with the necessary categorical background. Next, we study (wedge powers of) the cotangent complex of some complete intersections in §2. This analysis is used in §3.1 to provide examples of some singular projective varieties (such as nodal curves, or lci toric varieties) whose crystalline cohomology is always infinitely generated; all these examples admit local lifts to W_2 where Frobenius also lifts. Examples which are not obviously liftable (such as ordinary double points in high dimensions) are discussed in §3.2. We conclude with the aforementioned Question 3.15.

Notation. Let k and W be as above, and set $W_2 = W/p^2$. For a k -scheme X , let $X^{(1)}$ denote the Frobenius-twist of X ; we identify the étale topology on X and $X^{(1)}$. We use $H_{\text{crys}}^n(X/k)$ and $H_{\text{crys}}^n(X/W)$ to denote Berthelot’s crystalline cohomology groups relative to k and W respectively. All sheaves are considered with respect to the Zariski topology (unless otherwise specified), and all tensor products are derived. We say that X lifts to W_2 compatibly with Frobenius if there exists a flat W_2 -scheme \mathcal{X} lifting X , and a map $\mathcal{X} \rightarrow \mathcal{X}$ lifting the Frobenius map on X and lying over the canonical Frobenius lift on W_2 . For fixed integers $a \leq b \in \mathbf{Z}$, we say that a complex K over some abelian category has amplitude in $[a, b]$ if $H^i(K) = 0$ for $i \notin [a, b] \subset \mathbf{Z}$. A complex K of abelian groups is connected (resp.

simply connected) if $H^i(K) = 0$ for $i > 0$ (resp. for $i \geq 0$). An infinitely generated module over a ring is one that is not finitely generated. All gradings are indexed by \mathbf{Z} unless otherwise specified. If A is a graded ring, then $A(-j)$ is the graded A -module defined $A(-j)_i = A_{i-j}$; we set $M(-j) := M \otimes_A A(-j)$ for any graded A -complex M . We use Δ for the category of simplices, and $\text{Ch}(\mathcal{A})$ for the category of chain complexes over an abelian category \mathcal{A} .

Acknowledgements. I thank Johan de Jong, Davesh Maulik, and Mircea Mustața for inspiring conversations. In particular, Example 3.5 was discovered in conversation with de Jong and Maulik, and was the genesis of this note. Both Pierre Berthelot and Arthur Ogus had also independently calculated a variant of this example (unpublished), and I thank them for their prompt response to email inquiries.

1. REVIEW OF DERIVED DE RHAM THEORY

In this section, we summarise some structure results in derived de Rham theory that will be relevant in the sequel. We begin by recalling in §1.1 some standard techniques for working with filtrations in the derived category; this provides the language necessary for the work in [Bha12] reviewed in §1.2.

1.1. Some homological algebra. In the sequel, we will discuss filtrations on objects of the derived category. To do so in a homotopy-coherent manner, we use the following model structure:

Construction 1.1. Fix a small category I , a Grothendieck abelian category \mathcal{B} , and set $\mathcal{A} = \text{Fun}(I, \mathcal{B})$. We endow $\text{Ch}(\mathcal{B})$ with the model structure of [Lur11, Proposition 1.3.5.3]: the cofibrations are termwise monomorphisms, while weak equivalences are quasi-isomorphisms. The category $\text{Fun}(I, \text{Ch}(\mathcal{B})) = \text{Ch}(\text{Fun}(I, \mathcal{B})) = \text{Ch}(\mathcal{A})$ inherits a projective model structure by [Lur09, Proposition A.2.8.2] where the fibrations and weak equivalences are defined termwise. By [Lur09, Proposition A.2.8.7], the pullback $D(\mathcal{B}) \rightarrow D(\mathcal{A})$ induced by the constant map $I \rightarrow \{1\}$ has a left Quillen adjoint $D(\mathcal{A}) \rightarrow D(\mathcal{B})$ that we call a “homotopy-colimit over I ”. In fact, exactly the same reasoning shows: given a map $\phi : I \rightarrow J$ of small categories, the pullback $\phi^* : D(\text{Fun}(J, \mathcal{B})) \rightarrow D(\text{Fun}(I, \mathcal{B}))$ induced by composition with ϕ has a left Quillen adjoint $\phi_! : D(\text{Fun}(I, \mathcal{B})) \rightarrow D(\text{Fun}(J, \mathcal{B}))$ if $\text{Ch}(\text{Fun}(I, \mathcal{B}))$ and $\text{Ch}(\text{Fun}(J, \mathcal{B}))$ are given the projective model structures as above; we often refer to $\phi_!$ as a “homotopy-colimit along fibres of ϕ .” The most relevant examples of ϕ for us are: the projections $\Delta^{\text{opp}} \rightarrow \{1\}$, $\Delta^{\text{opp}} \times \mathbf{N} \rightarrow \mathbf{N}$ and $\mathbf{N} \rightarrow \{1\}$.

Using Construction 1.1, we can talk about increasing filtrations on objects of derived categories.

Construction 1.2. Let \mathcal{B} be a Grothendieck abelian category, and let $\mathcal{A} := \text{Fun}(\mathbf{N}, \mathcal{B})$, where \mathbf{N} is the category associated to the poset \mathbf{N} with respect to the usual ordering. There is a homotopy-colimit functor $F : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ which is left Quillen adjoint to the pullback $D(\mathcal{B}) \rightarrow D(\mathcal{A})$ induced by the constant map $\mathbf{N} \rightarrow \{1\}$; we informally refer to an object $K \in D(\mathcal{A})$ as an increasing (or \mathbf{N} -indexed) exhaustive filtration on the object $F(K) \in D(\mathcal{B})$. There are also restriction functors $[n]^* : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ for each $n \in \mathbf{N}$, and maps $[n]^* \rightarrow [m]^*$ for $n \leq m$ coherently compatible with composition. For each $n \in \mathbf{N}$, the cone construction defines a functor $\text{gr}_n : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ and an exact triangle $[n-1]^* \rightarrow [n]^* \rightarrow \text{gr}_n$ of functors $D(\mathcal{A}) \rightarrow D(\mathcal{B})$; for a filtered object $K \in D(\mathcal{B})$, we often use $\text{gr}_n(K)$ to denote gr_n applied to the specified lift of K to $D(\mathcal{A})$. A map $K_1 \rightarrow K_2$ in $D(\mathcal{A})$ is an equivalence if and only if $[n]^*K_1 \rightarrow [n]^*K_2$ is so for all $n \in \mathbf{N}$ if and only if $\text{gr}_n(K_1) \rightarrow \text{gr}_n(K_2)$ is so for all $n \in \mathbf{N}$. Given a cochain complex K over \mathcal{B} , the association $n \mapsto \tau_{\leq n} K$ defines an object of $D(\mathcal{A})$ lifting the image of $K \in D(\mathcal{B})$ under F .

Remark 1.3. The “cone construction” used in Construction 1.2 to define gr_n needs clarification: there is no functor $\text{Fun}([0 \rightarrow 1], D(\mathcal{B})) \rightarrow D(\mathcal{B})$ which incarnates the chain-level construction of the cone. However, the same construction *does* define a functor $D(\text{Fun}([0 \rightarrow 1], \mathcal{B})) \rightarrow D(\mathcal{B})$, which suffices for the above application (as there are restriction functors $D(\mathcal{A}) \rightarrow D(\text{Fun}([0 \rightarrow 1], \mathcal{B}))$ for each map $[0 \rightarrow 1] \rightarrow \mathbf{N}$ in \mathbf{N}).

1.2. The derived de Rham complex and the conjugate filtration. We first recall the definition:

Construction 1.4. For a morphism $f : X \rightarrow S$ of schemes, following [Ill72, §VIII.2], the derived de Rham complex $\text{dR}_{X/S} \in \text{Ch}(\text{Mod}_{f^{-1}\mathcal{O}_S})$ is defined as the homotopy-colimit over Δ^{opp} of the simplicial cochain complex $\Omega_{P_\bullet/f^{-1}\mathcal{O}_S}^* \in \text{Fun}(\Delta^{\text{opp}}, \text{Ch}(\text{Mod}_{f^{-1}\mathcal{O}_S}))$, where P_\bullet is a simplicial free $f^{-1}\mathcal{O}_S$ -algebra resolution of \mathcal{O}_X . When S is an \mathbf{F}_p -scheme, the de Rham differential is linear over the p^{th} -powers, so $\text{dR}_{X/S}$ can be viewed as an object of $\text{Ch}(\text{Mod}_{\mathcal{O}_{X^{(1)}}})$, where $X^{(1)} = X \times_{S, \text{Frob}} S$ is the (derived) Frobenius-twist of X (which is the usual one if f is flat).

The following theorem summarises the relevant results from [Bha12] about this construction:

Theorem 1.5. *Let X be a k -scheme. Then:*

- (1) The complex $dR_{X/k} \in \text{Ch}(\text{Mod}_{\mathcal{O}_{X^{(1)}}})$ comes equipped with a canonical increasing bounded below separated exhaustive filtration $\text{Fil}_{\bullet}^{\text{conj}}$ called the conjugate filtration. The graded pieces are computed by

$$\text{Cartier}_i : \text{gr}_i^{\text{conj}}(dR_{X/k}) \simeq \wedge^i L_{X^{(1)}/k}[-i].$$

In particular, if X is lci, then $\text{Fil}_i^{\text{conj}}(dR_{X/k})$ is a perfect $\mathcal{O}_{X^{(1)}}$ -complex for all i .

- (2) The formation of $dR_{X/k}$ and the conjugate filtration commutes with étale localisation on $X^{(1)}$.
(3) There exists a canonical morphism

$$\text{R}\Gamma(X^{(1)}, dR_{X/k}) \rightarrow \text{R}\Gamma_{\text{crys}}(X/k, \mathcal{O})$$

that is an isomorphism when X is an lci k -scheme.

- (4) If there is a lift of X to W_2 together with a compatible lift of Frobenius, then the conjugate filtration is split, i.e., there is an isomorphism

$$\bigoplus_{i \geq 0} \wedge^i L_{X^{(1)}/k}[-i] \simeq dR_{X/k}$$

whose restriction to the i^{th} summand on the left splits Cartier_i .

Remark 1.6. Theorem 1.5 can be regarded as an analogue of the results of Cartier (as explained in [DI87], say) and Berthelot [Ber74] to the singular case. In particular, when X is quasi-compact, quasi-separated and lci, parts (1) and (3) of Theorem 1.5 together with the end of Remark 1.7 yield a “conjugate” spectral sequence

$$E_2^{p,q} : H^p(X^{(1)}, \wedge^q L_{X^{(1)}/k}) \Rightarrow H_{\text{crys}}^{p+q}(X/k).$$

In the sequel, instead of using this spectral sequence, we will directly use the filtration on $dR_{X/k}$ and the associated exact triangles; this simplifies bookkeeping of indices.

Remark 1.7. We explain the interpretation of Theorem 1.5 using the language of §1.1. Let $\mathcal{B} = \text{Mod}(\mathcal{O}_{X^{(1)}})$, and let $\mathcal{A} = \text{Fun}(\mathbf{N}, \mathcal{B})$. The construction of the derived de Rham complex $dR_{X/k} \in D(\mathcal{B})$ naturally lifts to an object $\mathcal{E} \in D(\mathcal{A})$ under F : if $P_{\bullet} \rightarrow \mathcal{O}_X$ is the canonical free k -algebra resolution of \mathcal{O}_X , then $\Omega_{P_{\bullet}/k}^* \otimes_{P_{\bullet}^{(1)}} \mathcal{O}_{X^{(1)}}$ defines an object of $D(\text{Fun}(\Delta^{\text{opp}} \times \mathbf{N}, \mathcal{B}))$ via $(m, n) \mapsto (\tau_{\leq n} \Omega_{P_m/k}^*) \otimes_{P_m^{(1)}} \mathcal{O}_{X^{(1)}}$, and its homotopy-colimit over Δ^{opp} (i.e., its pushforward along $D(\text{Fun}(\Delta^{\text{opp}} \times \mathbf{N}, \mathcal{B})) \rightarrow D(\text{Fun}(\mathbf{N}, \mathcal{B}))$) defines the desired object $\mathcal{E} \in D(\mathcal{A})$. This construction satisfies $[n]^* \mathcal{E} \simeq \text{Fil}_n^{\text{conj}}(dR_{X/k})$, so $\text{gr}_n(\mathcal{E}) \simeq \text{gr}_n^{\text{conj}}(dR_{X/k})$ for all $n \in \mathbf{N}$. This lift $\mathcal{E} \in D(\mathcal{A})$ of $dR_{X/k} \in D(\mathcal{B})$ is implicit in any discussion of the conjugate filtration on $dR_{X/k}$ in this note (as in Theorem 1.5 (1), for example). In the sequel, we abuse notation to let $dR_{X/k}$ also denote $\mathcal{E} \in D(\mathcal{A})$. When X is quasi-compact and quasi-separated, cohomology commutes with filtered colimits, so $\text{R}\Gamma(X^{(1)}, dR_{X/k}) \simeq \text{colim}_n \text{R}\Gamma(X^{(1)}, \text{Fil}_n^{\text{conj}}(dR_{X/k}))$. In particular, when restricted to proper varieties, derived de Rham cohomology can be written as a filtered colimit of (complexes of) finite dimensional vector spaces *functorially* in X .

2. SOME FACTS ABOUT LOCAL COMPLETE INTERSECTIONS

In order to apply Theorem 1.5 to compute crystalline cohomology, we need good control on (wedge powers of) the cotangent complex of an lci singularity. The following lemma collects most of the results we will use in §3.1.

Lemma 2.1. *Let (A, \mathfrak{m}) be an essentially finitely presented local k -algebra with an isolated lci singularity at $\{\mathfrak{m}\}$. Let $N = \dim_k(\mathfrak{m}/\mathfrak{m}^2)$ be the embedding dimension. Then:*

- (1) $\wedge^n L_{A/k}$ is a perfect complex for all n . For $n \geq N$, $\wedge^n L_{A/k}$ can be represented by a complex of finite free A -modules lying between cohomological degrees $-n$ and $-n + N$ with differentials that are 0 modulo \mathfrak{m} .
- (2) For any $n \geq N$, the complex $\wedge^n L_{A/k}$ has finite length cohomology groups.
- (3) For any $n > N$, the group $H^{-n+N}(\wedge^n L_{A/k})$ is non-zero.
- (4) For any $n > N$, there exists an integer $0 < i \leq N$ such that $H^{-n+N-i}(\wedge^n L_{A/k})$ is non-zero.
- (5) If $\dim(A) > 0$ and $n > N$, then $H^{-n}(\wedge^n L_{A/k}) = 0$, so the integer i in (4) is strictly less than N .

Proof. Choose a polynomial algebra $P = k[x_1, \dots, x_N]$ and a map $P \rightarrow A$ such that $\Omega_{P/k}^1 \otimes_P A \rightarrow \Omega_{A/k}^1$ is surjective. By comparing dimensions, the induced map $\Omega_{P/k}^1 \otimes_P A \otimes_A A/\mathfrak{m} \rightarrow \Omega_{A/k}^1 \otimes_A A/\mathfrak{m}$ is an isomorphism. Now consider the exact triangle

$$L_{A/P}[-1] \rightarrow \Omega_{P/k}^1 \otimes_P A \rightarrow L_{A/k}.$$

The lci assumption on A and the choice of P ensure that $L_{A/P}[-1]$ is a free A -module of some rank r . Since $\mathrm{Spec}(A)$ is singular at \mathfrak{m} , we must have $r > 0$. The previous triangle then induces a (non-canonical) equivalence

$$\left(A^{\oplus r} \xrightarrow{T} A^{\oplus N} \right) \simeq L_{A/k}.$$

The map T must be 0 modulo \mathfrak{m} as $A^{\oplus N} \rightarrow L_{A/k}$ induces an isomorphism on H^0 after reduction modulo \mathfrak{m} , so the above presentation yields an identification

$$L_{A/k} \otimes_A k \simeq (k^{\oplus r}[1]) \oplus k^{\oplus N}.$$

Computing wedge powers gives

$$\wedge^n(L_{A/k}) \otimes_A k \simeq \bigoplus_{a=0}^N \left(\wedge^a(k^{\oplus N}) \otimes \Gamma^{n-a}(k^{\oplus r}) \right) [n-a], \quad (*)$$

where Γ^* is the divided-power functor; here we use that $\wedge^n(V[1]) = \Gamma^n(V)[n]$ for a flat k -module V over a ring k (see [Qui70, §7]). We now show the desired claims:

- (1) The perfectness of $\wedge^n L_{A/k}$ follows from the perfectness of $L_{A/k}$. The desired representative complex can be constructed as a Koszul complex on the map T above (see the proof of Lemma 2.5 (4) below); all differentials will be 0 modulo \mathfrak{m} by functoriality since T is so.
- (2) We must show that $(\wedge^n L_{A/k})_{\mathfrak{p}} = 0$ for any $\mathfrak{p} \in \mathrm{Spec}(A) - \{\mathfrak{m}\}$ and $n \geq N$. The functor $\wedge^n L_{-/k}$ commutes with localisation, so we must show that $\wedge^n L_{A_{\mathfrak{p}}/k} = 0$ for \mathfrak{p} and n as before, but this is clear: $A_{\mathfrak{p}}$ is the localisation of smooth k -algebra of dimension $\leq \dim(A) < N$ for any such \mathfrak{p} .
- (3) By (1), $H^{-n+N}(\wedge^n L_{A/k}) = 0$ if and only if $\wedge^n L_{A/k}$ has amplitude in $[-n, -n + N - 1]$. However, in the latter situation, the complex $\wedge^n L_{A/k} \otimes_A k$ would have no cohomology in degree $-n + N$, contradicting formula (*); note that $r \geq 1$ by the assumption that $\mathrm{Spec}(A)$ is singular at \mathfrak{m} .
- (4) Assume the assertion of the claim is false. Then (3) shows that $\wedge^n L_{A/k}$ is concentrated in a single degree, so $\wedge^n L_{A/k} \simeq M[-n + N]$ for some finite length A -module M . By (1), M has finite projective dimension. The Auslander-Buchsbaum formula and the fact that A is Cohen-Macaulay then show that the projective dimension of M is actually $\dim(A)$. Hence, $M \otimes_A k$ has at most $\dim(A) + 1$ non-zero cohomology groups. On the other hand, formula (*) shows that $\wedge^n L_{A/k} \otimes_A k$ has $N + 1$ distinct cohomology groups. Hence, $N \leq \dim(A)$, which contradicts the assumption that $\mathrm{Spec}(A)$ is singular at \mathfrak{m} .
- (5) Set $M := H^{-n}(\wedge^n L_{A/k})$, and assume $M \neq 0$. Then M has finite length by (2), and occurs as the kernel of a map of free A -modules by (1). Non-zero finite length R -modules cannot be found inside free R -modules for any S_1 -ring R of positive dimension, which is a contradiction since complete intersections are S_1 . \square

Remark 2.2. The assumption $\dim(A) > 0$ is necessary in Lemma 2.1 (5). For example, set $A = k[\epsilon]/(\epsilon^p)$. Then $N = \dim_k(\mathfrak{m}/\mathfrak{m}^2) = 1$, and $L_{A/k} \simeq A[1] \oplus A$. Applying \wedge^n for $n > 0$, we get

$$\wedge^n(L_{A/k}) \simeq \Gamma^n(A)[n] \oplus \Gamma^{n-1}(A)[n-1],$$

which certainly has non-zero cohomology in degree $-n$.

Using Lemma 2.1, we can show that the crystalline cohomology of an isolated lci singularity is infinitely generated in a very strong sense:

Corollary 2.3. *Let (A, \mathfrak{m}) be as in Lemma 2.1. Assume that A admits a lift to W_2 compatible with Frobenius. Then*

- (1) $H_{\mathrm{crys}}^i(\mathrm{Spec}(A)/k) \simeq \bigoplus_{j \geq 0} H^0(\mathrm{Spec}(A)^{(1)}, \wedge^j L_{A^{(1)}/k}[i-j])$ for all i .
- (2) $H_{\mathrm{crys}}^N(\mathrm{Spec}(A)/k)$ is infinitely generated as an $A^{(1)}$ -module.

Proof. Note that $H_{\mathrm{crys}}^*(\mathrm{Spec}(A)/k)$ is an $A^{(1)}$ -module since any divided-power thickening of A is an $A^{(1)}$ -algebra.

- (1) This follows from Theorem 1.5 (4) and the vanishing of higher quasi-coherent sheaf cohomology on affines.
- (2) This follows from Lemma 2.1 (3). \square

Remark 2.4. Let us explain the phrase ‘‘strong sense’’ appearing before Corollary 2.3. If A is an essentially smooth k -algebra, then $H_{\mathrm{crys}}^*(\mathrm{Spec}(A)/k)$ is infinitely generated over k , but not over $A^{(1)}$: the Cartier isomorphism shows that $H_{\mathrm{crys}}^i(\mathrm{Spec}(A)/k) \simeq \Omega_{A^{(1)}/k}^i$, which is a finite (and even locally free) $A^{(1)}$ -module. It is this latter finiteness that also breaks down in the singular setting of Corollary 2.3.

We also record a more precise result on the wedge powers of the cotangent complex for the special case of the co-ordinate ring of a smooth hypersurface; this will be used in §3.2.

Lemma 2.5. *Let A be the localisation at 0 of $k[x_0, \dots, x_N]/(f)$, where f is a homogeneous degree d polynomial defining a smooth hypersurface in \mathbf{P}^N . Assume $p \nmid d$. Then*

- (1) A is graded.
- (2) The quotient $M = A/(\frac{\partial f}{\partial x_0}, \dots, \frac{\partial f}{\partial x_N})$ is a finite length graded A -module whose non-zero weights j are contained in the interval $0 \leq j \leq (d-2)(N+1)$.
- (3) The A -linear Koszul complex $K := K_A(\{\frac{\partial f}{\partial x_i}\})$ of the sequence of partials is equivalent to $M \oplus M(-d)[1]$ as a graded A -complex.
- (4) For $n > N$, we have an equivalence of graded A -complexes

$$\wedge^n L_{A/k}[-n] \simeq M((N+1)(d-1) - nd)[-N-1] \oplus M((N+1)(d-1) - nd - d)[-N].$$

- (5) Assume that N and d satisfy $N(d-2) < d+2$. Fix j and n with $N < j < n$. Then all graded k -linear maps

$$\wedge^n L_{A/k}[-n] \rightarrow \wedge^j L_{A/k}[-j][1]$$

are nullhomotopic as graded k -linear maps.

Proof. Let $S = k[x_0, \dots, x_N]$ denote the polynomial ring. We note first the assumption $p \nmid d$ implies (by the Euler relation) that f lies in the ideal $J(f) \subset S$ generated by the sequence $\{\frac{\partial f}{\partial x_i}\}$ of partials. Since f defines a smooth hypersurface, the preceding sequence cuts out a zero-dimensional scheme in S , and hence must be a regular sequence by Auslander-Buchsbaum. In particular, each $\frac{\partial f}{\partial x_i}$ is non-zero of degree $d-1$. We now prove the claims:

- (1) This is clear.
- (2) Since $f \in J(f)$, the quotient M is identified with $S/J(f)$, so the claim follows from [Voi07, Corollary 6.20].
- (3) Consider the S -linear Koszul complex $L := K_S(\{\frac{\partial f}{\partial x_i}\})$ of the sequence of partials. Since the partials span a regular sequence in S , we have an equivalence $L \simeq S/J(f) \simeq A/J(f) \simeq M$ of graded S -modules. Now the complex K is simply $L \otimes_S A \simeq M \otimes_S A$. Since M is already an A -module, we get an identification $K \simeq M \otimes_A (A \otimes_S A)$ as graded A -modules, where the right hand side is given the A -module structure from the last factor. The resolution $(S(-d) \xrightarrow{f} S) \simeq A$ then shows that $K \simeq M \otimes_A (A(-d) \xrightarrow{0} A) \simeq M \oplus M(-d)[1]$.
- (4) Set $L = (f)/(f^2)$, $E = \Omega_{S/k}^1 \otimes_S A$, and $c : L \rightarrow E$ to be the map defined by differentiation. Then the two-term complex defined by c is identified with $L_{A/k}$. Taking wedge powers for $n > N$ then shows (see [KS04, Corollary 1.2.7], for example) that the complex

$$\Gamma^n(L) \otimes_A \wedge^0(E) \rightarrow \Gamma^{n-1}(L) \otimes_A \wedge^1(E) \rightarrow \dots \rightarrow \Gamma^{n-(N+1)}(L) \otimes_A \wedge^{N+1}(E) \quad (**)$$

computes $\wedge^n L_{A/k}[-n]$; here the term on the left is placed in degree 0. Explicitly, the differential

$$\Gamma^i(L) \otimes_A \wedge^k(E) \rightarrow \Gamma^{i-1}(L) \otimes_A \wedge^{k+1}(E)$$

is given by

$$\gamma_i(f) \otimes \omega \mapsto \gamma_{i-1}(f) \otimes (c(f) \wedge \omega) = (-1)^k \cdot \gamma_{i-1}(f) \otimes (\omega \wedge df).$$

In particular, if we trivialise $\Gamma^i(L)$ using $\gamma_i(f)$, then this differential is identified with left-multiplication by df in the exterior algebra $\wedge^*(E)$. We leave it to the reader to check that the complex $(**)$ above is isomorphic to $K((N+1)(d-1) - nd)[-N-1]$; the rest follows from (3).

- (5) Let $M' = M((N+1)(d-1) - nd)[-N-1]$. Then M' is, up to a shift, a graded A -module whose weights lie in an interval size of $(d-2)(N+1)$ by (2). By (4), we have

$$\wedge^n L_{A/k}[-n] \simeq M'(-nd) \oplus M'(-nd-d)[1]$$

and

$$\wedge^j L_{A/k}[-j][1] \simeq M'(-jd)[1] \oplus M'(-jd-d)[2].$$

Thus, we must check that all graded k -linear maps $M'(-nd-d) \rightarrow M'(-jd)$ are nullhomotopic. Twisting, it suffices to show M' and $M'((n+1-j)d)$ do not share a weight. If they did, then $(n+1-j)d \leq (d-2)(N+1)$. Since $j < n$, this implies $2d \leq (d-2)(N+1)$, i.e., $d+2 \leq N(d-2)$, which contradicts the assumption. \square

Remark 2.6. The assumption $N(d-2) < d+2$ in Lemma 2.5 (5) is satisfied in exactly the following cases: $N \geq 5$ with $d = 2$, $N = 3, 4$ with $d \leq 3$, $N = 2$ with $d \leq 5$, and $N = 1$ with any $d \geq 1$. In particular, an ordinary double point of any dimension satisfies the assumptions of Lemma 2.5 in any odd characteristic. We also remark that in this case (i.e., when $d = 2$), the proof of Lemma 2.5 (5) shows that the space of graded k -linear maps $\wedge^n L_{A/k}[-n] \rightarrow \wedge^j L_{A/k}[-j][1]$ is simply connected.

Remark 2.7. Lemma 2.5 (5) only refers to space of graded k -linear maps $\wedge^n L_{A/k}[-n] \rightarrow \wedge^j L_{A/k}[-j][1]$, and not the space of such graded A -linear maps. In particular, it can happen that a graded A -linear map $\wedge^n L_{A/k}[-n] \rightarrow \wedge^j L_{A/k}[-j][1]$ is nullhomotopic as a graded k -linear map, but not as an A -linear map.

Theorem 1.5 will be used to control on the mod p crystalline cohomology of an lci k -scheme. To lift these results to W , we will use the following base change isomorphism; see [BdJ11] for more details.

Lemma 2.8. *Let X be a finite type lci k -scheme. Then the W -complex $\mathrm{R}\Gamma_{\mathrm{crys}}(X/W)$ has finite amplitude, and there is a base change isomorphism*

$$\mathrm{R}\Gamma_{\mathrm{crys}}(X/W) \otimes_W k \simeq \mathrm{R}\Gamma_{\mathrm{crys}}(X/k).$$

Proof. By a Mayer-Vietoris argument, we immediately reduce to the case where $X = \mathrm{Spec}(A)$ is affine. In this case, let D be the p -adic completion of the divided power envelope of a surjection $P \rightarrow A$ from a finite type polynomial W -algebra P . Then $\mathrm{R}\Gamma_{\mathrm{crys}}(X/W)$ is computed by

$$\Omega_{P/W}^* \otimes_P D.$$

Since this complex has finite amplitude, the first claim is proven. Next, if $P_0 = P/p$, and D_0 is the divided power envelope of $P_0 \rightarrow A$, then $\mathrm{R}\Gamma_{\mathrm{crys}}(X/k)$ is computed by

$$\Omega_{P_0/k}^* \otimes_{P_0} D_0.$$

The claim now follows from the well-known fact that D is W -flat (since A is lci), and $D \otimes_W k \simeq D_0$. \square

3. EXAMPLES

We come to the main topic of this note: examples of singular proper lci k -varieties with large crystalline cohomology. In §3.1, using lifts of Frobenius, we show that certain singular proper varieties (such as nodal curves, or singular lci toric varieties) have infinitely generated crystalline cohomology. In §3.2, we show that a single ordinary double point (or worse) on an lci proper variety forces crystalline cohomology to be infinitely generated.

3.1. Frobenius-liftable examples. We start with a general proposition which informally says: a proper lci k -variety has large crystalline cohomology if it contains an isolated singular point whose étale local ring lifts to W_2 compatibly with Frobenius. Note that lci k -algebras always lift to W_2 , so this is really a condition on Frobenius.

Proposition 3.1. *Let X be proper lci k -scheme. Assume:*

- (1) *There is a closed point $x \in X$ that is an isolated singular point (but there could be other singularities on X).*
- (2) *There is a lift to W_2 of the Frobenius endomorphism of the henselian ring $\mathcal{O}_{X,x}^h$.*

Set $N = \dim_k(\mathfrak{m}_x/\mathfrak{m}_x^2)$. Then there exists an integer $0 < i \leq N$ such that:

- (1) *$H_{\mathrm{crys}}^N(X/k)$ is infinitely generated over k .*
- (2) *$H_{\mathrm{crys}}^{N-i}(X/k)$ is infinitely generated over k .*
- (3) *At least one of $H_{\mathrm{crys}}^{N+1}(X/W)[p]$ and $H_{\mathrm{crys}}^N(X/W)/p$ is infinitely generated over k .*
- (4) *At least one of $H_{\mathrm{crys}}^{N+1-i}(X/W)[p]$ and $H_{\mathrm{crys}}^{N-i}(X/W)/p$ is infinitely generated over k .*

If $\dim(\mathcal{O}_{X,x}) > 0$, then the integer i above can be chosen to be strictly less than N .

Proof. The desired integer i will be found in the proof of (2) below.

- (1) Consider the exact triangle

$$\mathrm{Fil}_N^{\mathrm{conj}}(\mathrm{dR}_{X/k}) \rightarrow \mathrm{dR}_{X/k} \rightarrow \mathcal{Q}$$

in the $\mathcal{O}_{X(1)}$ -complexes, where \mathcal{Q} is defined as the homotopy-cokernel. Theorem 1.5 (1) and the lci assumption on X show that $\mathrm{Fil}_N^{\mathrm{conj}}(\mathrm{dR}_{X/k})$ is a perfect complex on $X^{(1)}$, so $H^i(X^{(1)}, \mathrm{Fil}_N^{\mathrm{conj}}(\mathrm{dR}_{X/k}))$ is a finite dimensional vector space for all i by properness. By Theorem 1.5 (3), to show that $H_{\mathrm{crys}}^N(X/k)$ is infinitely generated, it suffices to show that $H^N(X^{(1)}, \mathcal{Q})$ is an infinite dimensional k vector space. First, we show:

Claim 3.2. *The natural map $R\Gamma(X, \mathcal{Q}) \rightarrow \mathcal{Q}_x$ is a projection onto a summand as k -complexes.*

Proof of Claim. Let $j : U \rightarrow X$ be an affine open neighbourhood of x such that U has an isolated singularity at x , and let $j' : V = X - \{x\} \subset X$. By Theorem 1.5 (2), $\mathcal{Q}|_{U \cap V} \simeq 0$ since $U \cap V$ is k -smooth. Hence, the Mayer-Vietoris sequence for the cover $\{U, V\}$ of X and the complex \mathcal{Q} degenerates to show

$$R\Gamma(X, \mathcal{Q}) \simeq R\Gamma(U, \mathcal{Q}) \oplus R\Gamma(V, \mathcal{Q}).$$

It now suffices to show that $R\Gamma(U, \mathcal{Q}) \simeq \mathcal{Q}_x$. By Theorem 1.5 (1), $\mathcal{Q}|_U$ admits an increasing bounded below separated exhaustive filtration with graded pieces $\wedge^n L_{U/k}[-k]$ for $n > N$. Since cohomology commutes with filtered colimits (as U is affine), $R\Gamma(U, \mathcal{Q})$ also inherits such a filtration with graded pieces computed by $R\Gamma(U, \wedge^n L_{U^{(1)}/k}[-k])$ for $n > N$. Applying the same analysis to \mathcal{Q}_x reduces us to checking that $R\Gamma(U, \wedge^n L_{U^{(1)}/k}[-n]) \simeq \wedge^n L_{\mathcal{O}_{X,x}^{(1)}/k}[-n]$ for $n > N$. But this is clear: for $n > N$, $\wedge^n L_{U^{(1)}/k}[-n]$ is a perfect complex on $U^{(1)}$ that is supported only at x and has stalk $\wedge^n L_{\mathcal{O}_{X,x}^{(1)}/k}[-n]$. \square

To compute the stalk \mathcal{Q}_x , define \mathcal{Q}' via the exact triangle

$$\mathrm{Fil}_N^{\mathrm{conj}}(\mathrm{dR}_{\mathcal{O}_{X,x}^h/k}) \rightarrow \mathrm{dR}_{\mathcal{O}_{X,x}^h/k} \rightarrow \mathcal{Q}'.$$

Then $\mathcal{Q}_x = \mathcal{Q}'$ by Theorem 1.5 (2), the finite length property of \mathcal{Q}_x , and the fact that $\mathcal{O}_{X,x}^h \otimes_{\mathcal{O}_{X,x}} M \simeq M$ for any finite length $\mathcal{O}_{X,x}$ -module M . Moreover, \mathcal{Q}' can be computed using the Frobenius lifting assumption and Theorem 1.5 (4):

$$\mathcal{Q}_x \simeq \mathcal{Q}' \simeq \bigoplus_{n=N+1}^{\infty} \wedge^n L_{\mathcal{O}_{X,x}^{(1),h}/k}[-n].$$

Thus, to prove that $H^N(X^{(1)}, \mathcal{Q})$ is infinitely generated, it suffices to show that $H^N(\mathcal{Q}_x)$ is infinitely generated. This follows from the formula above and Lemma 2.1 (3).

- (2) By combining the proof of (1) with Lemma 2.1 (4) and the pigeonhole principle, one immediately finds an integer $0 < i \leq N$ such that $H_{\mathrm{crys}}^{N-i}(X/k)$ is infinitely generated over k . Lemma 2.1 (5) shows that we can choose such an i with $i < N$ if $\dim(\mathcal{O}_{X,x}) > 0$.
- (3) The base change isomorphism from Lemma 2.8 gives a short exact sequence

$$0 \rightarrow H_{\mathrm{crys}}^N(X/W)/p \rightarrow H_{\mathrm{crys}}^N(X/k) \rightarrow H_{\mathrm{crys}}^{N+1}(X/W)[p] \rightarrow 0,$$

so the claim follows from (1).

- (4) The same argument as (3) works using (2) instead of (1). \square

We need the following elementary result on Frobenius liftings:

Lemma 3.3. *Let A be a k -algebra that admits a lift to W_2 together with a compatible lift of Frobenius. Then the same is true for any ind-étale A -algebra B (such as the henselisation A at a point).*

Proof. This follows by deformation theory since $L_{B/A} = 0$ for B as above. \square

Specialising Proposition 3.1 leads to the promised examples.

Example 3.4. Let $X = \mathrm{Spec}(k[x]/x^n)$ for some $n > 1$. Then $H_{\mathrm{crys}}^1(X/k)$, $H_{\mathrm{crys}}^0(X/k)$, $H_{\mathrm{crys}}^1(X/W)/p$, and $H_{\mathrm{crys}}^1(X/W)[p]$ are all infinitely generated. To see this, note first that Proposition 3.1 applies directly since X is a proper lci k -scheme with a lift of Frobenius to W_2 . Moreover, since X can be realised as a subscheme of \mathbf{A}^1 , the only non-zero cohomology groups are H_{crys}^1 and H_{crys}^0 (over W , as well as over k). The rest follows directly from Proposition 3.1 and the observation that $H_{\mathrm{crys}}^0(X/W) = W$ (which is true for any lci k -scheme).

Example 3.5. Let X be a proper nodal k -curve with at least one node. Then $H_{\mathrm{crys}}^1(X/k)$ and $H_{\mathrm{crys}}^2(X/k)$ are infinitely generated. Moreover, $H_{\mathrm{crys}}^2(X/W)/p$, and at least one of $H_{\mathrm{crys}}^1(X/W)/p$ and $H_{\mathrm{crys}}^2(X/W)[p]$, are infinitely generated. Most of these claims follow directly from Proposition 3.1: a nodal curve is always lci, and the henselian local ring at a node on X is isomorphic to the henselisation of $k[x, y]/(xy)$ at the origin, which is a one-dimensional local ring that admits a lift to W_2 compatible with Frobenius by Lemma 3.3. It remains to show that $H_{\mathrm{crys}}^3(X/W)[p]$ is finitely generated. As pointed out by de Jong, the stronger statement $H_{\mathrm{crys}}^3(X/W) = 0$ is true. If $u : (X/W)_{\mathrm{crys}} \rightarrow X_{\mathrm{Zar}}$ is the natural map (i.e., $u_*(\mathcal{F})(U \subset X) = \Gamma((U/W)_{\mathrm{crys}}, \mathcal{F}|_U)$), then $R^i u_* \mathcal{O}_{X/W, \mathrm{crys}}$ is non-zero only for $0 \leq i \leq 2$,

and $R^2 u_* \mathcal{O}_{X/W, \text{crys}}$ is supported only at the nodes¹. The rest follows from the Leray spectral sequence as X_{Zar} has cohomological dimension 1.

Example 3.6. Let X be a proper lci k -scheme. Assume that $x \in X(k)$ is an isolated singular point (but there could be other singularities on X) such that $\mathcal{O}_{X,x}^h$ is toric of embedding dimension N . Then $H_{\text{crys}}^N(X/k)$ is infinitely generated, and at least one of $H_{\text{crys}}^N(X/W)$ and $H_{\text{crys}}^{N+1}(X/W)[p]$ is infinitely generated over W . This follows from Proposition 3.1 and Lemma 3.3 since toric rings lift to W_2 compatibly with Frobenius (use multiplication by p on the defining monoid). Some specific examples are: any proper toric variety with isolated lci singularities, or any proper singular k -scheme of dimension ≤ 3 with at worst ordinary double points.

Example 3.7. Let (E, e) be an ordinary elliptic curve over k , and let X be a proper lci k -surface with a singularity at $x \in X(k)$ isomorphic to the one on the affine cone over $E \subset \mathbf{P}_k^2$ embedded via $\mathcal{O}(3[e])$; for example, we could take X to be the projective cone on $E \subset \mathbf{P}_k^2$. Then $H_{\text{crys}}^3(X/k)$ and one of $H_{\text{crys}}^2(X/k)$ or $H_{\text{crys}}^1(X/k)$ are infinitely generated over k . This can be proven using Proposition 3.1 and the theory of Serre-Tate canonical lifts. Since we prove a more general and shaper result in Example 3.13, we leave details of this argument to the reader.

3.2. Conical examples. Our goal here is to show that the presence of an ordinary double point forces crystalline cohomology to be infinitely generated. In fact, more generally, we show the same for any proper lci variety that has a singularity isomorphic to the cone on a low degree smooth hypersurface. We start with an *ad hoc* definition.

Definition 3.8. A local k -algebra A is called a *low degree cone* if its henselisation is isomorphic to the henselisation at the origin of the ring $k[x_0, \dots, x_N]/(f)$, where f is a homogeneous degree d polynomial defining a smooth hypersurface in \mathbf{P}^N such that $N(d-2) < d+2$. The integer d is called the *degree* of this cone; if $d=2$, we also call A an *ordinary double point*. A closed point $x \in X$ on a finite type k -scheme X is called *low degree conical singularity* (respectively, an *ordinary double point*) if $\mathcal{O}_{X,x}^h$ is a low degree cone (respectively, an ordinary double point).

We start by showing that the conjugate spectral sequence must eventually degenerate for low degree cones:

Proposition 3.9. *Let A be low degree cone of degree d . Assume $p \nmid d$. Then for $n > \dim(A)$, the extensions*

$$\text{gr}_n^{\text{conj}}(\text{dR}_{A/k}) \rightarrow \text{Fil}_{n-1}^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k})[1]$$

occurring in the conjugate filtration are nullhomotopic when viewed as k -linear extensions. In particular, there exist k -linear isomorphisms

$$\text{Fil}_n^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k}) \simeq \bigoplus_{j=\dim(A)+1}^n \text{gr}_j^{\text{conj}}(\text{dR}_{A/k}).$$

splitting the conjugate filtration for any $n > \dim(A)$.

Proof. By replacing A with its henselisation and then using the étale invariance of cotangent complexes and Theorem 1.5 (2), we may assume A is the localisation of $k[x_0, \dots, x_N]/(f)$ at the origin for some homogeneous degree d polynomial f defining a smooth hypersurface in \mathbf{P}^N . In particular, A is graded. Also, by functoriality, the conjugate filtration is compatible with the grading. Recall that the extensions in question arise from the triangles

$$\text{Fil}_{n-1}^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k}) \rightarrow \text{Fil}_n^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k}) \rightarrow \text{gr}_n^{\text{conj}}(\text{dR}_{A/k}).$$

These triangles (and thus the corresponding extensions) are viewed as living in the derived category of graded k -vector spaces. By induction, we have to show the following: assuming a graded splitting

$$s_{n-1} : \text{Fil}_{n-1}^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k}) \simeq \bigoplus_{j=\dim(A)+1}^{n-1} \text{gr}_j^{\text{conj}}(\text{dR}_{A/k})$$

of the conjugate filtration, there exists a graded splitting

$$s_n : \text{Fil}_n^{\text{conj}}(\text{dR}_{A/k})/\text{Fil}_{\dim(A)}^{\text{conj}}(\text{dR}_{A/k}) \simeq \bigoplus_{j=\dim(A)+1}^n \text{gr}_j^{\text{conj}}(\text{dR}_{A/k}),$$

of the conjugate filtration compatible with s_{n-1} . Chasing extensions, it suffices to show: for $\dim(A) < j < n$, all graded maps

$$\text{gr}_n^{\text{conj}}(\text{dR}_{A/k}) \rightarrow \text{gr}_j^{\text{conj}}(\text{dR}_{A/k})[1]$$

are nullhomotopic. This comes from Lemma 2.5 (5) and Theorem 1.5 (1). \square

¹Proof sketch: Replace the Zariski topology with the Nisnevich topology in the foundations of crystalline cohomology, and then use that every nodal curve is Nisnevich locally planar. This observation yields a three-term de Rham complex computing the stalks of $R^i u_* \mathcal{O}_{X/W, \text{crys}}$.

Remark 3.10. An inspection of the proof of Proposition 3.9 coupled with Remark 2.6 shows that if A is an ordinary double point, then the isomorphism

$$\mathrm{Fil}_n^{\mathrm{conj}}(\mathrm{dR}_{A/k})/\mathrm{Fil}_{\dim(A)}^{\mathrm{conj}}(\mathrm{dR}_{A/k}) \simeq \bigoplus_{j=\dim(A)+1}^n \mathrm{gr}_j^{\mathrm{conj}}(\mathrm{dR}_{A/k})$$

is unique, up to non-unique homotopy. We do not know any applications of this uniqueness.

Using Proposition 3.9, we can prove infiniteness of crystalline cohomology for some cones:

Corollary 3.11. *Let X be a proper lci k -scheme. Assume that there is low degree conical singularity at a closed point $x \in X$ with degree d and embedding dimension N . If $p \nmid d$, then $H_{\mathrm{crys}}^N(X/k)$ and $H_{\mathrm{crys}}^{N-1}(X/k)$ are infinitely generated k -vector spaces.*

Proof. We combine the proof strategy of Proposition 3.1 with Proposition 3.9. More precisely, following the proof of Proposition 3.1 (1), it suffices to show that \mathcal{Q}' is infinitely generated when regarded as a complex of k -vector spaces. Now \mathcal{Q}' admits an increasing bounded below separated exhaustive filtration with graded pieces given by $\mathrm{gr}_n^{\mathrm{conj}}(\mathrm{dR}_{\mathcal{O}_{X,x}^h/k})$ for $n > N$. By Proposition 3.9, there is a (non-canonical) isomorphism

$$\mathcal{Q}' \simeq \bigoplus_{n>N} \wedge^n L_{\mathcal{O}_{X,x}^{(1),h}/k}[-n].$$

The rest follows from Lemma 2.5 (4) (note that embedding dimension in *loc. cit.* is $N+1$, so we must shift by 1). \square

We can now give the promised example:

Example 3.12. Let X be any proper lci variety that contains an ordinary double point $x \in X(k)$ of embedding dimension N with p odd; for example, we could take X to be the projective cone over a smooth quadric in \mathbf{P}^{N-1} . Then $H_{\mathrm{crys}}^N(X/k)$ and $H_{\mathrm{crys}}^{N-1}(X/k)$ are infinitely generated by Corollary 3.11.

All examples given so far have rational singularities, so we record an example that is not even log canonical.

Example 3.13. Let X be any proper lci surface that contains a closed point $x \in X(k)$ with $\mathcal{O}_{X,x}^h$ isomorphic to the henselisation at the vertex of the cone over a smooth curve $C \subset \mathbf{P}^2$ of degree ≤ 5 . If $p \geq 7$, then $H_{\mathrm{crys}}^3(X/k)$ and $H_{\mathrm{crys}}^2(X/k)$ are infinitely generated by Corollary 3.11.

Remark 3.14. We do not know whether the ordinary double point from Example 3.12 admits a lift to W_2 compatible with Frobenius in arbitrary dimensions; similarly for the cones in Example 3.13 (except for ordinary elliptic curves).

We end with a question that has not been answered in this note.

Question 3.15. Do there exist singular proper k -varieties with finite dimensional crystalline cohomology over k ? Do there exist singular finite type k -algebras A whose crystalline cohomology relative to k is finitely generated over $A^{(1)}$?

REFERENCES

- [BdJ11] Bhargav Bhatt and Aise Johan de Jong. Crystalline cohomology and de Rham cohomology. 2011. Available at <http://arxiv.org/abs/1110.5001>.
- [Ber74] Pierre Berthelot. *Cohomologie cristalline des schémas de caractéristique $p > 0$* . Lecture Notes in Mathematics, Vol. 407. Springer-Verlag, Berlin, 1974.
- [Bha12] Bhargav Bhatt. p -adic derived de Rham cohomology. 2012. Available at <http://arxiv.org/abs/1204.6560>.
- [BO78] Pierre Berthelot and Arthur Ogus. *Notes on crystalline cohomology*. Princeton University Press, Princeton, N.J., 1978.
- [BO83] Pierre Berthelot and Arthur Ogus. F -isocrystals and de Rham cohomology. I. *Invent. Math.*, 72(2):159–199, 1983.
- [DI87] Pierre Deligne and Luc Illusie. Relèvements modulo p^2 et décomposition du complexe de de Rham. *Invent. Math.*, 89(2):247–270, 1987.
- [Gro68] Alexander Grothendieck. Crystals and the de Rham cohomology of schemes. In *Dix Exposés sur la Cohomologie des Schémas*, pages 306–358. North-Holland, Amsterdam, 1968.
- [Ill72] Luc Illusie. *Complexe cotangent et déformations. II*. Lecture Notes in Mathematics, Vol. 283. Springer-Verlag, Berlin, 1972.
- [Ked06] Kiran S. Kedlaya. Fourier transforms and p -adic ‘Weil II’. *Compos. Math.*, 142(6):1426–1450, 2006.
- [KM74] Nicholas Katz and William Messing. Some consequences of the Riemann hypothesis for varieties over finite fields. *Invent. Math.*, 23:73–77, 1974.
- [KS04] Kazuya Kato and Takeshi Saito. On the conductor formula of Bloch. *Publ. Math. Inst. Hautes Études Sci.*, (100):5–151, 2004.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur11] Jacob Lurie. Higher Algebra. Available at <http://math.harvard.edu/~lurie>, 2011.
- [Qui70] Daniel Quillen. Homology of commutative rings. Unpublished notes from an M.I.T. course., 1970.
- [Voi07] Claire Voisin. *Hodge theory and complex algebraic geometry. II*, volume 77 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, english edition, 2007. Translated from the French by Leila Schneps.