

## Viewpoints

# How the Change from FLM to FACS Affected Our Understanding of the G<sub>1</sub>-Phase of the Cell Cycle

Stephen Cooper

Correspondence to: Stephen Cooper, Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor Michigan 48109-0620 USA; Tel.: 734.764.4215; Fax: 734.764.3562; Email: cooper@umich.edu; Website: www.umich.edu/~cooper

Received 01/14/03; Accepted 01/17/03

Previously published online as a *Cell Cycle* "Paper In Press" at <http://www.landesbioscience.com/journals/cc/>

### KEY WORDS

Cell cycle, Flow assisted cell sorting, Frequency of Labeled Mitoses, G<sub>1</sub>-phase, G<sub>0</sub>

### ABSTRACT

The frequency of labeled mitoses (FLM) method for analyzing cell-cycle phases necessitates a determination of cell-cycle interdivision times and the absolute lengths of the cell-cycle phases. The change to flow sorting (FACS) analysis, a simpler, less labor intensive, and more rapid method, eliminated determinations of absolute phase times, yielding only percents of cells exhibiting particular DNA contents. Without an interdivision time value, conversion of these fractions into absolute phase lengths is not possible. This change in methodology has led to an alteration in how the cell cycle is viewed. The FLM method allowed the conclusion that G<sub>1</sub>-phase variability resulted from constancy of S and G<sub>2</sub> phase lengths. In contrast, with FACS analysis, slow growing cells exhibiting a large fraction of cells with a G<sub>1</sub>-phase amount of DNA appeared to be "arrested in G<sub>1</sub> phase". The loss of absolute phase length determinations has therefore led to the proposals of G<sub>1</sub>-phase arrest, G<sub>1</sub>-phase controls, restriction points, and G<sub>0</sub> phase. It is suggested that these G<sub>1</sub>-phase controls and phenomena require a critical reevaluation in the light of an alternative cell-cycle model that does not require or postulate such G<sub>1</sub>-phase controls.

After the original identification of cell-cycle phases by Howard and Pelc in the early 1950s, the next two decades brought forth a large number of measurements of cell-cycle phase lengths using the method of frequency of labeled mitoses (FLM). FLM is essentially the method used by Howard and Pelc.<sup>1,2</sup>

The FLM method is illustrated in Figure 1. Because this method is not now in common use, it is of interest to explain the method and the type of results obtained by this method. For FLM, growing cells are pulse-labeled with tritiated thymidine. After removal of label, growing cells are sampled at intervals and subjected to autoradiography. The fraction of mitotic cells that are radioactive is determined. As shown in Figure 1, there is an initial period with no labeled mitoses. This period gives the length, in absolute time, of the G<sub>2</sub> phase. Then there is a rise in labeled mitoses, a plateau, and a fall in labeled mitoses. The time interval of these labeled mitoses indicates the S-phase length. The G<sub>1</sub> phase length is determined from the G<sub>1</sub> + G<sub>2</sub> period—the period between the decrease in labeled mitoses and the subsequent increase in labeled mitoses—by subtracting the initially determined G<sub>2</sub> phase time. The total interdivision time (IDT) is determined by the time between the rise times of labeled mitoses.

The major generalization to come out of a large number of FLM analyses in the 1950–1970 period was that the G<sub>1</sub>-phase is the most variable cell-cycle phase. In contrast, S and G<sub>2</sub> phases were relatively constant. Some studies demonstrated not only a large variability of the G<sub>1</sub> phase, but also a definite constancy of S and G<sub>2</sub> phases as interdivision time was varied. This observation of constant S and G<sub>2</sub> phases led to the postulation of an analogy between the bacterial and eukaryotic models for DNA replication.<sup>3</sup> In bacteria the constant C and D periods are analogous to the constant S and G<sub>2</sub> phases.

Examples of idealized FLM results are presented in Table 1. Consider a given cell type growing in four different growth conditions (A, B, C, D), each with a different interdivision time. If S and G<sub>2</sub> are invariant, then G<sub>1</sub> phase length is necessarily the difference between the interdivision time and the S phase length plus G<sub>2</sub> phase length. That is,  $G_1 = IDT - (S + G_2)$ . These results (Table 1) led to the conclusion that the G<sub>1</sub> phase length was determined by the overall growth rate or interdivision time. This reasoning led to a model of cell-cycle control that did not invoke any particular or unique events occurring in the G<sub>1</sub>-phase.<sup>4-8</sup>




Figure 1 (left). Frequency of Labeled Mitoses (FLM) method for determination of cell-cycle phase lengths. To determine the cell-cycle phases using FLM, exponentially growing cells are pulse-labeled for a short period with tritiated thymidine. Only cells in S phase incorporate thymidine label. After removal of the exogenous label, cells are removed at intervals and analyzed by autoradiography. The fraction of mitotic cells with labeled DNA is determined. At the beginning of collection those cells in the G<sub>2</sub> phase at the time of pulse labeling produce dividing or mitotic cells over a time equivalent to the G<sub>2</sub> phase; there are no labeled mitoses during this initial period. After a period of time equal to the G<sub>2</sub> phase length, cells that were in S phase at the time of labeling now enter mitosis and one now sees labeled mitoses.

There are labeled mitoses for a period of time equal to the length of an S phase. Then cells that were in the G<sub>1</sub> phase at the time of labeling produce mitotic cells and these are unlabeled. The period of unlabeled mitoses continues for a time equal to the sum of the G<sub>1</sub> phase and the G<sub>2</sub> phase. This is because the cells that were originally in G<sub>2</sub> phase at the time of pulse labeling now divide again. After the second G<sub>2</sub> phase period, cells again produce labeled mitoses. The length of the G<sub>1</sub> phase is determined by subtracting the G<sub>2</sub> phase length from the G<sub>2</sub>+G<sub>1</sub> phase length. The total interdivision time (IDT) can be determined by the time from the first rise in labeled mitoses to the second rise in labeled mitoses. As shown here, the G<sub>1</sub>, S, and G<sub>2</sub> phases are 50%, 25%, and 25% of the total IDT.

The wide-spread use of FLM, a very labor-intensive method, was eventually supplanted by FACS (fluorescence assisted cell sorting) analysis, a simpler, less labor-intensive, and more rapid method. In FACS analysis, cells are merely sampled from a growing cell culture and the total fraction of cells with different DNA contents is determined. An example of idealized results obtained by FACS analysis is presented in Figure 2. As seen in Figure 2, FACS analysis gives the fraction of cells with a G<sub>1</sub>, S, or G<sub>2</sub>/M phase amount of DNA. The absolute lengths of these phases are not given by FACS analysis. Determination of phase lengths requires knowledge of the total interdivision time. Multiplying the total interdivision time by the fractional amount of cells in each phase gives the absolute phase lengths. (The age distribution has been ignored in this simplified analysis.)

Idealized results for the same cultures (A, B, C, and D) as in Table 1 are presented in Table 2 as determined by FACS analysis. The absence of interdivision times leads to the description of the cultures by the conditions used to produce the different cell-cycle patterns. The data of Table 2 are actually the same as the data in Table 1, but the results are given as percentages of the cells in different phases. As the serum concentration is decreased (a classic method for cell growth "arrest") more cells have a G<sub>1</sub>-phase amount of DNA. Without knowing the interdivision times, it has been generally concluded that the increase in cells with a G<sub>1</sub>-phase amount of DNA indicates that the cells are arrested at a particular point or control element in the G<sub>1</sub> phase.

FACS analysis therefore gives a very different picture of the cell cycle than FLM methodology. The constancy of S and G<sub>2</sub> phases is not apparent with FACS analysis. The dominant result with FACS analysis is the increase in the fraction of cells with a G<sub>1</sub>-phase amount of DNA.

The results in Table 2 for slow growing cells (low, lower, or very low serum) have been interpreted as indicating arrest of cells in the G<sub>1</sub> phase or arrest "at a point in G<sub>1</sub> phase" (e.g., the restriction point), or entrance of the cells into an "out-of-cycle-phase", G<sub>0</sub>. But it is important to emphasize that it is not postulated that the cells analyzed in Tables 1 and 2 are arrested. The cells described in Tables




Figure 2. Flow-cytometric (FACS) determination of fraction of cells with particular DNA contents. For FACS analysis, exponentially growing cells are labeled with a fluorophore that reflects the DNA content of cells. As each cell is assayed, the DNA contents are measured. As shown here, for cells with 50% G<sub>1</sub> phase cells, 25% S phase cells, and 25% G<sub>2</sub> phase cells, one can measure the total number of cells in the specific portions of the DNA contents. (N. B. For simplification, no account has been made of the age distribution of cells.)

Table 1 **FLM ANALYSIS OF CELL CYCLE**

	Interdivision Time	G <sub>1</sub> <sup>a</sup>	S <sup>a</sup>	G <sub>2</sub> <sup>a</sup>
A	20 hours	10	5	5
B	100 hours	90	5	5
C	1,000 hours	990	5	5
D	10,000 hours	9,990	5	5

<sup>a</sup>Phase lengths in hours

Table 2 **FACS ANALYSIS OF CELL CYCLE**

	Treatment	G <sub>1</sub> <sup>a</sup>	S <sup>a</sup>	G <sub>2</sub> <sup>a</sup>
A	Full serum	50	25	25
B	Low serum	90	5	5
C	Lower serum	99	0.5	0.5
D	Very low serum	99.9	0.05	0.05

<sup>a</sup>Percent of cells observed with specific DNA contents.

1 and 2 are assumed to be growing steadily but at different rates of growth. As shown in Table 1, the cells are merely growing slowly as the serum is reduced (described in Table 2). The imputation of cell “arrest” is merely due to the experimental difficulty of following cell growth over many hours and determining whether or not cells are growing slowly.

To be precise, the concepts of G<sub>0</sub> and the restriction point and G<sub>1</sub>-phase arrest points pre-dated the development of FACS analysis. What has happened, however, is that the facile use of this instrumentation has led to appearance of a large amount of support for these G<sub>1</sub>-phase control elements. The point made here is that the ubiquity of FACS analysis has made it appear as though a large amount of experimental evidence independently supports these G<sub>1</sub>-phase arrest phenomena. As shown in Tables 1 and 2, this is not the case.

We see that the advent of the newer, simpler, more rapid, and quantitative method, FACS analysis, displacing the FLM method, has led to omission of a major point, the constancy and relative invariance of the S and G<sub>2</sub> phases. Thus, FACS analysis can give an incorrect view of the cell cycle. As shown for bacteria, the constancy of DNA replication time (S phase in eukaryotes, C period in bacteria) and the constancy of the period between termination of DNA replication and cell division (G<sub>2</sub> in eukaryotes, D period in bacteria) leads to a model of cell-cycle regulation that does not postulate the importance of passage through G<sub>1</sub>-phase specific events, restriction points, or checkpoints. The experimental and theoretical support for this view of cell cycle control has been reviewed.<sup>4-7</sup>

Instead of envisioning specific G<sub>1</sub>-phase controls, this alternative model proposes that control of cell-cycle events is based on a continuous accumulation occurring in all phases of the cell cycle. The variable G<sub>1</sub> phase is a consequence of variation in growth rate throughout the cell cycle. The variable G<sub>1</sub> phase is not a consequence of how rapidly or slowly specific G<sub>1</sub>-phase events are carried out. FACS analysis, in contrast to FLM analysis, leads to the idea of G<sub>1</sub>-phase arrest, G<sub>1</sub>-phase control elements, and other G<sub>1</sub>-phase phenomena. The current model of the cell cycle should therefore be reevaluated and reconsidered in the light of this historical analysis.

### References

1. Howard A, Pelc S. Nuclear incorporation of P32 as demonstrated by autoradiographs. *Exp Cell Res* 1951; 2:178-87.
2. Howard A, Pelc S. Synthesis of nucleoprotein in bean root cells. *Nature* 1951; 167:599-600.
3. Helmstetter C, Cooper S, Pierucci O, Revelas E. On the bacterial life sequence. *Cold Spring Harb Symp Quant Biol* 1968; 33:809-22.
4. Cooper S. Rethinking synchronization of mammalian cells for cell-cycle analysis. *Cellular and Molecular Life Sciences* 2003; Accepted for publication.
5. Cooper S. Reappraisal of serum starvation, the restriction point, G<sub>0</sub>, and G<sub>1</sub>-phase arrest points. *FASEB J* 2003; 17:333-40.
6. Cooper S, Shayman JA. Revisiting retinoblastoma protein phosphorylation during the mammalian cell cycle. *Cell Mol Life Sci*. Apr 2001; 58:580-95.
7. Cooper S. The continuum model and G<sub>1</sub>-control of the mammalian cell cycle. *Prog Cell Cycle Res* 2000; 4:27-39.
8. Cooper S. On the interpretation of the shortening of the G<sub>1</sub>-phase by overexpression of cyclins in mammalian cells. *Exp Cell Res* 1998; 238:110-15.