

Jackson Theorems for Erdős Weights in $L_p(0 < p \leq \infty)$

S. B. Damelin and D. S. Lubinsky

Department of Mathematics, Witwatersrand University,
P.O. Wits 2050, South Africa

e-mail: 036SBD@COSMOS.WITS.AC.ZA and 036DSL@COSMOS.WITS.AC.ZA

February 4, 2000

Abstract

An *Erdős weight* is of the form $W := e^{-Q}$ where Q is even and of faster than polynomial growth at ∞ . For example, we can take

$$Q(x) := \exp_k(|x|^\alpha), \quad k \geq 1, \alpha > 0, \quad x \in \mathbb{R}$$

where \exp_k denotes the k th iterated exponential. We prove Jackson theorems in weighted L_p spaces with norm $\|fW\|_{L_p(\mathbb{R})}$ for all $0 < p \leq \infty$. These are the first proper Jackson theorems for Erdős weights even in L_∞ . An interesting feature is a Timan/Nikolskii/Brudnyi effect: The degree of approximation improves towards the endpoints of a certain interval. By contrast, there is no such feature for Freud weights.

AMS(MOS) Classification: Primary 41A10, 42C05

Keywords: Erdős weights, Jackson theorems, polynomial approximation

1 Statement of Results

In recent years, there have been many advances in the theory of weighted polynomial approximation, and orthonormal polynomials, associated with the weights

$$W := e^{-Q}. \quad (1.1)$$

Here $Q : \mathbb{R} \rightarrow \mathbb{R}$ is even, and typically grows at least as fast as $|x|^\lambda$, some $\lambda > 1$, at infinity. In some contexts, there has been a distinction between the case where Q is of polynomial growth at infinity (the so-called *Freud case*) and where Q is of faster than polynomial growth at infinity (the so-called *Erdős case*). To some extent, this is similar to the distinction between entire functions of finite, and infinite, order. For further orientation on this topic, see [7], [10], [15], [16], [21], [22].

In this paper, we discuss Jackson theorems for Erdős weights. That is, we estimate

$$E_n[f]_{W,p} := \inf_{P \in P_n} \|(f - P)W\|_{L_p(\mathbb{R})}, \quad (1.2)$$

$0 < p \leq \infty$, where P_n denote the polynomials of degree at most n .

Our methods are similar to those in [6], where Jackson theorems were proved for Freud weights. The approach involves approximating f by a spline (or piecewise polynomial), representing the piecewise polynomial in terms of certain characteristic functions, and then approximating the characteristic functions (in a suitable sense) by polynomials. This method has the advantage of involving only hypotheses on Q' , in contrast with the more complicated approach via orthogonal polynomials and de la Vallée Poussin sums, that typically involves hypotheses on Q'' [7], [10], [17], [21]. In the Erdős weight context, some new features arise: the degree of approximation improves toward the endpoints of the Mhaskar-Saff interval, and to reflect this, we need a more complicated modulus of continuity, and some proofs become more involved.

To state our result, we need to define our class of weights, as well as various quantities. First, we say that a function $f : (a, b) \rightarrow (0, \infty)$ is *quasi-increasing* if $\exists C > 0$ such that

$$a < x < y < b \implies f(x) \leq Cf(y).$$

Definition 1.1 Let $W := e^{-Q}$, where

(a) $Q : \mathbb{R} \rightarrow \mathbb{R}$ is even, continuous, and Q' is positive in $(0, \infty)$.

(b) $xQ'(x)$ is strictly increasing in $(0, \infty)$ with right limit 0 at 0.

(c) The function

$$T(x) := \frac{xQ'(x)}{Q(x)} \quad (1.3)$$

is quasi-increasing in (C, ∞) for some $C > 0$, and

$$\lim_{x \rightarrow \infty} T(x) = \infty. \quad (1.4)$$

(d) $\exists C_1, C_2, C_3 > 0$ such that

$$\frac{yQ'(y)}{xQ'(x)} \leq C_1 \left(\frac{Q(y)}{Q(x)} \right)^{C_2}, \quad y \geq x \geq C_3. \quad (1.5)$$

Then we write $W = e^{-Q} \in \mathcal{E}_1$.

The archetypal example of $W \in \mathcal{E}_1$ is

$$W(x) := W_{k,\alpha}(x) := \exp(-\exp_k(|x|^\alpha)), \quad k \geq 1, \alpha > 0, \quad (1.6)$$

where $\exp_k = \exp(\exp(\dots))$ denotes the k th iterated exponential. For this weight, we see

$$T(x) = \alpha x^\alpha \prod_{j=1}^{k-1} \exp_j(x^\alpha), \quad x > 0.$$

It is not too difficult to see that we can choose $C_2 > 1$ in (1.5) arbitrarily close to 1 in this case. Another example is

$$W(x) := \exp\left(-\exp[\log(2+x^2)]^\beta\right), \quad \beta > 1.$$

Here

$$T(x) = \frac{2\beta x^2}{2+x^2} [\log(2+x^2)]^{\beta-1}, \quad x > 0.$$

Again, we can choose C_2 arbitrarily close to 1.

The function $T(x)$ measures the regularity of growth of $Q(x)$. In particular, (1.4) forces Q to be of faster than polynomial growth at ∞ . The reader is cautioned that in other papers on Erdős weights [14], [17] the function

$$T_1(x) := 1 + \frac{xQ''(x)}{Q'(x)}$$

was used (and denoted by T), but it has essentially the same rate of growth as T , for “nice” weights.

We need the condition that $xQ'(x)$ be strictly increasing to guarantee the existence of the *Mhaskar-Rakhmanov-Saff number* a_u , the positive root of the equation

$$u = \frac{2}{\pi} \int_0^1 a_u t Q'(a_u t) \frac{dt}{\sqrt{1-t^2}}, \quad u > 0. \quad (1.7)$$

If we used something other than a_u , we could require less of $xQ'(x)$, namely that it be quasi-increasing for large x . However this would complicate formulations, so is omitted. For those to whom a_u is new, its significance lies partly in the identity [18], [19], [20]

$$\|PW\|_{L_\infty(\mathbb{R})} = \|PW\|_{L_\infty[-a_n, a_n]}, \quad P \in \mathcal{P}_n, \quad (1.8)$$

and that a_n is the “smallest” such number.

Our modulus of continuity involves two parts, a “main part” and a “tail”. The “main part” involves r th symmetric differences over a suitable interval, and the tail involves an error of weighted polynomial approximation over the remainder of the real line. The size of this “suitable interval” is determined by the decreasing function of t ,

$$\sigma(t) := \inf \left\{ a_u : \frac{a_u}{u} \leq t \right\}, \quad t > 0. \quad (1.9)$$

Thus σ is essentially the inverse function of the function $u \rightarrow \frac{a_u}{u}$, which decays to 0 as $u \rightarrow \infty$.

For $h > 0$, an interval J , and $r \geq 1$, we define the r th symmetric difference

$$\Delta_h^r(f, x, J) := \sum_{i=0}^r \binom{r}{i} (-1)^i f \left(x + \frac{rh}{2} - ih \right), \quad (1.10)$$

provided all arguments of f lie in J , and 0 otherwise. Sometimes the increment h will depend on x , and on the function

$$\Phi_t(x) := \sqrt{\left|1 - \frac{|x|}{\sigma(t)}\right|} + T(\sigma(t))^{-1/2}, \quad x \in \mathbb{R}. \quad (1.11)$$

This is the case in our modulus of continuity

$$\begin{aligned} \omega_{r,p}(f, W, t) : &= \sup_{0 < h \leq t} \|W \Delta_{h\Phi_t(x)}^r(f, x, \mathbb{R})\|_{L_p(|x| \leq \sigma(2t))} \\ &+ \inf_{P \in \mathcal{P}_{r-1}} \|(f - P)W\|_{L_p(|x| \geq \sigma(4t))}. \end{aligned} \quad (1.12a)$$

and its averaged ‘cousin’

$$\begin{aligned} \bar{\omega}_{r,p}(f, W, t) : &= \left(\frac{1}{t} \int_0^t \|W \Delta_{h\Phi_t(x)}^r(f, x, \mathbb{R})\|_{L_p(|x| \leq \sigma(2t))}^p dh \right)^{\frac{1}{p}} \\ &+ \inf_{P \in \mathcal{P}_{r-1}} \|(f - P)W\|_{L_p(|x| \geq \sigma(4t))}. \end{aligned} \quad (1.12b)$$

If $p = \infty$, we set :

$$\bar{\omega}_{r,p}(f, W, ;) = \omega_{r,p}(f, W, ;).$$

Observe that

$$\bar{\omega}_{r,p}(f, W, t) \leq \omega_{r,p}(f, W, t)$$

for every fixed $t \in \mathbb{R}$.

The inf in the tail is at first disconcerting, but note that it is over polynomials of degree at most $r - 1$, not n . Its presence ensures that for $f \in \mathcal{P}_{r-1}$, $\omega_{r,p}(f, W, t) \equiv 0$. The modulus of continuity is rather difficult to assimilate (as is the case with all its cousins [6], [7] for weighted approximation on \mathbb{R}). A good way to view the function $\sigma(t)$, is that for purposes of approximation by polynomials of degree at most n , essentially $t = \frac{an}{n}$, the main part of the modulus is taken over the range $[-a\frac{n}{2}, a\frac{n}{2}]$, and the tail is taken over $\mathbb{R} \setminus [-a_{n/2}, a_{n/2}]$. Moreover, the function $\Phi_t(x)$ describes the improvement in the degree of approximation near $\pm a_{n/2}$, in much the same way that $\sqrt{1 - x^2}$ does for weights on $[-1, 1]$.

It is possible to replace $\sigma(2t)$ by the somewhat larger term $\sigma(t) - At$ and $\sigma(4t)$ by the somewhat smaller term $\sigma(t) - Bt$ for suitable A, B in our modulus, under additional conditions on Q . However, it hardly seems worth the effort, as the resulting modulus is almost certainly equivalent to the above one. As evidence of this, we note that in [3], the first author proves that the above modulus is equivalent to a natural K-functional/realization functional.

Following is our main Jackson theorem:

Theorem 1.2 *Let $W := e^{-Q} \in \mathcal{E}_1$. Let $r \geq 1$ and $0 < p \leq \infty$. Then for $f : \mathbb{R} \rightarrow \mathbb{R}$ for which $fW \in L_p(\mathbb{R})$, (and for $p = \infty$, we require f to be continuous, and fW to vanish at $\pm\infty$), we have for $n \geq C_3$,*

$$E_n[f]_{W,p} \leq C_1 \bar{\omega}_{r,p} \left(f, W, C_2 \frac{a_n}{n} \right) \leq C_1 \omega_{r,p} \left(f, W, C_2 \frac{a_n}{n} \right) \quad (1.13)$$

where C_j , $j = 1, 2, 3$, do not depend on f or n .

Remark.

We remark that it is possible using the methods of [6] and [3] to prove Theorem 1.2 for $n \geq r - 1$.

Unfortunately, the modulus $\omega_{r,p}(f, W, t)$ is not obviously monotone increasing in t . So we also present a result involving the increasing modulus

$$\begin{aligned} \omega_{r,p}^*(f, W, t) : &= \sup_{\substack{0 < h \leq t \\ 0 < \tau \leq L}} \|W \Delta_{\tau h \Phi_n(x)}^r(f, x, \mathbb{R})\|_{L_p(|x| \leq \sigma(2h))} \\ &+ \inf_{P \in \mathcal{P}_{r-1}} \|(f - P)W\|_{L_p(|x| \geq \sigma(4t))}. \end{aligned} \quad (1.14)$$

Here L is a fixed (large enough) number independent of f, t .

Theorem 1.3 *Under the hypotheses of Theorem 1.2,*

$$E_n[f]_{W,p} \leq C_3 \omega_{r,p}^* \left(f, W, C_4 \frac{a_n}{n} \right), \quad (1.15)$$

where C_j , $j = 3, 4$ do not depend on f or n .

It seems likely that one should only really need $\tau = L$ in the definition of $\omega_{r,p}^*$, but we have only been able to prove this under additional conditions, see Section 7.

The modulus of continuity is analyzed in [3], and in particular the relationship to K-functionals/realization functionals, is discussed. These have the consequence, that at least for $p \geq 1$, we can dispense with the constant C_2 inside the modulus $\omega_{r,p}$ in (1.13) or (1.15). For $p < 1$, this requires extra hypotheses on W .

The paper is organized as follows: In Section 2, we present some technical details related to Q , a_u , and so on. In Section 3, we present estimates involving $\sigma(t)$ and Φ_t . In Section 4, we obtain polynomial approximations to W^{-1} over suitable intervals, and then in Section 5, we present our crucial approximations to characteristic functions. We prove Theorem 1.2 in Section 6 and Theorem in Section 1.3 in Section 7. Moreover, we discuss simplification of the modulus $\omega_{r,p}^*$ in Section 7.

We close this section with a little more notation. Throughout, C, C_1, C_2, \dots denote positive constants independent of n, x and $P \in \mathcal{P}_n$. The same symbol does not necessarily denote the same constant in different occurrences. We write $C \neq C(L)$ to indicate that C is independent of L . The notation $c_n \sim d_n$ means that $C_1 \leq c_n/d_n \leq C_2$ for the relevant range of n . Similar notation is used for functions and sequences of functions. In the sequel, we assume that $W = e^{-Q} \in \mathcal{E}_1$.

2 Technical Lemmas

Lemma 2.1

- (a) For some C_j , $j = 1, 2, 3$, and $s \geq r \geq C_3$,

$$\left(\frac{s}{r}\right)^{C_2 T(r)} \leq \frac{Q(s)}{Q(r)} \leq \left(\frac{s}{r}\right)^{C_1 T(s)}. \quad (2.1)$$

Moreover,

$$\left(\frac{s}{r}\right)^{C_2 T(r)} \frac{T(s)}{T(r)} \leq \frac{sQ'(s)}{rQ'(r)} \leq \frac{T(s)}{T(r)} \left(\frac{s}{r}\right)^{C_1 T(s)}. \quad (2.2)$$

- (b) Given $\delta > 0$, there exists C such that

$$T(y) \sim T\left(y\left(1 - \frac{\delta}{T(y)}\right)\right), \quad y \geq C. \quad (2.3)$$

- (c) Given $A \geq 0$, the functions $Q'(u)u^{-A}$ and $Q(u)u^{-A}$ are quasi-increasing for large enough u .

Proof

- (a) Firstly, (2.1) follows from the identity

$$\log \frac{Q(s)}{Q(r)} = \int_r^s \frac{T(t)}{t} dt$$

and the fact that T is quasi-increasing. Then the definition (1.3) of T gives (2.2).

- (b) We can reformulate (1.5) as

$$\frac{T(y)}{T(x)} \leq C_1 \left(\frac{Q(y)}{Q(x)} \right)^{C_2-1}.$$

Hence for $x = y(1 - \frac{\delta}{T(y)})$, the quasi-increasing nature of T gives

$$\begin{aligned} C_4 \leq \frac{T(y)}{T(x)} &\leq C_1 \exp \left((C_2 - 1) \int_x^y \frac{T(t)}{t} dt \right) \\ &\leq C_1 \exp \left(C_5 T(y) \log \frac{y}{x} \right) \leq C_6. \end{aligned}$$

Recall here that $T(y)$ is large for large y .

- (c) From (2.2) if $s \geq r \geq C$,

$$\frac{Q'(s)s^{-A}}{Q'(r)r^{-A}} \geq \frac{T(s)}{T(r)} \left(\frac{s}{r} \right)^{C_2 T(r) - 1 - A} \geq C_7.$$

Here we have used the quasi-monotonicity of T , and also that if C is large enough, then $C_2 T(r) - 1 - A \geq 0$. Similarly for $Q(s)s^{-A}$. \square

Next, some properties of a_u :

Lemma 2.2

(a) a_u is uniquely defined and continuous for $u \in (0, \infty)$, and is a strictly increasing function of u .

(b) For $u \geq C$,

$$a_u Q'(a_u) \sim u T(a_u)^{1/2}; \quad (2.4)$$

$$Q(a_u) \sim u T(a_u)^{-1/2}. \quad (2.5)$$

(c) Given fixed $\beta > 0$, we have for large u ,

$$T(a_{\beta u}) \sim T(a_u). \quad (2.6)$$

(d) Given fixed $\alpha > 1$,

$$\frac{a_{\alpha u}}{a_u} - 1 \sim \frac{1}{T(a_u)}. \quad (2.7)$$

(e) If C_2 is as in (1.5),

$$T(a_u) \leq C_6 u^{2\left(\frac{C_2-1}{C_2+1}\right)} = C_6 u^{2(1-\delta)} \quad (2.8)$$

with $\delta > 0$.

(f) If $\alpha > 1$, then for large enough u ,

$$\frac{Q(a_{\alpha u})}{Q(a_u)} \geq C_7 > 1. \quad (2.9)$$

(g) For some $C_8, C_9, C_{10}, C_{11}, C_{12}$, $u \geq C_8$, and $L \geq 1$,

$$\exp\left(C_{11} \frac{\log(C_{12}L)}{T(a_u)}\right) \geq \frac{a_{Lu}}{a_u} \geq 1 + C_9 \frac{\log(C_{10}L)}{T(a_{Lu})}. \quad (2.10)$$

Proof

(a) The function $u \rightarrow a_u$ is the inverse of the strictly increasing continuous function

$$a \rightarrow \frac{2}{\pi} \int_0^1 at Q'(at) \frac{dt}{\sqrt{1-t^2}}, \quad a \in (0, \infty),$$

which has right limit 0 at 0 and limit ∞ at ∞ . (Note that this function is continuous even if Q' is not). So the assertion follows.

(b) For u so large that $T(a_u) > 2$, we have

$$\begin{aligned}
\frac{u}{a_u Q'(a_u)} &= \frac{2}{\pi} \left[\int_0^{1-1/T(a_u)} + \int_{1-1/T(a_u)}^1 \right] \frac{a_u t Q'(a_u t)}{a_u Q'(a_u)} \frac{dt}{\sqrt{1-t^2}} \\
&\leq \frac{2}{\pi} T(a_u)^{1/2} \int_0^{1-1/T(a_u)} \frac{a_u Q'(a_u t)}{a_u Q'(a_u)} dt + \frac{2}{\pi} \int_{1-1/T(a_u)}^1 \frac{dt}{\sqrt{1-t^2}} \\
&\leq \frac{2}{\pi} T(a_u)^{1/2} \frac{Q(a_u) - Q(0)}{a_u Q'(a_u)} + \frac{4}{\pi} T(a_u)^{-1/2} \\
&\leq \frac{4}{\pi} T(a_u)^{1/2} \frac{Q(a_u)}{a_u Q'(a_u)} + \frac{4}{\pi} T(a_u)^{-1/2} = \frac{8}{\pi} T(a_u)^{-1/2}.
\end{aligned}$$

Here we also need u so large that $Q(a_u) \geq |Q(0)|$. So we have

$$a_u Q'(a_u) \geq \frac{\pi}{8} u T(a_u)^{1/2}.$$

In the other direction, (2.2) gives for large u ,

$$\begin{aligned}
\frac{u}{a_u Q'(a_u)} &= \frac{2}{\pi} \int_0^1 \frac{a_u t Q'(a_u t)}{a_u Q'(a_u)} \frac{dt}{\sqrt{1-t^2}} \\
&\geq C_1 \int_{1/2}^1 \frac{T(a_u t)}{T(a_u)} t^{C_1 T(a_u)} \frac{dt}{\sqrt{1-t^2}} \\
&\geq C_2 \frac{T\left(a_u \left(1 - \frac{1}{T(a_u)}\right)\right)}{T(a_u)} \left(1 - \frac{1}{T(a_u)}\right)^{C_1 T(a_u)} \times \\
&\quad \times \int_{1-1/T(a_u)}^1 \frac{dt}{\sqrt{1-t^2}} \\
&\geq C_3 T(a_u)^{-1/2}.
\end{aligned}$$

Here we have used (2.3) and the quasi-monotonicity of T . So we have (2.4). Then (2.5) follows from the definition of T .

(c) We can assume $\beta > 1$. Then by (2.5), and quasi-monotonicity of T ,

$$C_1 \leq \frac{T(a_{\beta u})}{T(a_u)} \sim \left[\frac{\beta u}{Q(a_{\beta u})} \right]^2 / \left[\frac{u}{Q(a_u)} \right]^2 \leq \beta^2.$$

(d) Now

$$\begin{aligned} \alpha u &= \frac{2}{\pi} \int_0^1 a_{\alpha u} t Q'(a_{\alpha u} t) \frac{dt}{\sqrt{1-t^2}} \\ &\geq \frac{2}{\pi} \int_{a_u/a_{\alpha u}}^1 a_u Q'(a_u) \frac{dt}{\sqrt{1-t^2}} \\ &\geq C_2 u T(a_u)^{1/2} \left(1 - \frac{a_u}{a_{\alpha u}}\right)^{1/2} \end{aligned}$$

by (2.4). Hence

$$1 - \frac{a_u}{a_{\alpha u}} \leq C_3 / T(a_u).$$

In the other direction,

$$\begin{aligned} \alpha u &= \frac{2}{\pi} \left[\int_0^{a_u/a_{\alpha u}} + \int_{a_u/a_{\alpha u}}^1 \right] a_{\alpha u} t Q'(a_{\alpha u} t) \frac{dt}{\sqrt{1-t^2}} \\ &\leq \frac{2}{\pi} \int_0^{a_u/a_{\alpha u}} a_{\alpha u} t Q'(a_{\alpha u} t) \frac{dt}{\sqrt{1 - \left(\frac{a_{\alpha u} t}{a_u}\right)^2}} + \frac{2}{\pi} a_{\alpha u} Q'(a_{\alpha u}) \int_{a_u/a_{\alpha u}}^1 \frac{dt}{\sqrt{1-t^2}} \\ &\leq \frac{a_u}{a_{\alpha u}} \left[\frac{2}{\pi} \int_0^1 a_u s Q'(a_u s) \frac{ds}{\sqrt{1-s^2}} \right] + \frac{4}{\pi} a_{\alpha u} Q'(a_{\alpha u}) \left(1 - \frac{a_u}{a_{\alpha u}}\right)^{1/2} \\ &\leq u + C u T(a_u)^{1/2} \left(1 - \frac{a_u}{a_{\alpha u}}\right)^{1/2} \end{aligned}$$

by (2.4) and (2.6). Then

$$1 - \frac{a_u}{a_{\alpha u}} \geq \left(\frac{\alpha - 1}{C}\right)^2 \frac{1}{T(a_u)}.$$

(e) We apply (1.5) with $y = a_u$ and $x = C_3$, so that

$$\begin{aligned} a_u Q'(a_u) &\leq C_4 Q(a_u)^{C_2} \\ \Rightarrow u T(a_u)^{1/2} &\leq C_5 (u T(a_u)^{-1/2})^{C_2}. \end{aligned}$$

Rearranging this gives (2.8).

(f) For large enough u ,

$$\begin{aligned} \frac{Q(a_{\alpha u})}{Q(a_u)} &= \exp\left(\int_{a_u}^{a_{\alpha u}} \frac{T(t)}{t} dt\right) \\ &\geq \exp\left(C_6 T(a_u) \log\left(\frac{a_{\alpha u}}{a_u}\right)\right) \geq \exp(C_7) > 1, \end{aligned}$$

by (d) of this lemma.

(g) From (1.5) with $y = a_{Lu}$ and $x = a_u$,

$$\frac{T(a_{Lu})}{T(a_u)} \leq C \left(\frac{Q(a_{Lu})}{Q(a_u)}\right)^{C_2-1}.$$

This forces $C_2 > 1$, as the left-hand side $\rightarrow \infty$ as $L \rightarrow \infty$. Then with the constants in \sim independent of L , (2.5) gives

$$\begin{aligned} \frac{Q(a_{Lu})}{Q(a_u)} &\sim \frac{Lu T(a_{Lu})^{-1/2}}{u T(a_u)^{-1/2}} \\ &\geq CL \left(\frac{Q(a_{Lu})}{Q(a_u)}\right)^{-(C_2-1)/2} \\ \Rightarrow \frac{Q(a_{Lu})}{Q(a_u)} &\geq CL^{\frac{2}{1+C_2}}. \end{aligned}$$

Then using (2.1),

$$\left(\frac{a_{Lu}}{a_u}\right)^{C_1 T(a_{Lu})} \geq CL^{\frac{2}{1+C_2}}.$$

We deduce the right-hand inequality in (2.10) from this last inequality and the inequality $\log t \leq t - 1, t \geq 1$. In the other direction, (2.1) and

then (2.5) give

$$\begin{aligned} \frac{a_{Lu}}{a_u} &\leq \left(\frac{Q(a_{Lu})}{Q(a_u)} \right)^{1/(C_2 T(a_u))} \\ &\leq \left(C_1 \frac{LuT(a_{Lu})^{-1/2}}{uT(a_u)^{-1/2}} \right)^{1/(C_2 T(a_u))} \leq (C_3 L)^{1/(C_2 T(a_u))}. \end{aligned}$$

Here the constants are independent of L and u . Then the left inequality in (2.10) follows. \square

We finish this section with an infinite finite-range inequality: We provide a proof as those in the literature [18], [20], [13], ...don't quite match our needs/hypotheses:

Lemma 2.3 Let $0 < p \leq \infty$, $s > 1$. Then for some $L, C_1, C_2 > 0$, $n \geq 1$, and $P \in \mathcal{P}_n$,

$$\|PW\|_{L_p(\mathbb{R})} \leq C_1 \|PW\|_{L_p(-a_{sn}, a_{sn})}. \quad (2.11)$$

Moreover,

$$\|PW\|_{L_p(|x| \geq a_{sn})} \leq C_1 e^{-C_2 n T(a_n)^{-1/2}} \|PW\|_{L_p(-a_{sn}, a_{sn})}. \quad (2.12)$$

Remark Note that (2.8) of Lemma 2.2(e) shows that for some $C_3 > 0$, and large enough n ,

$$nT(a_n)^{-1/2} \geq n^{C_3}.$$

Proof We may change Q in a finite interval without affecting (2.11), (2.12) apart from increasing the constants. Note too that the affect on a_u is marginal, and is absorbed into the fact that $s > 1$. Thus we may assume that Q' is continuous in $[-1, 1]$. This and the strict monotonicity of $tQ'(t)$ in $(0, \infty)$, allow us to apply existing sup-norm inequalities to deduce that for $P \in \mathcal{P}_n$,

$$\|PW\|_{L_\infty(\mathbb{R})} \leq C \|PW\|_{L_\infty[-a_{sn}, a_{sn}]}$$

For a precise reference, see [25] and [9,Thm.4.5]. Moreover, the proof of Lemma 5.1 in [13,pp.231-232] gives without change for $p < \infty$

$$|PW|^p(a_n x) \leq \frac{1}{\pi} \frac{2x}{x-1} \int_{-1}^1 |PW|^p(a_n t) dt, \quad x > 1. \quad (2.13)$$

Let $\langle x \rangle$ denote the greatest integer $\leq x$. Let δ be small and positive, let $l := \langle \delta n \rangle$ and let $T_l(x)$ denote the Chebyshev polynomial of degree l . Using the identity

$$T_l(x) = \frac{1}{2} \left[(x + \sqrt{x^2 - 1})^l + (x - \sqrt{x^2 - 1})^l \right], \quad x > 1, \quad (2.14)$$

it is not difficult to see that

$$T_l(x) \geq \begin{cases} \frac{1}{2} \exp\left(\frac{l}{\sqrt{2}} \sqrt{x-1}\right), & x \in \left(1, \frac{9}{8}\right) \\ \frac{1}{2} x^l, & x \geq 1 \end{cases}. \quad (2.15)$$

We now let $m := n + l = n + \langle \delta n \rangle$, $m' := n + 2l = n + 2\langle \delta n \rangle$ and apply (2.13) to $P(x)T_l(x/a_m) \in \mathcal{P}_m$. We obtain for $x > 1$,

$$|PW|^p(a_m x) \leq T_l(x)^{-p} \frac{1}{\pi} \frac{2x}{x-1} \int_{-1}^1 |PW|^p(a_m t) dt.$$

Replacing $a_m x$ by y , and integrating from $a_{m'}$ gives

$$\int_{a_{m'}}^{\infty} |PW|^p(y) dy \leq \left(\int_{-a_m}^{a_m} |PW|^p(s) ds \right) \left(\frac{2}{\pi} \int_{a_{m'}}^{\infty} \frac{y}{y-a_m} T_l\left(\frac{y}{a_m}\right)^{-p} \frac{dy}{a_m} \right).$$

Here using (2.15),

$$\begin{aligned} \int_{a_{m'}}^{\infty} \frac{y}{y-a_m} T_l\left(\frac{y}{a_m}\right)^{-p} \frac{dy}{a_m} &= \int_{a_{m'}/a_m}^{\infty} \frac{x}{x-1} T_l(x)^{-p} dx \\ &\leq C \left(\int_{a_{m'}/a_m}^{9/8} \frac{1}{x-1} \exp\left(-\frac{lp}{\sqrt{2}} \sqrt{x-1}\right) dx + \int_{9/8}^{\infty} x^{-lp} dx \right) \\ &\leq C_1 \left(\log\left(\frac{8}{\frac{a_{m'}}{a_m} - 1}\right) \exp\left(-C_2 l \left(\frac{a_{m'}}{a_m} - 1\right)^{1/2}\right) + \left(\frac{9}{8}\right)^{-lp} \right) \\ &\leq C_3 \exp\left(-C_4 n T(a_n)^{-1/2}\right). \end{aligned}$$

Here we have used (2.7) and our choice of l . Now if δ is small enough, $m' \leq sn$. Then (2.12) follows easily, and in turn yields (2.11). The proof for $p = \infty$ is similar but easier. \square

3 Technical lemmas on Φ_t

In this section, we present various estimates involving the functions σ and Φ_t . Throughout, we assume that $W = e^{-Q} \in \mathcal{E}_1$. Recall that

$$\sigma(t) := \inf \left\{ a_u : \frac{a_u}{u} \leq t \right\}, \quad t > 0;$$

$$\Phi_t(x) = \sqrt{\left| 1 - \frac{|x|}{\sigma(t)} \right|} + T(\sigma(t))^{-1/2}, \quad x > 0.$$

Lemma 3.1

- (a) There exists s_0, v_0 such that for $s \in (0, s_0)$, we can write $s = \frac{a_v}{v}$, where $v \geq v_0$. Moreover, we can write

$$\sigma(s) = \sigma\left(\frac{a_v}{v}\right) = a_{\beta(v)} \quad (3.1)$$

where

$$1 \geq \sigma\left(\frac{a_v}{v}\right) / a_v = a_{\beta(v)} / a_v \geq 1 - C/T(a_v). \quad (3.2)$$

In particular,

$$\lim_{v \rightarrow \infty} \frac{\beta(v)}{v} = 1. \quad (3.3)$$

- (b) There exist $C_1, C_2 > 0$ such that for $\frac{s}{2} \leq t \leq s$, and $s \leq C_1$,

$$1 \leq \frac{\sigma(t)}{\sigma(s)} \leq 1 + \frac{C_2}{T(\sigma(t))}. \quad (3.4)$$

- (c) There exist C_1, C_2 independent of s, t, x , such that for $0 < t < s \leq C_1$,

$$\Phi_s(x) \leq C_2 \Phi_t(x), \quad |x| \leq \sigma(s). \quad (3.5)$$

(d) There exists C_1 , such that for $0 < s \leq C_1$, and $\frac{s}{2} \leq t \leq s$,

$$\Phi_s(x) \sim \Phi_t(x), \quad x \in \mathbb{R}. \quad (3.6)$$

(e) Uniformly for $x \in \mathbb{R}$, and $n \geq 1$,

$$\Phi_{\frac{a_n}{n}}(x) \sim \sqrt{\left|1 - \frac{|x|}{a_n}\right|} + T(a_n)^{-1/2}. \quad (3.7)$$

Proof

(a) The existence of v for the given s , follows from the fact that $u \rightarrow a_u$ is continuous and

$$\frac{a_u}{u} \rightarrow 0, \quad u \rightarrow \infty.$$

The latter in turn follows from the faster than polynomial growth of Q and (2.5), which implies $Q(a_u) = o(u)$. The continuity of a_u allows us to write $\sigma(s) = a_{\beta(v)}$, some $\beta(v)$. Since

$$\sigma(s) = \sigma\left(\frac{a_v}{v}\right) \leq a_v$$

the left inequality in (3.2) follows. For the other direction, we note that by definition of $\sigma\left(\frac{a_v}{v}\right)$ and $\beta(v)$, we have $\beta(v) \leq v$ and

$$\frac{a_{\beta(v)}}{\beta(v)} \leq \frac{a_v}{v}$$

so

$$1 \leq \frac{v}{\beta(v)} \leq \frac{a_v}{a_{\beta(v)}} \leq \left(\frac{Q(a_v)}{Q(a_{\beta(v)})}\right)^{1/2}$$

for large enough v , by (2.1). Using (2.5), we obtain

$$1 \leq \frac{v}{\beta(v)} \leq C \left(\frac{vT(a_v)^{-1/2}}{\beta(v)T(a_{\beta(v)})^{-1/2}}\right)^{1/2} \leq C_1 \left(\frac{v}{\beta(v)}\right)^{1/2}.$$

It follows that $v \leq C_2\beta(v)$ and so $v \sim \beta(v)$. Then

$$1 \leq \frac{v}{\beta(v)} \leq \frac{a_v}{a_{\beta(v)}} \rightarrow 1, \quad v \rightarrow \infty,$$

by (2.7), so we have (3.3). Then (2.7) also gives the right inequality in (3.2).

(b) Write $s = \frac{a_u}{u}$ and $t = \frac{a_v}{v}$. Then as σ is decreasing,

$$1 \geq \frac{\sigma(s)}{\sigma(t)} = \frac{a_{\beta(u)}}{a_{\beta(v)}}$$

If we can show that

$$u \sim v \tag{3.8}$$

which in turn implies that

$$\beta(u) \sim \beta(v),$$

then (2.7) gives

$$1 \geq \frac{\sigma(s)}{\sigma(t)} \geq 1 - \frac{C}{T(a_v)}$$

which together with (2.6) gives the result. We proceed to establish (3.8). Suppose that it is not true, say, for example, we can have

$$\frac{u}{v} \rightarrow \infty.$$

For the corresponding s, t , our hypothesis is

$$\frac{1}{2} \leq \frac{t}{s} = \frac{a_v}{a_u} \frac{u}{v} \leq 1.$$

Then

$$\frac{a_v}{a_u} \rightarrow 0 \tag{3.9}$$

and (2.1) gives

$$\frac{Q(a_u)}{Q(a_v)} \geq \left(\frac{a_u}{a_v}\right)^{C_2 T(a_v)} \geq \left(\frac{a_u}{a_v}\right)^2,$$

for large u, v . But from (2.5),

$$\begin{aligned} \left(\frac{a_u}{a_v}\right)^2 &\leq \frac{Q(a_u)}{Q(a_v)} \sim \frac{uT(a_u)^{-1/2}}{vT(a_v)^{-1/2}} \\ &\leq C \frac{u}{v} \leq C \frac{a_u}{a_v}, \end{aligned}$$

again by our hypotheses on s, t . This contradicts (3.9). So we have (3.8) and the result.

(c) Let $\delta > 0$ be fixed. Firstly for $1 - |x|/\sigma(s) \geq \delta/T(\sigma(s))$,

$$\Phi_s(x) \sim \sqrt{1 - \frac{|x|}{\sigma(s)}} \leq \sqrt{1 - \frac{|x|}{\sigma(t)}} \leq \Phi_t(x).$$

Next, for $|1 - |x|/\sigma(s)| \leq \delta/T(\sigma(s))$,

$$\Phi_s(x) \sim T(\sigma(s))^{-1/2}.$$

This is bounded by $C\Phi_t(x)$ if $|1 - |x|/\sigma(t)| \geq \delta/T(\sigma(s))$, for a fixed $\delta > 0$. Otherwise, we have $|1 - |x|/\sigma(s)| \leq \delta/T(\sigma(s))$ and $|1 - |x|/\sigma(t)| \leq \delta/T(\sigma(s))$, so

$$\begin{aligned} \left|1 - \frac{\sigma(t)}{\sigma(s)}\right| &= \left|\left(1 - \frac{|x|}{\sigma(s)}\right) - \frac{|x|}{\sigma(s)}\left(\frac{\sigma(t)}{|x|} - 1\right)\right| \\ &\leq C_1\delta/T(\sigma(s)). \end{aligned}$$

If δ is small enough, we deduce from (2.7) and (2.6) that

$$T(\sigma(t)) \sim T(\sigma(s))$$

and again (3.5) follows.

(d) Write $s = \frac{au}{v}$ and $t = \frac{av}{u}$. Then we have (3.8), so

$$\begin{aligned} \left|1 - \frac{|x|}{\sigma(t)}\right| &= \left|1 - \frac{|x|}{\sigma(s)} + \left[\frac{|x|}{\sigma(s)} - 1 + 1\right] \left(1 - \frac{\sigma(s)}{\sigma(t)}\right)\right| \\ &\leq \left|1 - \frac{|x|}{\sigma(s)}\right| \left[1 + O\left(\frac{1}{T(\sigma(s))}\right)\right] + O\left(\frac{1}{T(\sigma(s))}\right). \end{aligned}$$

Then we obtain for $x \in \mathbb{R}$,

$$\left|1 - \frac{|x|}{\sigma(t)}\right|^{1/2} \leq C\Phi_s(x).$$

Also $T(\sigma(t)) \sim T(\sigma(s))$, so

$$\Phi_t(x) \leq C\Phi_s(x).$$

The converse inequality follows similarly.

(e) By (a), we can write

$$\sigma\left(\frac{a_n}{n}\right) = a_{\beta(n)} = a_n(1+o(1)).$$

Recall that

$$\Phi_{\frac{a_n}{n}}(x) = \sqrt{\left|1 - \frac{|x|}{\sigma\left(\frac{a_n}{n}\right)}\right|} + T\left(\sigma\left(\frac{a_n}{n}\right)\right)^{-1/2}.$$

Here by (2.6) and (a) of this lemma,

$$T\left(\sigma\left(\frac{a_n}{n}\right)\right) \sim T(a_n)$$

and much as in (d),

$$\left|1 - \frac{|x|}{\sigma\left(\frac{a_n}{n}\right)}\right| \sim \left|1 - \frac{|x|}{a_n}\right|$$

for large n and $|x| \leq a_{n/2}$ or $|x| \geq a_{2n}$. In the range $a_{n/2} \leq |x| \leq a_{2n}$, both the left and right-hand side of (3.7) are $\sim T(a_n)^{-1/2}$. \square

Lemma 3.2

(a) Let $L > 0$. Uniformly for $u \geq 1$, and $|x|, |y| \leq a_u$, such that

$$|x - y| \leq L \frac{a_u}{u} \sqrt{\left|1 - \frac{|y|}{a_u}\right|}, \quad (3.10)$$

we have

$$W(x) \sim W(y) \quad (3.11)$$

and

$$1 - \frac{|x|}{a_{2u}} \sim 1 - \frac{|y|}{a_{2u}}. \quad (3.12)$$

(b) Let $L, M > 0$. For $t \in (0, t_0)$, $|x|, |y| \leq \sigma(Mt)$ such that

$$|x - y| \leq Lt\Phi_t(x), \quad (3.13)$$

we have (3.11) and

$$\Phi_t(x) \sim \Phi_t(y). \quad (3.14)$$

Proof

- (a) It suffices to prove (3.11), (3.12) for large u . Moreover, (3.11) and (3.12) are immediate for $|x| \leq C$, and large u . Let us suppose that $C \leq x \leq y \leq x + L \frac{a_u}{u} \sqrt{\left|1 - \frac{|y|}{a_u}\right|}$. Then as $Q'(s)$ is quasi-increasing for large s ,

$$0 \leq Q(y) - Q(x) \leq C_1 Q'(y)(y - x).$$

We have then (3.11) for

$$y - x = O\left(\frac{1}{Q'(y)}\right). \quad (3.15)$$

We shall show that

$$a_u Q'(y) \sqrt{\left|1 - \frac{y}{a_u}\right|} \leq C_2 u, \quad (3.16)$$

so that (3.10) implies (3.15) and hence (3.11). If firstly, $0 < y \leq \frac{a_u}{2}$, then

$$\begin{aligned} a_u Q'(y) \sqrt{\left|1 - \frac{y}{a_u}\right|} &\leq C_3 a_u Q'(y) \int_{1/2}^1 \frac{dt}{\sqrt{1-t^2}} \\ &\leq C_4 \int_{1/2}^1 a_u t Q'(a_u t) \frac{dt}{\sqrt{1-t^2}} \leq C_5 u. \end{aligned}$$

If on the other hand, $\frac{a_u}{2} \leq y \leq a_u$,

$$a_u Q'(y) \sqrt{\left|1 - \frac{y}{a_u}\right|} \leq C_6 \int_{y/a_u}^1 a_u t Q'(a_u t) \frac{dt}{\sqrt{1-t^2}} \leq C_7 u.$$

So we have (3.16) in all cases. Next from (3.10) and as $y \leq a_u$,

$$\begin{aligned} 1 &\leq \frac{1 - \frac{x}{a_{2u}}}{1 - \frac{y}{a_{2u}}} = 1 + \frac{y - x}{a_{2u} \left(1 - \frac{y}{a_{2u}}\right)} = 1 + O\left(\frac{1}{u \sqrt{1 - \frac{y}{a_{2u}}}}\right) \\ &= 1 + O\left(\frac{1}{u \sqrt{1 - \frac{a_u}{a_{2u}}}}\right) = 1 + O\left(\frac{T(a_u)^{1/2}}{u}\right) = 1 + o(1), \end{aligned}$$

by (2.7) and (2.8).

(b) Write $Mt = \frac{a_u}{u}$, so that $|x|, |y| \leq \sigma(Mt) \leq a_u$, and we can recast (3.13) as

$$|x - y| \leq C_1 \frac{a_u}{u} \left[\sqrt{1 - \frac{|x|}{a_u}} + T(a_u)^{-1/2} \right] \leq C_2 \frac{a_{2u}}{2u} \sqrt{1 - \frac{|x|}{a_{2u}}}$$

by (2.7), (3.6) and (3.7). Then (a) gives (3.11), and (3.14) follows easily from (3.12). \square

4 Polynomial Approximation of W^{-1}

The result of this section is:

Theorem 4.1 *For $n \geq 1$, there exist polynomials G_n of degree at most Cn , such that*

$$0 \leq G_n(x) \leq W^{-1}(x), \quad x \in \mathbb{R}; \quad (4.1)$$

and

$$G_n(x) \sim W^{-1}(x), \quad |x| \leq a_n. \quad (4.2)$$

We remark that this does not follow from existing results in the literature on approximation by weighted polynomials of the form $P_n(x)W(a_n x)$ [14], [26] as our weights do not satisfy their hypotheses. The methods of Totik [26] can be applied to give sharper results but we base our proof on:

Lemma 4.2 *There exists an even entire function*

$$G(x) = \sum_{j=0}^{\infty} g_j x^{2j}, \quad g_j \geq 0 \quad \forall j, \quad (4.3)$$

such that

$$G(x) \sim W^{-1}(x), \quad x \in \mathbb{R}. \quad (4.4)$$

Proof Set

$$Q_1(r) := Q(\sqrt{r});$$

and

$$\psi(r) := rQ'_1(r) = \frac{1}{2} \sqrt{r} Q'(\sqrt{r}).$$

Then ψ is increasing in $(0, \infty)$, and if $\lambda > 1$, $r \geq r_0$, the quasi-increasing nature of Q' gives for some $C \neq C(\lambda)$,

$$\psi(\lambda r) - \psi(r) \geq \frac{1}{2} \sqrt{r} Q'(\sqrt{r}) (\sqrt{\lambda} C - 1) \geq 1$$

if λ is large enough. Moreover, $\phi(r) := e^{Q_1(r)}$ admits the representation

$$\phi(r) = \phi(1) \exp\left(\int_1^r \frac{\psi(s)}{s} ds\right), \quad r \geq 1.$$

By a theorem of Clunie and Kövari [2,Thm4,p.19], there exists entire

$$G_1(r) = \sum_{j=0}^{\infty} g_j r^j, \quad g_j \geq 0 \quad \forall j$$

such that

$$G_1(r) \sim \phi(r) := \exp(Q(\sqrt{r})), \quad r \geq r_0.$$

Then assuming $g_0 > 0$ as we can, we see that

$$G(r) := G_1(r^2)$$

satisfies (4.4). \square

In the analogous construction for Freud weights, the second author and Z. Ditzian used as the polynomials G_n the partial sums of G . However in the Erdős case, for partials sums of degree $O(n)$, we only have

$$G_n(x) \sim W^{-1}(x)$$

for $|x| \leq q_n$, where q_n is Freud's quantity, the root of the equation

$$n = q_n Q'(q_n).$$

Although $a_n/q_n \rightarrow 1$, $n \rightarrow \infty$ for Erdős weights, in effect, q_n is significantly smaller than a_n . (We cannot properly describe, using only q_n , the

improvement in the degree of approximation near $\pm a_n$). So we use a more sophisticated interpolant:

Proof of Theorem 4.1 Let J be a positive even integer (to be chosen large enough later) and let $T_n(x)$ denote the classical Chebyshev polynomial on $[-1, 1]$. Let G_n denote the Lagrange interpolant to G at the zeros of $T_n(x/a_n)^J$ so that G_n has degree at most $Jn - 1$, and admits the error representation

$$(G - G_n)(x) = \frac{1}{2\pi i} \int_{\Gamma} \frac{G(t)}{t - x} \left(\frac{T_n(x/a_n)}{T_n(t/a_n)} \right)^J dt$$

for x inside Γ . We shall choose Γ to be the ellipse with foci at $\pm a_n$, intersecting the real and imaginary axes at $\frac{a_n}{2}(\rho + \rho^{-1})$ and $\frac{a_n}{2}(\rho - \rho^{-1})$ respectively. Here we shall choose for some fixed small $\varepsilon > 0$,

$$\rho := 1 + \left(\frac{\varepsilon}{T(a_n)} \right)^{1/2}.$$

Since G has non-negative Maclaurin series coefficients, and satisfies (4.4), we deduce that

$$\delta_n := \|G_n/G - 1\|_{L^\infty[-a_n, a_n]} \leq C_1 \frac{W^{-1}\left(\frac{a_n}{2}(\rho + \rho^{-1})\right)}{(\rho - 1)^2} \frac{1}{\min_{t \in \Gamma} |T_n(t/a_n)|^J}.$$

Now for $t \in \Gamma$, we can write $t = \frac{a_n}{2}(z + z^{-1})$ where $|z| = \rho$, so that

$$\begin{aligned} |T_n(t/a_n)| &= \left| T_n\left(\frac{1}{2}(z + z^{-1})\right) \right| = \left| \frac{1}{2}(z^n + z^{-n}) \right| \\ &\geq \frac{1}{2}(\rho^n - \rho^{-n}) \geq \exp(C_2 n T(a_n)^{-1/2}). \end{aligned}$$

(Recall that $nT(a_n)^{-1/2} \rightarrow \infty$ as $n \rightarrow \infty$ and in fact grows faster than a power of n). It is important here that C_2 is independent of J . Next

$$\frac{a_n}{2}(\rho + \rho^{-1}) \leq a_n \left(1 + C_3 \frac{\varepsilon}{T(a_n)} \right) \leq a_{2n}$$

if ε is small enough, and n is large enough, by (2.7). Then

$$W^{-1} \left(\frac{a_n}{2} (\rho + \rho^{-1}) \right) \leq \exp(C_4 Q(a_{2n})) \leq \exp(C_5 n T(a_n)^{-1/2})$$

where again it is important that C_5 is independent of J . Since $(\rho - 1)^{-2} \sim T(a_n)$ grows no faster than a power of n , we see that choosing J large enough, gives

$$\delta_n \rightarrow 0, \quad n \rightarrow \infty.$$

Then (4.4) gives (4.2).

We now turn to proving (4.1). It suffices to prove

$$0 \leq G_n \leq CW^{-1}$$

for then (4.1) follows on multiplying G_n by a suitable constant. Firstly, we can assume n is even (for odd n , we can use G_{n+1}) so that $H_n(x) := G_n(\sqrt{x})$ is a polynomial of degree at most $\frac{Jn}{2} - 1$ (recall T_n and J are even) that interpolates to the entire function $H(x) := G(\sqrt{x})$ at the $\frac{Jn}{2}$ zeros of $T_n \left(\frac{\sqrt{x}}{a_n} \right)^J$ that lie in $(0, a_n^2)$. Thus $H_n(x)$ is determined entirely by interpolation conditions. Let γ_n denote the leading coefficient of $T_n \left(x/\sqrt{a_n} \right)$. Then the usual derivative-error formula for Hermite interpolation gives for $x \in (0, \infty)$ and some $\xi = \xi(x) \in (0, \infty)$,

$$(H - H_n)(x) = \gamma_n^{-J} T_n \left(\frac{\sqrt{x}}{a_n} \right)^J \frac{H^{(\frac{Jn}{2})}(\xi)}{\left(\frac{Jn}{2} \right)!} \geq 0.$$

(Recall that H is entire and has non-negative Maclaurin series coefficients). So in \mathbb{R}

$$G_n \leq G \leq CW^{-1}.$$

To show that $G_n \geq 0$ in \mathbb{R} , we note that it is true in $[-a_n, a_n]$ and we must establish it elsewhere. We use an idea employed in the Posse-Markov-Stieltjes inequalities [8,p.30, Lemma 5.3] (There the proof is for $(-\infty, \infty)$, but the proof goes through for $(0, \infty)$ with trivial changes). Now H is absolutely monotone in $(0, \infty)$ and $H - H_n$ has $\frac{Jn}{2}$ zeros in $(0, a_n^2]$. If m is the number of zeros of $H_n(x)$ in $[a_n^2, \infty)$, Lemma 5.3 in [8,p.30] gives

$$\frac{Jn}{2} + m \leq \deg(H_n) + 1 \leq \frac{Jn}{2}.$$

So $m = 0$, that is H_n has no zeros in (a_n^2, ∞) . Thus $H_n \geq 0$ there, so $G_n \geq 0$ in \mathbb{R} . \square

5 Polynomials approximating characteristic functions

Our Jackson theorem is based on polynomial approximations to the characteristic function $\chi_{[a,b]}$ of an interval $[a, b]$. We believe the following result is of independent interest:

Theorem 5.1 *Let l be a positive integer. There exist J, C_1, n_0 such that for $n \geq n_0$ and $\tau \in [-a_n, a_n]$, there exist polynomials $R_{n,\tau}$ of degree at most $2lJn$ such that for $x \in \mathbb{R}$,*

$$\left| \chi_{[\tau, a_n]} - R_{n,\tau} \right| (x) W(x) / W(\tau) \leq C_1 \left(1 + \frac{n|x - \tau|}{a_n \sqrt{1 - \frac{|\tau|}{a_{2n}}}} \right)^{-l}. \quad (5.1)$$

We emphasize that the constants J, C_1, n_0 are independent of n, τ, x .

Remark. The method of proof of Theorem 5.1 in the unweighted case goes back to an old paper of Brudnyi [1]. We also make heavy use of polynomials from [12] built on the Chebyshev polynomials.

Lemma 5.2 *There exist C_1, B, n_1 such that for $n \geq n_1$ and $|\zeta| \leq \cos \frac{\pi}{2n}$, there exists a polynomial $V_{n,\zeta}$ of degree at most $n - 1$ with*

$$\|V_{n,\zeta}\|_{L^\infty[-1,1]} = V_{n,\zeta}(\zeta) = 1; \quad (5.2)$$

$$|V_{n,\zeta}(t)| \leq \frac{B\sqrt{1 - |\zeta|}}{n|t - \zeta|}, \quad t \in (-1, 1) \setminus \{\zeta\}. \quad (5.3)$$

Moreover,

$$V_{n,\zeta}(t) \geq \frac{1}{2}, \quad |t - \zeta| \leq C_1 \frac{\sqrt{1 - |\zeta|}}{n}. \quad (5.4)$$

The constants are independent of n, ζ, t .

Proof The assertions (5.2), (5.3) are Proposition 13.1 in [12]. The estimate (5.4) follows from the classical Bernstein inequality. \square

The polynomials $R_{n,\tau}$ are determined as follows: Let us suppose that, say,

$$a_1 \leq \tau \leq a_n.$$

Later on, we shall suppose that τ exceeds a fixed positive constant. We define

$$\zeta := \frac{\tau}{a_{2lJn}} \quad (5.5)$$

and if G_n are the polynomials of Theorem 4.1,

$$R_{n,\tau}(x) := \frac{\int_0^x G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds}{\int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds}. \quad (5.6)$$

The parameters $\tau^* > \tau$ and J are defined as follows: Let $A \in (0, 1]$ denote the constant in the quasi-monotonicity of Q' , so that

$$Q'(y) \geq A Q'(x), \quad y \geq x \geq 1. \quad (5.7)$$

Let M denote a positive constant such that for say, $u \geq u_0$,

$$Q'(x) \leq M Q'(a_u), \quad 1 \leq x \leq a_{2u}. \quad (5.8)$$

The existence of such an M follows from (2.4), (2.6). We set

$$H := H(n, \tau, l) := \frac{2l n}{A a_n Q'(\tau) \sqrt{1 - \zeta}} \quad (5.9)$$

and if $\tau = a_r$,

$$\tau^* := \tau^*(n, \tau) := \min \left\{ a_{2r}, a_n, \tau + 2 \frac{a_n}{n} \sqrt{1 - \zeta} H \log H \right\}. \quad (5.10)$$

The reason for this (complicated!) choice will become clearer later. We assume that $J \geq 4$ is so large that G_n has degree at most $Jn - 1$, and also

$$J \geq 16M/A \quad (5.11)$$

where A, M are as above. We also assume that J is a multiple of 4. Note that then $R_{n,\tau}$ has degree at most $Jn + lJn$. We first record some estimates of the terms in (5.6):

Lemma 5.3

(a) For $n \geq n_1$, and $C_1 \leq \tau \leq a_n$, we have

$$W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds \geq C_2 \frac{a_n}{n} \sqrt{1-\zeta}, \quad (5.12)$$

where $C_2 \neq C_2(n, \tau)$.

(b) For $x \in (\tau, a_{2lJn})$,

$$\int_x^{a_{2lJn}} V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{\frac{lJ}{2}} ds \leq C_1 \frac{a_n}{n} \sqrt{1-\zeta} \left(1 + \frac{n|x-\tau|}{a_n \sqrt{1-\zeta}} \right)^{-l} \quad (5.13)$$

and for $x \in (-a_{2lJn}, \tau)$,

$$\int_{-a_{2lJn}}^x V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{\frac{lJ}{2}} ds \leq C_1 \frac{a_n}{n} \sqrt{1-\zeta} \left(1 + \frac{n|x-\tau|}{a_n \sqrt{1-\zeta}} \right)^{-l}. \quad (5.14)$$

Here $C_1 \neq C_1(n, \tau)$.

Proof

(a) Let us denote the left-hand side of (5.12) by Γ . By (4.2) and (5.4),

$$\Gamma \geq C_2 W(\tau) \int_{\tau - C_3 \frac{a_n}{n} \sqrt{1-\zeta}}^{\tau} W^{-1}(s) ds \geq C_4 \frac{a_n}{n} \sqrt{1-\zeta},$$

where we have used (3.11) of Lemma 3.2(a).

(b) These follow in a straightforward fashion from the estimates (5.2), (5.3) and the fact that $J \geq 4$, so $lJ/2 > l + 1$. \square

Now we begin the proof of Theorem 5.1. We first show that it suffices to consider τ in the range $[S, a_n]$, for some fixed S .

Proof of Theorem 5.1 for $|\tau| \leq S$, where S is fixed

Note first that since for such τ ,

$$W(x)/W(\tau) \leq W(0)/W(S), \quad x \in \mathbb{R},$$

we must only prove there exists $R_{n,\tau}$ of degree at most n such that

$$\left| \chi_{[\tau, a_n]} - R_{n,\tau} \right| (x) \leq C_1 \left(1 + \frac{n|x - \tau|}{a_n \sqrt{1 - \frac{|\tau|}{a_{2n}}}} \right)^{-l},$$

for $|x| \leq a_{2n}$, and then our infinite-finite range inequality Lemma 2.3 gives the rest. Setting here $\xi := \tau/a_n$, and $s := x/a_n$, and $U_{n,\xi}(s) := R_{n,\tau}(x) = R_{n,\tau}(a_n s)$, we see that it suffices to show

$$\left| \chi_{[\xi, 1]}(s) - U_{n,\xi}(s) \right| \leq C_2 (1 + n|s - \xi|)^{-l}, \quad s \in [-2, 2].$$

We have used here that $|\xi| \leq \frac{1}{2}$, for large n . The existence of such polynomials is classical. See for example [4]. One could also base them on the $V_{n,\zeta}$ above. \square

It suffices to consider $\tau \in [S, a_n]$, where S is fixed

For, once this is done, we have the result for all $\tau \in [0, a_n]$. With the result for $\tau \geq 0$, we set

$$R_{n,-\tau}(x) := 1 - R_{n,\tau}(-x), \quad x \in \mathbb{R}.$$

It is not difficult to check the result for $-\tau$ from the corresponding result for τ , using the identity

$$\chi_{[-\tau, a_n]}(x) = 1 - \chi_{(\tau, a_n]}(-x), \quad x \in [a_{-n}, a_n]. \quad \square$$

In the sequel, we define $R_{n,\tau}$ by (5.5)-(5.10).

It suffices to prove (5.1) for $\tau \in [S, a_n]$ and $|x| \leq a_{2lJn}$

For then (5.1) for this restricted range implies

$$\left\| \left(1 + \left[\frac{n(x - \tau)}{a_n \sqrt{1 - \frac{\tau}{a_{2n}}}} \right]^2 \right)^l R_{n,\tau}(x) \frac{W(x)}{W(\tau)} \right\|_{L_\infty[-a_{2lJn}, a_{2lJn}]} \leq C_3 n^{C_4}$$

where $C_4 \neq C_4(n, \tau)$. Since the polynomial in the left-hand side has degree at most $2l + Jn + lJn \leq \eta 2lJn$, some fixed $\eta < 1$, if $l \geq 2$ and n is large

enough (as we can assume), then the infinite-finite range inequality Lemma 2.3 gives

$$\left\| \left(1 + \left[\frac{n(x-\tau)}{a_n \sqrt{1 - \frac{\tau}{a_{2n}}}} \right]^2 \right)^l R_{n,\tau}(x) \frac{W(x)}{W(\tau)} \right\|_{L_\infty(|x| \geq a_{2lJ_n})} \leq C_5 \exp(-n^{C_6}).$$

Then (5.1) follows for $|x| \geq a_{2lJ_n}$. \square

We can now begin the proof of (5.1) proper. We consider 5 different ranges of x : $[0, \tau)$, $[\tau, \tau^*]$, $(\tau^*, a_n]$, $(a_n, a_{2lJ_n}]$, $[-a_{2lJ_n}, 0)$. Moreover, we set

$$\Delta(x) := \left| \chi_{[\tau, a_n]} - R_{n,\tau} \right| (x) W(x) / W(\tau).$$

Proof of (5.1) for $x \in [0, \tau)$

Here using (4.1), and then (5.12),

$$\begin{aligned} \Delta(x) &= \frac{W(x) \int_0^x G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds}{W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds} \\ &\leq C \frac{W(x) \int_0^x W^{-1}(s) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds}{\frac{a_n}{n} \sqrt{1-\zeta}} \\ &\leq C \frac{\int_0^x V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds}{\frac{a_n}{n} \sqrt{1-\zeta}} \end{aligned}$$

by the monotonicity of W . Then (5.14) gives the result. \square

Proof of (5.1) for $x \in [\tau, \tau^*]$

Here

$$\begin{aligned} \Delta(x) &= \frac{W(x) \int_x^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds}{W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds} \\ &\leq C \frac{\int_x^{\tau^*} \exp(Q(s) - Q(x)) V_{n,\zeta} \left(\frac{s}{a_{2lJ_n}} \right)^{lJ} ds}{\frac{a_n}{n} \sqrt{1-\zeta}} \end{aligned}$$

by (4.1) and (5.12). Now for $s \in (x, \tau^*)$, the property (5.8) of Q' gives (recall $\tau^* \leq a_{2r}$)

$$Q(s) - Q(x) \leq \overline{MQ'}(a_r)(s - x) \leq MQ'(\tau)(s - \tau).$$

Then using our bounds on $V_{n,\zeta}$ in (5.2), (5.3), we have

$$\begin{aligned} \Delta(x) &\leq C_1 \frac{\int_x^{\tau^*} \exp(MQ'(\tau)(s - \tau)) \min\left\{1, \frac{Ba_{2lJn}\sqrt{1-\zeta}}{n(s-\tau)}\right\}^{lJ} ds}{\frac{a_{2lJn}}{n}\sqrt{1-\zeta}} \\ &= C_1 B \int_{\frac{n(x-\tau)}{Ba_{2lJn}\sqrt{1-\zeta}}}^{\frac{n(\tau^*-\tau)}{Ba_{2lJn}\sqrt{1-\zeta}}} \exp\left(\frac{a_{2lJn}}{a_n} \frac{2lMBu}{AH}\right) \min\left\{1, \frac{1}{u}\right\}^{lJ} du \\ &\leq C_2 \int_{\frac{n(x-\tau)}{Ba_{2lJn}\sqrt{1-\zeta}}}^{\frac{2}{B}H \log H} g(u) \min\left\{1, \frac{1}{u}\right\}^{lJ/2} du \end{aligned}$$

for say $n \geq n_1 = n_1(J, l)$ by (5.10), and where

$$g(u) := \exp\left(\frac{4lMBu}{AH}\right) \min\left\{1, \frac{1}{u}\right\}^{lJ/2}.$$

We claim that if J is large enough,

$$g(u) \leq C_3, u \in \left[0, \frac{2}{B}H \log H\right],$$

with C_3 independent of τ, n . Firstly we claim that if l is large enough,

$$H \geq e; H \geq e^{B/2} \tag{5.15}$$

uniformly for $\tau \in [S, a_n]$ and $n \geq n_0(J, l)$. First recall that B, J, A, M are independent of l (see (5.3), (5.7), (5.8), (5.11)). Then also from (3.16) for $\tau \in [S, a_n]$

$$a_n Q'(\tau) \sqrt{1 - \frac{\tau}{a_{2n}}} \leq Cn,$$

with $C \neq C(n, \tau, l)$. Then from (5.9),

$$H \geq \frac{2l}{AC} \left(\frac{1 - \frac{\tau}{a_{2n}}}{1 - \frac{\tau}{a_{2lJn}}} \right)^{1/2}.$$

Here for $n \geq n_0(J, l)$, using $1 - u \leq \log \frac{1}{u}$, $u \in (0, 1]$, we obtain

$$\begin{aligned} \frac{1 - \frac{\tau}{a_{2lJn}}}{1 - \frac{\tau}{a_{2n}}} &= 1 + \frac{\tau}{a_{2n}} \frac{1 - \frac{a_{2n}}{a_{2lJn}}}{1 - \frac{\tau}{a_{2n}}} \\ &\leq 1 + \frac{\log \frac{a_{2lJn}}{a_{2n}}}{1 - \frac{a_n}{a_{2n}}} \leq 1 + C_1 \log(lJ), \end{aligned}$$

by (2.7) and the left inequality in (2.10). Thus for $n \geq n_0(J, l)$, uniformly for $\tau \in [S, a_n]$,

$$H \geq \frac{C_2 l}{\sqrt{\log lJ}}.$$

So (5.15) follows if we choose l large enough. Then

$$g(u) \leq \exp\left(\frac{4lMB}{Ae}\right), \quad u \in (0, 1].$$

Next, by elementary calculus, g has at most one local extremum in $[1, \infty)$, and this is a minimum. Thus in any subinterval of $[1, \infty)$, g attains its maximum at the endpoints of that interval. In particular, we must only check that $g\left(\frac{2}{B}H \log H\right)$ is bounded. Note that by (5.15), $\frac{2}{B}H \log H \geq e > 1$. So

$$g\left(\frac{2}{B}H \log H\right) = \exp\left(l \log H \left\{ \frac{8M}{A} - \frac{J}{2} \right\} - \frac{Jl}{2} \log \left[\frac{2}{B} \log H \right]\right) \leq 1$$

as $J \geq 16M/A$ (see (5.11)) and $H \geq e^{B/2}$. So we have

$$\Delta(x) \leq C_4 \int_{\frac{n(x-\tau)}{Ba_{2lJn}\sqrt{1-\zeta}}}^{\infty} \min\left\{1, \frac{1}{u}\right\}^{lJ/2} du$$

and then (5.1) follows as $J \geq 4$. \square

Proof of (5.1) for $x \in (\tau^*, a_n]$

Here

$$\begin{aligned}
\Delta(x) &= \frac{W(x) \int_{\tau^*}^x G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds}{W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds} \\
&\leq C_1 \frac{\int_{\tau^*}^x \exp(Q(s) - Q(x)) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds}{\frac{a_n}{n} \sqrt{1-\zeta}} \\
&\leq C_2 \frac{n}{a_n \sqrt{1-\zeta}} \left(e^{Q\left(\frac{\tau+x}{2}\right) - Q(x)} \int_{\tau^*}^{\frac{\tau+x}{2}} V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds \right. \\
&\quad \left. + \int_{\frac{\tau+x}{2}}^x V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds \right) \\
&\leq C_3 \left\{ e^{Q\left(\frac{\tau+x}{2}\right) - Q(x)} \left[1 + \frac{n(\tau^* - \tau)}{a_n \sqrt{1-\zeta}} \right]^{-l} \right. \\
&\quad \left. + \left[1 + \frac{n(x - \tau)}{a_n \sqrt{1-\zeta}} \right]^{-l} \right\} \tag{5.16}
\end{aligned}$$

by (5.3) and (5.13). Here if $\tau^* > \frac{\tau+x}{2}$, the first term in the last two lines can be dropped and we already have the desired estimate. In the contrary case, we must estimate the first term. We note that we can assume that $\tau^* < a_n$, for otherwise the current range of x is empty. We consider two subcases (recall the definition (5.10) of τ^*):

(I) $\tau^* = \tau + 2\frac{a_n}{n} \sqrt{1-\zeta} H \log H$

We shall show that

$$\Gamma := \frac{Q(x) - Q\left(\frac{\tau+x}{2}\right)}{l \log \left(1 + \frac{n(x-\tau)}{a_n \sqrt{1-\zeta}} \right)} \geq 1. \tag{5.17}$$

Then the first part of the first term in the right-hand side of (5.16) already gives the desired estimate; the second part of that first term

can be bounded by 1. By quasi-monotonicity (5.7) of Q' ,

$$Q(x) - Q\left(\frac{\tau + x}{2}\right) \geq AQ'(\tau)\left(\frac{x - \tau}{2}\right).$$

Setting

$$u := \frac{n(x - \tau)}{a_n \sqrt{1 - \zeta}},$$

we have

$$\Gamma \geq \frac{AQ'(\tau) \frac{a_n}{n} \sqrt{1 - \zeta} u}{2l \log(1 + u)} = \frac{u}{H \log(1 + u)}.$$

(Recall that H was defined at (5.9)). But

$$u \geq \frac{n(\tau^* - \tau)}{a_n \sqrt{1 - \zeta}} = 2H \log H.$$

Recall from (5.15) that $H \geq e$. Then since the function $\frac{u}{\log(1+u)}$ is increasing for $u \geq 2H \log H \geq e$, we obtain

$$\Gamma \geq \frac{2H \log H}{H \log(1 + 2H \log H)}.$$

Using the inequality $1 + 2t \log t \leq t^2$, $t \geq 2$, we have

$$\Gamma \geq \frac{2 \log H}{\log(H^2)} = 1.$$

So we have (5.17) and the result.

(II) $\tau^* = a_{2r}$

In this case, from (2.7),

$$\tau^* - \tau = a_{2r} - a_r \sim \frac{a_r}{T(a_r)} = \frac{\tau}{T(\tau)}.$$

Now if $\tau^* \leq x \leq \tau(1 + \frac{1}{T(\tau)})$, then

$$x - \tau \sim \tau^* - \tau$$

and the second part of the first term in the right-hand side of (5.16) already gives the desired estimate (the first part of the first term can be estimated by 1). If $x > \tau \left(1 + \frac{1}{T(\tau)}\right)$, then

$$\frac{x}{\left(\frac{x+\tau}{2}\right)} \geq 1 + \frac{1}{2T(\tau) + 1} \geq 1 + \frac{1}{3T(\tau)}$$

for large τ , so from (2.1),

$$\frac{Q(x)}{Q\left(\frac{x+\tau}{2}\right)} \geq \left(1 + \frac{1}{3T(\tau)}\right)^{C_2 T\left(\frac{x+\tau}{2}\right)} \geq C_3 > 1.$$

(Recall that $\frac{x+\tau}{2} > \tau$). Then

$$e^{Q\left(\frac{\tau+x}{2}\right) - Q(x)} \left[1 + \frac{n(\tau^* - \tau)}{a_n \sqrt{1 - \zeta}}\right]^{-l} \leq e^{-C_4 Q(x)} \left[1 + \frac{C_5 n \tau}{a_n T(\tau) \sqrt{1 - \zeta}}\right]^{-l}.$$

This will admit the desired estimate, namely

$$C_6 \left[1 + \frac{n(x - \tau)}{a_n \sqrt{1 - \zeta}}\right]^{-l}$$

provided

$$e^{C_4 Q(x)/l} \frac{\tau}{T(\tau)} \geq C_7(x - \tau).$$

But

$$e^{C_4 Q(x)/l} \frac{\tau}{T(\tau)} \geq C_8 \frac{e^{C_4 Q(x)/l}}{T(x)} \geq C_9 Q(x) \geq C_{10} x > C_{10}(x - \tau)$$

by (2.5), (2.8) and the faster than polynomial growth of Q , so we have the desired estimate. \square

Proof of (5.1) for $x \in (a_n, a_{2lJ_n}]$

Here, much as in the previous range,

$$\begin{aligned}
\Delta(x) &= \frac{W(x) \int_0^x G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds}{W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds} \\
&\leq C_2 \frac{n}{a_n \sqrt{1-\zeta}} \left(e^{Q\left(\frac{\tau+x}{2}\right)-Q(x)} \int_0^{\frac{\tau+x}{2}} V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds \right. \\
&\quad \left. + \int_{\frac{\tau+x}{2}}^x V_{n,\zeta} \left(\frac{s}{a_{2lJn}} \right)^{lJ} ds \right) \\
&\leq C_3 \left\{ e^{Q\left(\frac{\tau+x}{2}\right)-Q(x)} + \left[1 + \frac{n(x-\tau)}{a_n \sqrt{1-\zeta}} \right]^{-l} \right\}.
\end{aligned}$$

We must show that the first term on the last right-hand side admits a bound that is a constant multiple of the second term on the last right-hand side. Let us write $x = a_v$ (so $v \geq n$) and $\frac{\tau+x}{2} = a_u$ (so that $u < v$). If firstly $u \geq \frac{n}{2}$, then

$$\begin{aligned}
Q(x) - Q\left(\frac{\tau+x}{2}\right) &\geq C_4 Q'(a_{n/2})(x-\tau) \\
&\geq C_5 \frac{n}{a_n} T(a_n)^{1/2} (x-\tau) \\
&\geq C_6 \frac{n(x-\tau)}{a_n \sqrt{1-\zeta}} \\
&\geq C_7 l \log \left(1 + \frac{n(x-\tau)}{a_n \sqrt{1-\zeta}} \right)
\end{aligned}$$

by (2.4),(2.7). (Recall that $\zeta = \tau/a_{2lJn}$.) In this case the result follows. If $u < \frac{n}{2}$,

$$\begin{aligned}
Q(x) - Q\left(\frac{\tau+x}{2}\right) &\geq Q(a_n) - Q(a_{n/2}) \\
&\geq C_8 Q(a_n) \geq C_9 n T(a_n)^{-1/2} \geq C_{10} n^{C_{11}}
\end{aligned}$$

by (2.5), (2.8). Since

$$\left[1 + \frac{n(x - \tau)}{a_n \sqrt{1 - \zeta}}\right]^{-l} \geq n^{-C_{11}}$$

the result again follows. \square

Proof of (5.1) for $x \in [-a_{2lJn}, 0)$

Here using the evenness of W and (4.1), (5.12) as before gives

$$\begin{aligned} \Delta(x) &= \frac{W(x) \int_x^0 G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}}\right)^{lJ} ds}{W(\tau) \int_0^{\tau^*} G_n(s) V_{n,\zeta} \left(\frac{s}{a_{2lJn}}\right)^{lJ} ds} \\ &\leq C_2 \frac{n}{a_n \sqrt{1 - \zeta}} \left(\int_x^{\frac{x}{2}} V_{n,\zeta} \left(\frac{s}{a_{2lJn}}\right)^{lJ} ds + e^{Q(\frac{x}{2}) - Q(x)} \int_{\frac{x}{2}}^0 V_{n,\zeta} \left(\frac{s}{a_{2lJn}}\right)^{lJ} ds \right) \\ &\leq C_3 \left\{ \left[1 + \frac{n \left|\frac{x}{2} - \tau\right|}{a_n \sqrt{1 - \zeta}}\right]^{-l} + e^{Q(\frac{x}{2}) - Q(x)} \left[1 + \frac{n\tau}{a_n \sqrt{1 - \zeta}}\right]^{-l} \right\}. \end{aligned}$$

Here $\left|\frac{x}{2} - \tau\right| = \frac{|x|}{2} + \tau \sim |x - \tau|$. Also, if $|x| \leq \tau$, then $\tau \sim \tau + |x| = |x - \tau|$. Otherwise (recall $\tau \geq S$), we have

$$e^{Q(\frac{x}{2}) - Q(x)} \leq e^{-C_4 Q(x)} \leq e^{-C_5 |x|} \leq (C_6 |x|)^{-l}.$$

Again as $|x|\tau \geq C_8 (\tau + |x|) = C_8 |x - \tau|$, the result follows. \square

6 The Proof of Theorem 1.2

In this section, we prove Theorem 1.2. Recall that our moduli of continuity are

$$\begin{aligned} \omega_{r,p}(f, W, t) : &= \sup_{0 < h \leq t} \left\| W \Delta_{h\Phi_t(x)}^r(f, x, \mathbb{R}) \right\|_{L_p(|x| \leq \sigma(2t))} \\ &\quad + \inf_{P \in \mathcal{P}_{r-1}} \left\| (f - P)W \right\|_{L(|x| \geq \sigma(4t))} \end{aligned}$$

and

$$\begin{aligned} \bar{\omega}_{r,p}(f, W, t) : &= \left(\frac{1}{t} \int_0^t \left\| W \Delta_{h\Phi_t(x)}^r(f, x, \mathbb{R}) \right\|_{L^p(|x| \leq \sigma(2t))}^p dh \right)^{\frac{1}{p}} \\ &+ \inf_{P \in \mathcal{P}_{r-1}} \left\| (f - P)W \right\|_{L(|x| \geq \sigma(4t))} \end{aligned}$$

where

$$\sigma(t) = \inf \left\{ a_u : \frac{a_u}{u} \leq t \right\}.$$

We need further moduli of continuity. If I is an interval, and $f : I \rightarrow \mathbb{R}$, we define for $t > 0$,

$$\Lambda_{r,p}(f, t, I) := \sup_{0 < h \leq t} \left(\int_I |\Delta_h^r(f, x, I)|^p dx \right)^{1/p} \quad (6.1)$$

and its averaged cousin

$$\Omega_{r,p}(f, t, I) := \left(\frac{1}{t} \int_0^t \int_I |\Delta_s^r(f, x, I)|^p dx ds \right)^{1/p}. \quad (6.2)$$

Note that for some C_1, C_2 depending only on r and p , (not on f, I, t)

$$C_1 \leq \Lambda_{r,p}(f, t, I) / \Omega_{r,p}(f, t, I) \leq C_2. \quad (6.3)$$

It seems that (6.3) first appeared in [23]. See also [4] and [24,p.191].

For large enough n , we choose a partition

$$-a_n = \tau_{0n} < \tau_{1n} < \dots < \tau_{nn} = a_n \quad (6.4)$$

such that if

$$I_{kn} := [\tau_{kn}, \tau_{k+1,n}], \quad 0 \leq k \leq n-1, \quad (6.5)$$

then uniformly in k and n ,

$$|I_{kn}| \sim \frac{a_n}{n} \sqrt{1 - \frac{|\tau_{kn}|}{a_{2n}}}. \quad (6.6)$$

($|I|$ denotes the length of the interval I). We also set $I_{nn} := \emptyset$. There are many ways to do this. For example, one can choose $\tau_{0,n} := -a_n$ and for $1 \leq k \leq n$, determine $\tau_{k,n}$ by

$$\frac{\int_{\tau_{k-1,n}}^{\tau_{k,n}} \frac{1}{\sqrt{1-\frac{|s|}{a_{2n}}}} ds}{\int_{-a_n}^{a_n} \frac{1}{\sqrt{1-\frac{|s|}{a_{2n}}}} ds} = \frac{1}{n}.$$

Let us set

$$I_n := [-a_n, a_n] = \bigcup_{k=0}^{n-1} I_{kn}, \quad (6.7)$$

$$\theta_{kn}(x) := \chi_{[\tau_{kn}, a_n]}(x) = \chi_{\bigcup_{i=k}^{n-1} I_{in}}(x) \quad (6.8)$$

and

$$I_{kn}^* := I_{kn} \cup I_{k+1,n}, \quad 0 \leq k \leq n-1. \quad (6.9)$$

By Whitney's theorem [24,p.195], we can find a polynomial p_k of degree at most r , such that

$$\|f - p_k\|_{L_p(I_{kn}^*)} \leq C_2 \Lambda_{r,p}(f, |I_{kn}^*|, I_{kn}^*) \quad (6.10)$$

with $C_2 \neq C_2(f, n, k, I_{kn}^*)$.

Now define an approximating piecewise polynomial/spline by

$$L_n[f](x) := p_0(x)\theta_{0n}(x) + \sum_{k=1}^{n-1} (p_k - p_{k-1})(x)\theta_{kn}(x). \quad (6.11)$$

We first show that $L_n[f]$ is a good approximation to f :

Lemma 6.1 Let $\Psi_n : [-a_n, a_n] \rightarrow \mathbb{R}$ be such that uniformly in n , and $x \in [-a_n, a_n]$,

$$\Psi_n(x) \sim \sqrt{1 - \frac{|x|}{a_{2n}}}. \quad (6.12)$$

Then for $0 < p < \infty$,

$$\|(f - L_n[f])W\|_{L_p(\mathbb{R})}^p$$

$$\begin{aligned}
&\leq C_1 \left\{ \frac{n}{a_n} \int_0^{C_2 \frac{a_n}{n}} \left\| W \Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R}) \right\|_{L_p[-a_n, a_n]}^p dh + \|fW\|_{L_p(|x| \geq a_n)}^p \right\} \\
&\leq C_3 \left(\sup_{0 < h \leq C_2 \frac{a_n}{n}} \left\| W \Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R}) \right\|_{L_p[-a_n, a_n]}^p \right. \\
&\quad \left. + \|fW\|_{L_p(|x| \geq a_n)}^p \right). \tag{6.13}
\end{aligned}$$

Here $C_j \neq C_j(f, n)$, $j = 1, 2, 3$. Moreover, the constants are independent of $\{\Psi_n\}$, depending only on the constants in \sim in (6.12). For $p = \infty$, (6.13) holds if we remove the exponents p .

Proof

We first deal with $p < \infty$. Now

$$\|(f - L_n[f])W\|_{L_p(\mathbb{R})}^p = \sum_{j=0}^{n-1} \Delta_{jn} + \|fW\|_{L_p(|x| \geq a_n)}^p, \tag{6.14}$$

where

$$\Delta_{jn} := \int_{I_{jn}} |f - L_n[f]|^p W^p. \tag{6.15}$$

Note that in $(\tau_{jn}, \tau_{j+1, n})$, $L_n[f] = p_j$, so that

$$\begin{aligned}
\Delta_{jn} &= \int_{I_{jn}} |f - p_j|^p W^p \\
&\leq \|W\|_{L_\infty(I_{jn})}^p C_2^p \Lambda_{r,p}^p(f, |I_{jn}^*|, I_{jn}^*) \quad (\text{by (6.10)}) \\
&\leq \|W\|_{L_\infty(I_{jn}^*)}^p \|W^{-1}\|_{L_\infty(I_{jn}^*)}^p \frac{C_3}{|I_{jn}^*|} \int_0^{|I_{jn}^*|} \int_{I_{jn}^*} |W \Delta_s^r(f, x, I_{jn}^*)|^p dx \, ds, \tag{6.16}
\end{aligned}$$

by (6.2), (6.3). Now from (3.11) of Lemma 3.2(a),

$$\|W\|_{L_\infty(I_{jn}^*)}^p \|W^{-1}\|_{L_\infty(I_{jn}^*)}^p \sim 1 \tag{6.17}$$

uniformly in j and n . Moreover, uniformly in j , n , and $x \in I_{jn}^*$,

$$|I_{jn}^*| \sim \frac{a_n}{n} \sqrt{1 - \frac{|x|}{a_{2n}}} \sim \frac{a_n}{n} \Psi_n(x).$$

Then we can continue (6.16) as

$$\begin{aligned}
\Delta_{jn} &\leq \frac{C_4}{|I_{jn}^*|} \int_{I_{jn}^*} \int_0^{|I_{jn}^*|} \left| W \Delta_s^r(f, x, I_{jn}^*) \right|^p ds dx \\
&= \frac{C_4}{|I_{jn}^*|} \int_{I_{jn}^*} \Psi_n(x) \int_0^{|I_{jn}^*|/\Psi_n(x)} \left| W \Delta_{t\Psi_n(x)}^r(f, x, I_{jn}^*) \right|^p dt dx \\
&\leq C_5 \frac{n}{a_n} \int_0^{C_6 \frac{an}{n}} \int_{I_{jn}^*} \left| W \Delta_{t\Psi_n(x)}^r(f, x, I_{jn}^*) \right|^p dx dt. \tag{6.18}
\end{aligned}$$

Adding over j gives

$$\sum_{j=0}^{n-1} \Delta_{jn} \leq C_5 \frac{n}{a_n} \int_0^{C_6 \frac{an}{n}} \int_{I_n} \left| W \Delta_{t\Psi_n(x)}^r(f, x, \mathbb{R}) \right|^p dx dt.$$

This and (6.14) give the result. Note that we have also effectively shown that

$$\sum_{j=0}^{n-1} \Omega_{r,p}^p(f, |I_{jn}^*|, I_{jn}^*) W^p(\tau_{jn}) \leq C_5 \frac{n}{a_n} \int_0^{C_6 \frac{an}{n}} \int_{I_n} \left| W \Delta_{t\Psi_n(x)}^r(f, x, \mathbb{R}) \right|^p dx dt. \tag{6.19}$$

For $p = \infty$, the proof is similar, but easier: We see that

$$\|(f - L_n[f])W\|_{L^\infty(\mathbb{R})} \leq \max \left\{ \max_{0 \leq j \leq n-1} \|(f - p_j)W\|_{L^\infty(I_{jn})}, \|fW\|_{L^\infty(|x| \geq a_n)} \right\}.$$

The rest of the proof is as before. \square

Now we can define our polynomial approximation to f :

$$P_n[f] := p_0(x)R_{n,\tau_{0n}}(x) + \sum_{k=1}^{n-1} (p_k - p_{k-1})(x)R_{n,\tau_{kn}}(x). \tag{6.20}$$

Note that this has been formed from $L_n[f]$ of (6.11) by replacing the characteristic function $\theta_{kn}(x) = \chi_{[\tau_{kn}, a_n]}(x)$ by its polynomial approximation $R_{n,\tau_{kn}}(x)$ formed in the previous section.

Lemma 6.2 Let $\{\Psi_n\}_n$ be as in the previous lemma. Then for $0 < p < \infty$,

$$\begin{aligned} & \| (L_n[f] - P_n[f])W \|_{L_p(\mathbb{R})} \\ & \leq C \left\{ \left(\frac{n}{a_n} \int_0^{\frac{c_1 a_n}{n}} \| W \Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R}) \|_{L_p[-a_n, a_n]}^p dh \right)^{\frac{1}{p}} \right. \\ & \quad \left. + \| fW \|_{L_p(I_{0n}^*)} \right\}. \end{aligned} \quad (6.21)$$

For $p = \infty$, this remains valid if we replace the p th powers by appropriate sup norms.

Proof

We see that if we define $p_{-1}(x) \equiv 0$,

$$(L_n[f] - P_n[f])(x) = \sum_{k=0}^{n-1} (p_k - p_{k-1})(x) (\theta_{kn}(x) - R_{n, \tau_{kn}}(x)). \quad (6.22)$$

We shall make substantial use of the following inequality: Let S be a polynomial of degree at most r , and $[a, b]$ be a real interval. Then for all $x \in \mathbb{R}$,

$$|S(x)| \leq C(b-a)^{-1/p} \left(1 + \frac{\min\{|x-a|, |x-b|\}}{b-a} \right)^r \|S\|_{L_p[a,b]}. \quad (6.23)$$

Here $C \neq C(a, b, x, S)$ but $C = C(p, r)$. This follows from standard Nikolskii inequalities and the Bernstein-Walsh inequality. See for example [24,p.193]. Hence for $x \in \mathbb{R}$, and $1 \leq k \leq n-1$,

$$|p_k - p_{k-1}|(x) \leq C |I_{kn}|^{-1/p} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^r \|p_k - p_{k-1}\|_{L_p(I_{kn})}.$$

This is still true for $k = 0$ if we recall that $p_{-1} \equiv 0$. Now for $1 \leq k \leq n-1$, (6.10) gives

$$\|p_k - p_{k-1}\|_{L_p(I_{kn})} \leq C_1 \sum_{i=k-1}^k \Lambda_{r,p}(f, |I_{in}^*|, I_{in}^*)$$

where $C_1 \neq C_1(f, k, n)$. This remains true for $k = 0$ if we set

$$|I_{-1,n}| := |I_{0n}|; |I_{-1,n}^*| := |I_{0n}^*|; \tau_{-1,n} := \tau_{0n};$$

and

$$\Lambda_{r,p}(f, |I_{-1,n}^*|, I_{-1,n}^*) := \|f\|_{L_p(I_{0n}^*)} =: \Omega_{r,p}(f, |I_{-1,n}^*|, I_{-1,n}^*)$$

Since (see (3.6), (3.7), (6.6)) uniformly in k, n , and $x \in \mathbb{R}$,

$$1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \sim 1 + \frac{|x - \tau_{k-1,n}|}{|I_{k-1,n}|}$$

we obtain from (6.23) and Theorem 5.1, uniformly for $0 \leq k \leq n-1$ and $x \in \mathbb{R}$,

$$\begin{aligned} & \left| (p_k - p_{k-1})(x)(\theta_{kn}(x) - R_{n,\tau_{kn}}(x)) \right| \frac{W(x)}{W(\tau_{kn})} \\ & \leq C_2 \sum_{i=k-1}^k |I_{in}|^{-1/p} \left(1 + \frac{|x - \tau_{in}|}{|I_{in}|} \right)^{r-l} \Omega_{r,p}(f, |I_{in}^*|, I_{in}^*). \end{aligned} \quad (6.24)$$

We consider three different ranges of p :

(I) $0 < p < 1$

Here from (6.22) and then (6.24),

$$\begin{aligned} \int_{\mathbb{R}} (|L_n[f] - P_n[f]|W)^p & \leq \sum_{k=0}^{n-1} \int_{\mathbb{R}} (|p_k - p_{k-1}| |\theta_{kn} - R_{n,\tau_{kn}}| W)^p \\ & \leq \sum_{k=-1}^{n-1} |I_{kn}|^{-1} \Omega_{r,p}^p(f, |I_{kn}^*|, I_{kn}^*) W^p(\tau_{kn}) \times \\ & \quad \times \int_{\mathbb{R}} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{(r-l)p} dx. \end{aligned} \quad (6.25)$$

Here if $(r-l)p < -1$,

$$|I_{kn}|^{-1} \int_{\mathbb{R}} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{(r-l)p} dx = \int_{\mathbb{R}} (1 + |u|)^{(r-l)p} du =: C_3 < \infty.$$

So

$$\int_{\mathbb{R}} \left(|L_n[f] - P_n[f] = \text{big}W \right)^p \leq C_4 \sum_{k=-1}^{n-1} \Omega_{r,p}^p(f, |I_{kn}^*|, I_{kn}^*) W^p(\tau_{kn}).$$

This is the same as our sum in (6.19) except for the term for $k = -1$. So the estimate (6.19) gives the estimate (6.21), keeping in mind our choice of $\Omega_{r,p}(f, |I_{-1,n}^*|, I_{-1,n}^*)$.

(II) $1 \leq p < \infty$

From (6.22) and (6.24) and then Hölder's inequality,

$$\begin{aligned} & \left\{ |L_n[f] - P_n[f]|(x)W(x) \right\}^p \\ & \leq C \left\{ \sum_{k=-1}^{n-1} |I_{kn}|^{-1/p} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{r-l} \Omega_{r,p}(f, |I_{kn}^*|, I_{kn}^*) W(\tau_{kn}) \right\}^p \\ & \leq C \sum_{k=-1}^{n-1} |I_{kn}|^{-1} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{(r-l)p/2} \times \\ & \quad \times \Omega_{r,p}^p(f, |I_{kn}^*|, I_{kn}^*) W^p(\tau_{kn}) \cdot S_n(x)^{p/q} \end{aligned} \quad (6.26)$$

where $q := p/(p-1)$ and

$$S_n(x) := \sum_{k=0}^{n-1} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{(r-l)q/2}.$$

We shall show that if $(r-l)q/2 < -1$, then

$$\sup_{n \geq 1} \sup_{x \in \mathbb{R}} S_n(x) \leq C_1 < \infty. \quad (6.27)$$

Note that $S_n(x)$ is a decreasing function of x for $x \geq a_n = \tau_{nn}$, so it suffices to consider $x \in [0, a_n]$. Recall that

$$|I_{kn}| \sim |I_{k+1,n}| \sim \frac{a_n}{n} \sqrt{1 - \frac{|\tau_{kn}|}{a_{2n}}}.$$

It is then not difficult to see that

$$\begin{aligned} S_n(x) & \leq C_2 \frac{n}{a_n} \int_{-a_n}^{a_n} \left(1 + \frac{n}{a_n} \frac{|x-u|}{\sqrt{1 - \frac{|u|}{a_{2n}}}} \right)^{(r-l)q/2} \frac{du}{\sqrt{1 - \frac{|u|}{a_{2n}}}} \\ & \leq C_3 n \int_{-1}^1 \left(1 + n \frac{|\bar{x}-s|}{\sqrt{1-s}} \right)^{(r-l)q/2} \frac{ds}{\sqrt{1-s}} \end{aligned}$$

where $\bar{x} := x/a_{2n}$, so that

$$1 - \bar{x} \geq 1 - a_n/a_{2n} \geq C_4 T(a_n)^{-1} \geq C_5 n^{-2}.$$

We make the substitution $(1 - s) = (1 - \bar{x})w$ to obtain

$$\begin{aligned} S_n(x) &\leq C_3 n \sqrt{1 - \bar{x}} \int_0^{\frac{2}{1 - \bar{x}}} \left(1 + n \sqrt{1 - \bar{x}} \frac{|w - 1|}{\sqrt{w}} \right)^{(r-l)q/2} \frac{dw}{\sqrt{w}} \\ &\leq C_4 n \sqrt{1 - \bar{x}} \left\{ \int_0^{1/2} \left[1 + \frac{n \sqrt{1 - \bar{x}}}{\sqrt{w}} \right]^{(r-l)q/2} \frac{dw}{\sqrt{w}} \right. \\ &\quad + \int_{1/2}^{3/2} \left[1 + n \sqrt{1 - \bar{x}} |w - 1| \right]^{(r-l)q/2} dw \\ &\quad \left. + \int_{3/2}^{2/(1 - \bar{x})} \left[1 + n \sqrt{(1 - \bar{x})w} \right]^{(r-l)q/2} \frac{dw}{\sqrt{w}} \right\}. \end{aligned}$$

(We can omit the third integral if $2/(1 - \bar{x}) \leq 3/2$.) We now make the substitutions $w = n^2(1 - \bar{x})v$ in the first integral, $v = n\sqrt{1 - \bar{x}}(w - 1)$ in the second integral, and $v = n^2(1 - \bar{x})w$ in the third integral. It is then not difficult to see that the resulting terms are bounded independent of n and x if l is large enough. (The least obvious is the first integral: there we need to ensure that $(r - l)q/4 - 1/2 \geq 0$, so that the integrand is bounded after the substitution). So we have (6.27). Then integrating (6.26) and using (6.19) gives our result.

(III) $p = \infty$

Now

$$\begin{aligned} |L_n[f] - P_n[f]|(x) &\leq C \sum_{k=0}^{n-1} |p_k - p_{k-1}|(x) |\theta_{kn} - R_{n, \tau_{kn}}|(x) W(x) \\ &\leq C \max_{-1 \leq k \leq n-1} \Omega_{r,p}(f, |I_{kn}^*|, I_{kn}^*) W(\tau_{kn}) \cdot \sum_{k=0}^{n-1} \left(1 + \frac{|x - \tau_{kn}|}{|I_{kn}|} \right)^{(r-l)}. \end{aligned}$$

As before, the sum is bounded if l is large enough. Then we can continue

this as

$$\begin{aligned}
&\leq C_1 \left\{ \sup_{0 \leq k \leq n-1} \sup_{0 < h \leq |I_{kn}^*|} \|\Delta_h^r(f, x, I_{kn}^*)W\|_{L_\infty(I_{kn}^*)} + \|fW\|_{L_\infty(I_{0n}^*)} \right\} \\
&\leq C_2 \left\{ \sup_{0 \leq k \leq n-1} \sup_{0 < h \leq Ca_n/n} \|\Delta_{h\Psi_n(x)}^r(f, x, I_{kn}^*)W\|_{L_\infty(I_{kn}^*)} + \|fW\|_{L_\infty(I_{0n}^*)} \right\} \\
&\leq C_3 \left\{ \sup_{0 < h \leq Ca_n/n} \|\Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R})W\|_{L_\infty(-a_n, a_n)} + \|fW\|_{L_\infty(I_{0n}^*)} \right\}. \quad \square
\end{aligned}$$

We can now turn to the

Proof of Theorem 1.2

Now recall that $R_{n,\tau}$ has degree at most $2lJn$, where J is as in the proof of Theorem 5.1. So $P_n[f]$ has degree at most $2lJn + r$. So, if $M := 3lJ$, we have for large n ,

$$\begin{aligned}
E_{Mn}[f]_{W,p} &\leq \|(f - P_n[f])W\|_{L_p(\mathbb{R})} \\
&\leq C \left\{ \|(f - L_n[f])W\|_{L_p(\mathbb{R})} + \|(L_n[f] - P_n[f])W\|_{L_p(\mathbb{R})} \right\} \\
&\leq C_1 \left\{ \left(\frac{n}{a_n} \int_0^{\frac{C_2 a_n}{n}} \|\Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R})W\|_{L_p(-a_n, a_n)}^p dh \right)^{\frac{1}{p}} \right. \\
&\quad \left. + \|fW\|_{L_p(|x| \geq a_n(1 - C_2[nT(a_n)^{1/2}] - 1))} \right\}. \quad (6.28)
\end{aligned}$$

Here we have used Lemmas 6.1 and 6.2, and also (6.6), which implies that

$$|I_{0n}^*| \sim \frac{a_n}{n} \sqrt{1 - \frac{a_n}{a_{2n}}} \sim \frac{a_n}{n} T(a_n)^{-1/2}.$$

Furthermore, at this stage, the functions $\{\Psi_n\}$ are any functions satisfying (6.12): they will be explicitly chosen later. Next for

$$Mn \leq j \leq M(n+1) \quad (6.29)$$

we write

$$n = \kappa j,$$

where $\kappa = \kappa(j, n)$. Note that

$$\kappa = \frac{n}{j} \rightarrow \frac{1}{M}, \quad j \rightarrow \infty. \quad (6.30)$$

We set

$$t := t(j) := \frac{Ma_j}{3j}.$$

Note that then

$$\frac{t}{a_n/n} = \frac{1}{3} \frac{Mn}{j} \frac{a_j}{a_n} = \frac{1}{3}(1 + o(1)), \quad n \rightarrow \infty. \quad (6.31)$$

Let $\beta > 3$. We claim that for large enough n ,

$$a_n \left(1 - C_2[nT(a_n)^{1/2}]^{-1}\right) \geq \sigma(\beta t). \quad (6.32)$$

To see this, note from (2.8) that

$$[nT(a_n)^{1/2}]^{-1} = o(T(a_n)^{-1})$$

so that by (2.7), if $1 > \alpha > 3/\beta$,

$$\begin{aligned} a_n \left(1 - C_2[nT(a_n)^{1/2}]^{-1}\right) &\geq a_n \left(1 - o\left(\frac{1}{T(a_n)}\right)\right) \geq a_{\alpha n} \\ &\geq \sigma\left(\frac{a_{\alpha n}}{\alpha n}\right) = \sigma\left(\frac{3t}{\alpha}[1 + o(1)]\right) \geq \sigma(\beta t), \end{aligned}$$

for large enough j , by first (3.2) and then (6.31). Next, we claim that if $0 < \gamma < 3$, then for n large enough,

$$a_n \leq \sigma(\gamma t). \quad (6.33)$$

To see this, note that by (6.31) if $1 < \delta < 3/\gamma$

$$\sigma(\gamma t) = \sigma\left(\frac{\gamma a_n}{3n}[1 + o(1)]\right) \geq \sigma\left(\frac{a_{\delta n}}{\delta n}\right) = a_{\delta n(1+o(1))} \geq a_n.$$

Here we also used the fact that σ is decreasing, and also (3.2), (3.3) with n large enough. Since also $a_n/n \leq 4t$ for large enough n , we can recast (6.28)

as

$$\begin{aligned}
E_j[f]_{W,p} &\leq E_{Mn}[f]_{W,p} \\
&\leq C_1 \left\{ \left(\frac{1}{2t} \int_0^{4Ct} \left\| \Delta_{h\Psi_n(x)}^r(f, x, \mathbb{R})W \right\|_{L_p(-\sigma(2t), \sigma(2t))}^p dh \right)^{\frac{1}{p}} \right. \\
&\quad \left. + \|fW\|_{L_p(|x| \geq \sigma(4t))} \right\}. \tag{6.34}
\end{aligned}$$

We now turn to our choice of $\{\Psi_n\}$: we must ensure that (6.12) holds with constants independent of x , j and n , that is

$$\Psi_n(x) \sim \sqrt{1 - \frac{|x|}{a_{2n}}}, \quad |x| \leq a_n.$$

But for this range of x ,

$$\sqrt{1 - \frac{|x|}{a_{2n}}} \sim \sqrt{1 - \frac{|x|}{a_{2n}}} + T(a_{2n})^{-1/2} \sim \Phi_{\frac{a_{2n}}{2n}}(x) \sim \Phi_t(x)$$

by Lemma 3.1(d), (e). We choose $h_1 := h/(4C)$ and $\Psi_n := \Phi_t/(4C)$ so that $h\Psi_n = h_1\Phi_t$, a choice satisfying (6.12). Then we rewrite (6.34) as

$$\begin{aligned}
E_j[f]_{W,p} &\leq C_1 \left\{ \left(\frac{4C}{2t} \int_0^t \left\| \Delta_{h_1\Phi_t(x)}^r(f, x, \mathbb{R})W \right\|_{L_p(-\sigma(2t), \sigma(2t))}^p dh_1 \right)^{\frac{1}{p}} \right. \\
&\quad \left. + \|fW\|_{L_p(|x| \geq \sigma(4t))} \right\}.
\end{aligned}$$

Replacing f by $f - P_0$ with a suitable choice of $P_0 \in \mathcal{P}_{r-1}$, we have for large enough j ,

$$\begin{aligned}
E_j[f]_{W,p} &= E_j[f - P_0]_{W,p} \\
&\leq C_3 \left\{ \left(\frac{1}{t} \int_0^t \left\| \Delta_{h_1\Phi_t(x)}^r(f, x, \mathbb{R})W \right\|_{L_p(-\sigma(2t), \sigma(2t))}^p dh_1 \right)^{\frac{1}{p}} \right. \\
&\quad \left. + \|(f - P_0)W\|_{L_p(|x| \geq \sigma(4t))} \right\}
\end{aligned}$$

$$\begin{aligned}
&\leq 2C_3 \left\{ \left(\frac{1}{t} \int_0^t \left\| \Delta_{h_1 \Phi_t(x)}^r(f, x, \mathbb{R}) W \right\|_{L_p(-\sigma(2t), \sigma(2t))}^p dh_1 \right)^{\frac{1}{p}} \right. \\
&\quad \left. + \inf_{P \in \mathcal{P}_{r-1}} \left\| (f - P) W \right\|_{L_p(|x| \geq \sigma(4t))} \right\} \\
&= C_3 \bar{\omega}_{r,p}(f, W, t) = C_3 \bar{\omega}_{r,p} \left(f, W, \frac{Ma_j}{3j} \right). \quad \square
\end{aligned}$$

For use in [3], we record the following form of Theorem 1.2:

Theorem 6.3 For $n \geq 1$, let $\lambda(n) \in [\frac{4}{5}, 1]$. Then

$$E_n[f]_{W,p} \leq C_1 \bar{\omega}_{r,p} \left(f, W, C_2 \lambda(n) \frac{a_n}{n} \right) \quad (6.35)$$

where C_1, C_2 do not depend on n or f or $\{\lambda(n)\}$. Moreover,

$$E_n[f]_{W,p} \leq C_1 \inf_{\rho \in [\frac{4}{5}, 1]} \bar{\omega}_{r,p} \left(f, W, C_2 \rho \frac{a_n}{n} \right). \quad (6.36)$$

Proof

Obviously (6.36) implies (6.35). The only difference to the above proof is that for $\rho \in [\frac{4}{5}, 1]$, we choose

$$t_1 := \rho t := \rho \frac{Ma_j}{3j}$$

to replace t above. Then from (6.31),

$$\frac{t_1}{a_n/n} = \frac{\rho}{3}(1 + o(1))$$

and here $\frac{\rho}{3} \in [\frac{4}{15}, \frac{1}{3}]$. Then as $4\rho > 3$, (6.32) above shows that

$$a_n \left(1 - C_2 [nT(a_n)^{1/2}]^{-1} \right) \geq \sigma(4\rho t) = \sigma(4t_1)$$

and as $\rho \leq 1$, (6.33) above shows that

$$a_n \leq \sigma(2\rho t) = \sigma(2t_1).$$

Moreover, for large enough n , $a_n/n \leq 3t(1 + o(1)) \leq 4t_1$. Choosing $h_1 := h/(4C)$ and $\Psi_n(x) := \Phi_{t_1}(x)/(4C)$ we note that (6.12) holds uniformly in ρ . We proceed as before to obtain

$$E_j[f]_{W,p} \leq C_1 \bar{\omega}_{r,p} \left(f, W, C_2 \frac{\rho a_j}{j} \right)$$

with constants independent of ρ, f, j . \square

7 The Proof of Theorem 1.3

We begin with a technical lemma, which refines part of Lemma 3.1:

Lemma 7.1

(a) For $0 < s < t \leq C$,

$$T(\sigma(t)) \left(1 - \frac{\sigma(t)}{\sigma(s)} \right) \leq C_1 \log \left(2 + \frac{t}{s} \right). \quad (7.1)$$

(b) For $0 < s < t \leq C$,

$$\sup_{x \in \mathbb{R}} \frac{\Phi_s(x)}{\Phi_t(x)} \leq C_2 \sqrt{\log \left(2 + \frac{t}{s} \right)}. \quad (7.2)$$

Hence, given $\gamma > 0$,

$$\sup_{x \in \mathbb{R}} \left(\frac{s}{t} \right)^\gamma \frac{\Phi_s(x)}{\Phi_t(x)} \leq C_3. \quad (7.3)$$

Proof

(a) We write $s = \frac{a_u}{u}$ and $t = \frac{a_v}{v}$. Note (with the notation of Lemma 3.1) that

$$a_{\beta(u)} = \sigma(s) \geq \sigma(t) = a_{\beta(v)},$$

so $\beta(u) \geq \beta(v)$. Using the inequality

$$1 - u \leq \log \frac{1}{u}, \quad u \in (0, 1]$$

we obtain

$$\begin{aligned} 1 - \frac{\sigma(t)}{\sigma(s)} &\leq \log \frac{\sigma(s)}{\sigma(t)} = \log \frac{a_{\beta(u)}}{a_{\beta(v)}} \\ &\leq C_1 \frac{\log C \frac{\beta(u)}{\beta(v)}}{T(a_{\beta(v)})} = C_1 \frac{\log C \frac{\beta(u)}{\beta(v)}}{T(\sigma(t))} \end{aligned} \quad (7.4)$$

by (2.10). Next, $\beta(u) = u(1+o(1))$, and similarly for $\beta(v)$, so it suffices to show that

$$\log \frac{u}{v} \leq C_2 \log \left(2 + \frac{t}{s} \right). \quad (7.5)$$

But from (2.1) for $s < t$ and small t , and then from (2.5),

$$\begin{aligned} \frac{u}{v} / \frac{t}{s} &= \frac{a_u}{a_v} \leq \left(\frac{Q(a_u)}{Q(a_v)} \right)^{1/2} \\ &\leq C_1 \left(\frac{uT(a_u)^{-1/2}}{vT(a_v)^{-1/2}} \right)^{1/2} \leq C_2 \left(\frac{uT(a_{\beta(u)})^{-1/2}}{vT(a_{\beta(v)})^{-1/2}} \right)^{1/2} \leq C_3 \left(\frac{u}{v} \right)^{1/2} \end{aligned}$$

as $\beta(u) \geq \beta(v)$. So

$$\left(\frac{u}{v} \right)^{1/2} \leq C_4 \frac{t}{s} \quad (7.6)$$

and we have (7.5).

(b) Now if $x \geq 0$,

$$\begin{aligned} \left| 1 - \frac{x}{\sigma(s)} \right| &\leq \left| 1 - \frac{x}{\sigma(t)} \right| + \frac{x}{\sigma(t)} \left| 1 - \frac{\sigma(t)}{\sigma(s)} \right| \\ &\leq \left| 1 - \frac{x}{\sigma(t)} \right| + \left(\left| 1 - \frac{x}{\sigma(t)} \right| + 1 \right) \left| 1 - \frac{\sigma(t)}{\sigma(s)} \right|. \end{aligned}$$

Using (a) of this lemma, we obtain

$$\left| 1 - \frac{x}{\sigma(s)} \right|^{1/2} \leq C_{12} \Phi_t(x) \sqrt{\log \left(2 + \frac{t}{s} \right)}.$$

Since $\sigma(s) \geq \sigma(t)$, also

$$T(\sigma(s))^{-1/2} \leq C_{13} T(\sigma(t))^{-1/2}.$$

So (7.2) follows. \square

We turn to the proof of Theorem 1.3. We provide full proofs only where the details are significantly different, and otherwise refer back. We begin with an analogue of Lemma 6.1 for $L_n[f]$ of (6.11).

Lemma 7.2

$$\begin{aligned} \|(f - L_n[f])W\|_{L_p(\mathbb{R})} &\leq C_1 \left[\sup_{\substack{0 < h \leq a_{3n}/(3n) \\ 0 < \tau \leq L}} \|W \Delta_{\tau h \Phi_h(x)}^r(f, x, \mathbb{R})\|_{L_p[-a_n, a_n]} \right. \\ &\quad \left. + \|fW\|_{L_p(|x| \geq a_n)} \right]. \end{aligned} \quad (7.7)$$

Here L is independent of f, n .

Proof

We do this for $p < \infty$. Recall that the crux of Lemma 6.1 is estimation of

$$\begin{aligned} \Delta_{jn} : &= \int_{I_{jn}} |f - p_j|^p W^p \leq C_1 \Omega_{r,p}(f, |I_{jn}^*|, I_{jn}^*)^p W^p(\tau_{jn}) \\ &\leq \frac{C_2}{|I_{jn}^*|} \int_{I_{jn}^*} \int_0^{|I_{jn}^*|} |W \Delta_s^r(f, x, I_{jn}^*)|^p ds dx. \end{aligned} \quad (7.8)$$

We now choose $L > 0$ such that for $0 < h \leq 1$,

$$\sup_{x \in \mathbb{R}} \frac{\frac{h}{L} \Phi_{\frac{h}{L}}(x)}{h \Phi_h(x)} \leq \frac{1}{2}. \quad (7.9)$$

This is possible by (7.2). Now we choose

$$\delta_{n,k}(x) := L^{1-k} \frac{a_{3n}}{3n} \Phi_{L^{1-k} \frac{a_{3n}}{3n}}(x), \quad k \geq 1.$$

Note that by (7.9),

$$\sup_{x \in \mathbb{R}} \frac{\delta_{n,k+1}(x)}{\delta_{n,k}(x)} \leq \frac{1}{2}. \quad (7.10)$$

In view of (6.6), (3.6) and (3.7), we may assume that L is so large that uniformly in $n, j, x \in I_{jn}^*$,

$$|I_{jn}^*| \leq L \frac{a_{3n}}{3n} \Phi_{\frac{a_{3n}}{3n}}(x) = L \delta_{n,1}(x); \quad |I_{jn}^*| \sim \delta_{n,1}(x).$$

Then from (7.8),

$$\begin{aligned} \Delta_{jn} &\leq C_4 \int_{I_{jn}^*} \int_0^{L\delta_{n,1}(x)} \frac{1}{\delta_{n,1}(x)} \left| W \Delta_s^r(f, x, I_{jn}^*) \right|^p ds dx \\ &= C_4 \int_{I_{jn}^*} \sum_{k=1}^{\infty} \int_{L\delta_{n,k+1}(x)}^{L\delta_{n,k}(x)} \frac{1}{\delta_{n,1}(x)} \left| W \Delta_s^r(f, x, I_{jn}^*) \right|^p ds dx \\ &= C_4 \int_{I_{jn}^*} \sum_{k=1}^{\infty} \int_{L\delta_{n,k+1}(x)/\delta_{n,k}(x)}^L \frac{\delta_{n,k}(x)}{\delta_{n,1}(x)} \left| W \Delta_{\tau\delta_{n,k}(x)}^r(f, x, I_{jn}^*) \right|^p d\tau dx \\ &\leq C_4 \int_{I_{jn}^*} \sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^{k-1} \int_0^L \left| W \Delta_{\tau\delta_{n,k}(x)}^r(f, x, I_{jn}^*) \right|^p d\tau dx. \end{aligned}$$

Then

$$\begin{aligned} \sum_{j=0}^{n-1} \Delta_{jn} &\leq C_4 \int_{-a_n}^{a_n} \sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^{k-1} \int_0^L \left| W \Delta_{\tau\delta_{n,k}(x)}^r(f, x, \mathbb{R}) \right|^p d\tau dx \\ &\leq 2C_4 \sup_{\substack{0 < h \leq a_{3n}/(3n) \\ 0 < \tau \leq L}} \int_{-a_n}^{a_n} \left| W \Delta_{\tau h \Phi_h(x)}^r(f, x, \mathbb{R}) \right|^p dx. \end{aligned}$$

The rest of the proof is as before. \square

The analogue of Lemma 6.2 is

Lemma 7.3

$$\|(L_n[f] - P_n[f])W\|_{L_p(\mathbb{R})}$$

$$\leq C_1 \left\{ \sup_{\substack{0 < h \leq a_{3n}/(3n) \\ 0 < \tau \leq L}} \left\| W \Delta_{\tau h \Phi_h(x)}^r(f, x, \mathbb{R}) \right\|_{L_p[-a_n, a_n]} + \|fW\|_{L_p(I_{\delta_n}^*)} \right\}.$$

Proof

This is exactly the same as the proof of Lemma 6.2, except that we substitute for (6.19) the estimate of Lemma 7.2. \square

Proof of Theorem 1.3

This follows from Lemma 7.2 and 7.3 exactly as Theorem 1.2 followed from Lemma 6.1 and 6.2. \square

Finally, we briefly show that under some additional conditions on Q , we can use the simpler modulus

$$\begin{aligned} \omega_{r,p}^\#(f, W, t) &= \sup_{0 < h \leq t} \left\| W \Delta_{Lh \Phi_h(x)}^r(f, x, \mathbb{R}) \right\|_{L_p(|x| \leq \sigma(2h))} \\ &\quad + \inf_{P \in \mathcal{P}_{r-1}} \|(f - P)W\|_{L_p(|x| \geq \sigma(4t))}. \end{aligned} \quad (7.11)$$

We shall assume in addition to $W \in \mathcal{E}$, that Q'' exists and is non-negative in $(0, \infty)$, and

$$\frac{Q''(x)}{Q'(x)} \sim \frac{Q'(x)}{Q(x)}, \quad x \in (0, \infty). \quad (7.12)$$

Moreover, we assume that

$$|T'(x)| \leq C_1 \frac{T^2(x)}{x}, \quad x \geq C_1. \quad (7.13)$$

Using (7.12) and the methods of proof of Lemma 2.2 in [13,p.209], we obtain

$$\frac{a'_u}{a_u} \sim \frac{1}{uT(a_u)}, \quad u \geq C_2 \quad (7.14)$$

and hence

$$\frac{d}{du} \left(\frac{a_u}{u} \right) \sim -\frac{a_u}{u^2}, \quad u \geq C_2. \quad (7.15)$$

Since $u \rightarrow \frac{a_u}{u}$ is then strictly decreasing for large u , we obtain the identity

$$\sigma \left(\frac{a_u}{u} \right) = a_u, \quad u \geq C_3. \quad (7.16)$$

Differentiating this, and using (7.14), (7.15) leads to

$$\frac{\sigma'(t)}{\sigma(t)} \sim -\frac{1}{tT(\sigma(t))}, \quad 0 < t \leq C_4 \quad (7.17)$$

and then using (7.13), we obtain

$$\left| t \frac{d}{dt} T(\sigma(t)) \right| \leq C_5 T(\sigma(t)), \quad 0 < t \leq C_4. \quad (7.18)$$

These last two bounds easily give

$$\left| \frac{d}{dt} [t\Phi_t(x)] \right| \leq C_5 \Phi_t(x) \quad (7.19)$$

for

$$0 < t \leq C_5; \quad \left| 1 - \frac{|x|}{\sigma(t)} \right| \geq \frac{\varepsilon}{T(\sigma(t))}. \quad (7.20)$$

Here ε is any fixed positive number. We now estimate Δ_{jn} a little differently from the way we proceeded after (7.8). Let us make the substitution $s = Lt\Phi_t(x)$ in the right-hand side of (7.8) and keep our choice of $L, \delta_{n,1}(x)$ to deduce that

$$\begin{aligned} \Delta_{jn} &\leq C_6 \int_{I_{jn}^*} \int_0^{a_{3n}/(3n)} \frac{1}{\delta_{n,1}(x)} \left| W \Delta_{Lt\Phi_t(x)}^r(f, x, I_{jn}^*) \right|^p \left| \frac{d}{dt} [t\Phi_t(x)] \right| dt dx \\ &\leq \frac{C_7 3n}{a_{3n}} \int_{I_{jn}^*} \int_0^{a_{3n}/(3n)} \sqrt{\log \left(2 + \frac{a_{3n}}{3nt} \right)} \left| W \Delta_{Lt\Phi_t(x)}^r(f, x, I_{jn}^*) \right|^p dt dx \end{aligned}$$

by (7.19) and (7.2). In applying (7.19) we must ensure that the range conditions in (7.20) must hold for $x \in I_{jn}^*$ and $t \leq a_{3n}/(3n)$. In fact if $|x| \leq a_n$, then

$$\begin{aligned} 1 - \frac{|x|}{\sigma(t)} &\geq 1 - \frac{a_n}{\sigma(a_{3n}/(3n))} \geq 1 - \frac{a_n}{a_{3n(1+o(1))}} \\ &\geq C_8 T(a_n)^{-1} \geq C_9 T(\sigma(t))^{-1} \end{aligned}$$

by (3.2), (3.3), then (2.7) and then (2.6). Thus

$$\begin{aligned} \sum_{j=0}^{n-1} \Delta_{jn} &\leq \frac{C_8 3n}{a_{3n}} \int_{-a_n}^{a_n} \int_0^{a_{3n}/(3n)} \sqrt{\log\left(2 + \frac{a_{3n}}{3nt}\right)} \left|W\Delta_{Lt\Phi_t(x)}^r(f, x, \mathbb{R})\right|^p dt dx \\ &\leq C_8 \sup_{0 < t \leq a_{3n}/(3n)} \int_{-a_n}^{a_n} \left|W\Delta_{Lt\Phi_t(x)}^r(f, x, \mathbb{R})\right|^p dx \int_0^1 \sqrt{\log\left(2 + \frac{1}{s}\right)} ds. \end{aligned}$$

So under the additional conditions on Q we obtain

$$E_n[f]_{W,p} \leq C_9 \omega_{r,p}^\# \left(f, W, C_{10} \frac{a_n}{n}\right). \quad (7.21)$$

We note that these additional conditions (7.12) and (7.13) are certainly satisfied for $W_{k,\alpha}$ of (1.6).

References

- [1] Yu. A. Brudnyi, Approximation of functions by algebraic polynomials, *Izvestiya AN SSSR, ser.mathem.*, 32, N4 (1968), 780-787.
- [2] J. Clunie, T. Kövari, On Integral Functions having prescribed asymptotic growth II, *Canad. J. Math.*, 20(1968), 7-20.
- [3] S. B. Damelin, Converse and Smoothness Theorems for Erdős Weights in $L_p(0 < p \leq \infty)$, to appear in *J. Approx. Th.*
- [4] R. A. DeVore and V. A. Popov, Interpolation of Besov Spaces, *Trans. Amer. Math. Soc.*, 305(1988), 397-414.
- [5] R. A. DeVore, D. Leviatan, X. M. Yu, Polynomial Approximation in $L_p(0 < p < 1)$, *Constructive Approximation*, 8(1992), 187-201.
- [6] Z. Ditzian and D. S. Lubinsky, Jackson and Smoothness Theorems for Freud Weights in $L_p, 0 < p \leq \infty$, *Constructive Approximation*, 13(1997), 99-152.
- [7] Z. Ditzian and V. Totik, *Moduli of Smoothness*, Springer Series in Computational Mathematics, Vol. 9, Springer, Berlin, 1987.

- [8] G. Freud, *Orthogonal Polynomials*, Pergamon Press/ Akademiai Kiado, Budapest, 1971.
- [9] M. von Golitschek, G. G. Lorentz and Y. Makovoz, *Asymptotics of Weighted Polynomials*, in *Progress in Approximation Theory*, A.A. Gonchar and E.B. Saff (eds.), Springer Series in Computational Mathematics, Vol. 19, Springer, New York, 1992, 431- - 451.
- [10] S. Jansche and R. L. Stens, *Best weighted polynomial approximation on the real line: a functional analytic approach*, *J. Comp. Appl. Math.*, **40**(1992), 199- -213.
- [11] D. Leviatan and X. M. Yu, *Shape Preserving Approximation by Polynomials in L_p* , manuscript.
- [12] A. L. Levin and D. S. Lubinsky, *Christoffel functions, orthogonal polynomials, and Nevai's conjecture for Freud weights*, *Constructive Approximation*, **8**(1992), 463- -535.
- [13] A. L. Levin, D. S. Lubinsky and T. Z. Mthembu, *Christoffel functions and orthogonal polynomials for Erdős weights on $(-\infty, \infty)$* , *Rendiconti di Matematica e delle sue Applicazioni (di Roma)*, **14**(1994), 199- -289.
- [14] D. S. Lubinsky, *Strong Asymptotics for Extremal Errors and Polynomials Associated with Erdős Type Weights*, *Pitman Research Notes in Mathematics*, Vol. 202, Longman, Harlow, Essex, 1989.
- [15] D. S. Lubinsky, *An update on orthogonal polynomials and weighted approximation on the real line*, *Acta Applic. Math.*, **33**(1993), 121- -164.
- [16] D. S. Lubinsky, *Ideas of Weighted Polynomial Approximation on $(-\infty, \infty)$* , *Approximation and Interpolation*, World Scientific, (1995), pp 371-396
- [17] D. S. Lubinsky and T. Z. Mthembu, *Orthogonal expansions and the error of weighted polynomial approximation for Erdős weights*, *Numer. Funct. Anal. and Optimiz.*, **13**(1992), 327- -347.

- [18] H. N. Mhaskar and E. B. Saff, Extremal problems for polynomials with exponential weights, *Trans. Amer. Math. Soc.*, **285**(1984), 203- -234.
- [19] H. N. Mhaskar and E. B. Saff, Where does the sup-norm of a weighted polynomial live?, *Constructive Approximation*, **1**(1985), 71- -91.
- [20] H. N. Mhaskar and E. B. Saff, Where does the L_p norm of a weighted polynomial live?, *Trans. Amer. Math. Soc.*, **303**(1987), 109- -124.
- [21] P. Nevai, Geza Freud, Orthogonal polynomials and Christoffel functions: a case study, *J. Approx. Theory*, **48**(1986), 3- -167.
- [22] P. Nevai (ed.), *Orthogonal Polynomials, Theory and Practice*, NATO ASI Series, Vol. 294, Kluwer, Dordrecht, 1990.
- [23] P. Oswald, Ungleichungen Vom Jackson-Typ für die algebraische beste Approximation in L_p , *J. Approx. Theory*, **23**, N2, (1978), 113-136.
- [24] P. P. Petrushev and V. Popov, *Rational Approximation of Real Functions*, Cambridge University Press, Cambridge, 1987.
- [25] E. B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer, to appear.
- [26] V. Totik, *Weighted Approximation with Varying Weight*, Springer Lecture Notes in Mathematics, Vol. 1569, Springer, Berlin, 1994.