

## Problem Set 3

Due: Tuesday, January 27 at the beginning of class

We will assume rings have identity and ring homomorphisms are unital (send 1 to 1) unless stated otherwise. Read 10.3 Problem 27. Turn in Problems 1–10.

**Problem 1.** An element  $e \in R$  is called a *central idempotent* if  $e^2 = e$  and  $er = re$  for all  $r \in R$ . If  $e$  is a central idempotent in  $R$ , prove that  $M = eM \oplus (1 - e)M$ .

**Problem 2.** An element  $m$  of the  $R$ -module  $M$  is called a *torsion element* if  $rm = 0$  for some nonzero element  $r \in R$ . The set of torsion elements is denoted

$$\text{Tor}(M) = \{m \in M \mid rm = 0 \text{ for some nonzero } r \in R\}.$$

- (a) Prove that if  $R$  is an integral domain then  $\text{Tor}(M)$  is a submodule of  $M$  (called the *torsion submodule* of  $M$ ).
- (b) Give an example of a ring  $R$  and an  $R$ -module  $M$  such that  $\text{Tor}(M)$  is not a submodule.
- (c) If  $R$  has zero divisors show that every nonzero  $R$ -module has nonzero torsion elements.

**Problem 3.** Let  $\phi : M \rightarrow N$  be an  $R$ -module homomorphism. Prove that  $\phi(\text{Tor}(M)) \subseteq \text{Tor}(N)$ .

**Problem 4.** A *torsion  $R$ -module* is an  $R$ -module  $M$  such that  $\text{Tor}(M) = M$ . For an  $R$ -module  $M$ , the *annihilator of  $M$  in  $R$*  is

$$\text{Ann}_R(M) := \{r \in R \mid rm = 0 \text{ for all } m \in M\}.$$

Give an example of an integral domain  $R$  and a nonzero torsion  $R$ -module  $M$  such that  $\text{Ann}_R(M) = 0$ . Prove that if  $N$  is a finitely generated torsion  $R$ -module then  $\text{Ann}_R(N) \neq 0$ .

**Problem 5.** Let  $R = \mathbb{Z}[x]$  and let  $M = (2, x)$  be the ideal generated by 2 and  $x$ , considered as a submodule of  $R$ . Show that  $\{2, x\}$  is not a basis of  $M$ . Show that the rank of  $M$  is 1 but that  $M$  is not free of rank 1.

**Problem 6.** Let  $\mathbb{F}_2$  denote the field with 2 elements. Determine (with proof) which of the following pairs of rings are isomorphic:

- $\mathbb{F}_2[x]/(x^3 + x^2 + x + 1)$
- $\mathbb{F}_2[x]/(x^3 + x + 1)$
- $\mathbb{F}_2[x]/(x^3 + x^2 + 1)$ .

Avoid using facts about finite fields we have not proved, though you can use them if you prove them.

**Problem 7.** Determine all nilpotent elements of  $M_2(\mathbb{C})$ .

**Problem 8.** Let  $F$  be a field. Give a simple description of the set of zero divisors of  $M_n(F)$  in terms of concepts from linear algebra.

**Problem 9.** Show that if  $R = \mathbb{Z}$ ,  $I = \mathbb{Z}_{>0}$ , and  $M_i = \mathbb{Z}/i\mathbb{Z}$  for each  $i \in I$ , then  $\bigoplus_{i \in I} M_i$  is not isomorphic to  $\prod_{i \in I} M_i$ .

**Problem 10.** Let  $R$  be a commutative ring. Prove that  $R^n \cong R^m$  if and only if  $n = m$ , i.e., two free  $R$ -modules of finite rank are isomorphic if and only if they have the same rank.