

## Problem Set 9

Due: Tuesday, March 10 at the beginning of class

We will assume rings have identity and ring homomorphisms are unital (send 1 to 1) unless stated otherwise. Turn in Problems 1–10.

**Problem 1.** Prove that elements  $x$  and  $y$  are conjugate in a group  $G$  if and only if  $\chi(x) = \chi(y)$  for all irreducible characters  $\chi$  of  $G$ .

**Problem 2.** Let  $H$  and  $K$  be finite groups and  $V$  be a  $\mathbb{C}$ -vector space. Let  $G = H \times K$  and let  $\phi : H \rightarrow GL(V)$  be an irreducible representation of  $H$  with character  $\chi$ . Then  $G \xrightarrow{\pi_H} H \xrightarrow{\phi} GL(V)$  gives an irreducible representation of  $G$ , where  $\pi_H$  is the natural projection; the character  $\tilde{\chi}$  of this representation is  $\tilde{\chi}((h, k)) = \chi(h)$ . Likewise any irreducible character  $\psi$  of  $K$  gives an irreducible character  $\tilde{\psi}$  of  $G$  with  $\tilde{\psi}((h, k)) = \psi(k)$ .  
Prove that the product  $\tilde{\chi}\tilde{\psi}$  is an irreducible character of  $G$ .

**Problem 3.** Show that the element “ $2 \otimes 1$ ” is 0 in  $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$  but is nonzero in  $2\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ .

**Problem 4.** An element of a tensor product  $M \otimes_R N$  is a *simple tensor* if it is of the form  $m \otimes n$  for  $m \in M, n \in N$ . Let  $F$  be a field and let  $V$  be an  $n$ -dimensional  $F$ -vector space.

- The vector space  $V \otimes_F V$  can be identified with  $M_n(F)$  via  $e_i \otimes e_j \mapsto E_{ij}$ , where  $e_1, \dots, e_n$  is a basis of  $V$  and  $E_{ij}$  is the matrix with a 1 in the  $i, j$ th spot and 0's elsewhere. Express what it means for an element of  $V \otimes_F V$  to be a simple tensor in terms of concepts from linear algebra.
- Suppose  $n \geq 2$ . Show that the element  $e_1 \otimes e_2 + e_2 \otimes e_1$  in  $V \otimes_F V$  is not a simple tensor.

**Problem 5.** Show that  $\mathbb{Q} \otimes_{\mathbb{Z}} \mathbb{Q}$  and  $\mathbb{Q} \otimes_{\mathbb{Q}} \mathbb{Q}$  are isomorphic left  $\mathbb{Q}$ -modules.

**Problem 6.** Suppose  $R$  is commutative and let  $I$  and  $J$  be ideals of  $R$ , so  $R/I$  and  $R/J$  are naturally  $R$ -modules.

- Prove that every element of  $R/I \otimes_R R/J$  can be written as a simple tensor of the form  $(1 \bmod I) \otimes (r \bmod J)$ .
- Prove that there is an  $R$ -module isomorphism  $R/I \otimes_R R/J \cong R/(I + J)$  mapping  $(r \bmod I) \otimes (r' \bmod J)$  to  $rr' \bmod (I + J)$ .

**Problem 7.** Let  $A$  be any ring, let  $L$  be any left  $A$ -module, and let  $L^{\oplus n}$  be the direct sum of  $n$  copies of  $L$  with itself.

- Prove the ring isomorphism  $\text{Hom}_A(L^{\oplus n}, L^{\oplus n}) \cong M_n(D)$ , where  $D = \text{Hom}_A(L, L)$ .
- Deduce that if  $L$  is an irreducible  $A$ -module, then  $\text{Hom}_A(L^{\oplus n}, L^{\oplus n})$  is isomorphic to a matrix ring over a division ring.

**Problem 8.** The alternating group  $A_4$  is a subgroup of  $S_4$ , hence  $\mathbb{C}A_4$  is a subalgebra of  $\mathbb{C}S_4$ . Therefore any  $\mathbb{C}S_4$ -module is a  $\mathbb{C}A_4$ -module by restriction. For each irreducible  $\mathbb{C}S_4$ -module  $V$ , determine the decomposition of  $V$ , regarded as a  $\mathbb{C}A_4$ -module, into irreducibles. Feel free to use the character table for  $A_4$  in 19.1.

**Problem 9.** By the Artin-Wedderburn Theorem and the character table of  $\mathcal{S}_3$ , we have the following isomorphism of rings:

$$\mathbb{C}\mathcal{S}_3 \cong M_1(\mathbb{C}) \times M_2(\mathbb{C}) \times M_1(\mathbb{C}).$$

Determine explicitly the elements  $\left(\begin{bmatrix} 1 \\ \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, [0]\right)$ ,  $\left(\begin{bmatrix} 0 \\ \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, [0]\right)$ ,  $\left(\begin{bmatrix} 0 \\ \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, [1]\right)$  on the right hand side in terms of the basis  $\{\pi : \pi \in \mathcal{S}_3\}$  of  $\mathbb{C}\mathcal{S}_3$ .

This is equivalent to finding three elements  $z_1, z_2, z_3$  in the center of  $\mathbb{C}\mathcal{S}_3$  such that

- $z_1 + z_2 + z_3 = 1$ ,
- $z_i^2 = z_i$ ,
- $z_i z_j = z_j z_i = 0$  for  $i \neq j$ .

**Problem 10.** Let  $F$  be a field and  $G$  a finite group. Let  $V$  and  $W$  be  $FG$ -modules.

(a) Consider the  $F$ -vector space  $V \otimes_F W$ . Show that if  $g \in G$  acts on  $V \otimes_F W$  by

$$g \cdot (v \otimes w) := gv \otimes gw \quad \text{for every } v \in V, w \in W,$$

then this gives  $V \otimes_F W$  the structure of an  $FG$ -module.

(b) Prove that the character of  $V \otimes_F W$  is given by  $\chi_{V \otimes_F W}(g) = \chi_V(g)\chi_W(g)$  for every  $g \in G$ .