

On the Kottwitz conjecture for local shtuka spaces

David Hansen, Tasho Kaletha, and Jared Weinstein

May 7, 2021

Abstract

Kottwitz’s conjecture describes the contribution of a supercuspidal representation to the cohomology of a local Shimura variety in terms of the local Langlands correspondence. A natural extension of this conjecture concerns Scholze’s more general spaces of local shtukas. Using a new Lefschetz-Verdier trace formula for v-stacks, we prove the extended conjecture, disregarding the action of the Weil group, and modulo a virtual representation whose character vanishes on the locus of elliptic elements. As an application, we show that for an irreducible smooth representation of an inner form of GL_n , the L -parameter constructed by Fargues-Scholze agrees with the usual semisimplified parameter arising from local Langlands.

Contents

1	Introduction	3
1.1	Remarks on the proof, and relation with prior work	5
2	Review of the objects appearing in Kottwitz’s conjecture	9
2.1	Basic notions	9
2.2	Construction of $\delta_{\pi,\rho}$ in a special case	9
2.3	Construction of $\delta_{\pi,\rho}$ in the general case	10
2.4	Spaces of local shtukas and their cohomology	12
3	Transfer of conjugation-invariant functions from $G(F)$ to $G_b(F)$	15
3.1	The space of strongly regular conjugacy classes in $G(F)$	15
3.2	Hecke correspondences	16
3.3	Interaction with orbital integrals	19
3.4	Proof of Theorem 3.2.9	22
4	The Lefschetz-Verdier trace formula for v-stacks	25
4.1	Small v-stacks and the six-functor formalism	26
4.2	Artin v-stacks	28
4.3	The category of cohomological correspondences	29

T.K. was supported in part by NSF grant DMS-1161489 and a Sloan Fellowship. J.W. was supported by NSF grant DMS-1303312 and a Sloan Fellowship.

4.4	Examples	34
4.5	A Künneth theorem for characteristic classes	35
4.6	The case of $[X/G]$ for G smooth	41
5	Local terms on the B_{dR}-affine Grassmannian	45
5.1	The main result	45
5.2	Strategy of proof	47
5.3	Local terms and base change	49
5.4	Independence of g	51
5.5	Degeneration to characteristic p	52
5.6	Local terms on the Witt vector affine Grassmannian	53
6	Application to the Hecke stacks	55
6.1	Bun_G , the local and global Hecke stacks, and their relation to shtuka spaces	55
6.2	The inertia stack of the Hecke stack; admissibility of elliptic fixed points	58
6.3	Transfer of distributions from G_b to G	60
6.4	Hecke operators on Bun_G and the cohomology of shtuka spaces	63
6.5	Proof of Theorem 1.0.2	69
6.6	Application to inner forms of GL_n	73
A	Endoscopy	74
A.1	Endoscopic character relations	74
A.2	The Kottwitz sign	77
B	Elementary Lemmas	77
B.1	Homological algebra	77
B.2	Sheaves on locally profinite sets	79
C	Some representation theory	80
C.1	Non-elliptic representations	81
C.2	Integral representations and parabolic inductions	81

1 Introduction

Let F be a finite extension of the field \mathbf{Q}_p of p -adic numbers, and let G be a connected reductive group defined over F . Scholze [SW20, §23] has introduced a tower of moduli spaces of mixed-characteristic shtukas

$$\mathrm{Sht}_{G,b,\mu} = \varprojlim_K \mathrm{Sht}_{G,b,\mu,K}$$

depending on a σ -conjugacy class of $b \in G(\check{F})$ (where \check{F} is the completion of the maximal unramified extension of F) and on a conjugacy class of cocharacters $\mu: \mathbf{G}_m \rightarrow G$ defined over \bar{F} . Here K ranges over open compact subgroups of $G(F)$. The inverse limit $\mathrm{Sht}_{G,b,\mu,K}$ is a locally spatial diamond defined over $\mathrm{Spd} \check{E}$, where E is the field of definition of the conjugacy class of μ .

When μ is minuscule, $\mathrm{Sht}_{G,b,\mu,K}$ is the diamond associated to a rigid-analytic variety $\mathcal{M}_{G,b,\mu,K}$ [SW20, §24]. The latter is a *local Shimura variety*, whose general existence was conjectured in [RV14]. The theory of Rapoport-Zink spaces [RZ96] provides instances of $\mathcal{M}_{G,b,\mu,K}$ admitting a moduli interpretation, as the generic fiber of a deformation space of p -divisible groups.

The *Kottwitz conjecture* [Rap95, Conjecture 5.1], [RV14, Conjecture 7.3] relates the cohomology of $\mathcal{M}_{G,b,\mu,K}$ to the local Langlands correspondence, in the case that b lies in the unique basic class in $B(G, \mu)$. There is a natural generalization of this conjecture for $\mathrm{Sht}_{G,b,\mu,K}$, as we now explain.

Let G_b the inner form of G associated to b . The tower $\mathrm{Sht}_{G,b,\mu,K}$ admits commuting actions of $G_b(F)$ and $G(F)$. The action of $G_b(F)$ preserves each $\mathrm{Sht}_{G,b,\mu,K}$, whereas the action of $g \in G(F)$ sends $\mathrm{Sht}_{G,b,\mu,K}$ to $\mathrm{Sht}_{G,b,\mu, gKg^{-1}}$. There is furthermore a (not necessarily effective) Weil descent datum on this tower from \check{E} down to E .

Let ℓ be a prime distinct from p . The Satake equivalence produces an object \mathcal{S}_μ in the derived category of étale \mathbf{Z}_ℓ -sheaves on $\mathrm{Sht}_{G,b,\mu,K}$; this is compatible with the actions of $G(F)$ and $G_b(F)$ on the tower. Let C be the completion of an algebraic closure of \check{E} . For a smooth representation ρ of $G_b(F)$ with coefficients in $\overline{\mathbf{Q}}_\ell$, we define:

$$R\Gamma(G, b, \mu)[\rho] = \varinjlim_K R\mathrm{Hom}_{G_b(F)}(R\Gamma_c(\mathrm{Sht}_{G,b,\mu,K,C}, \mathcal{S}_\mu), \rho).$$

Then $R\Gamma(G, b, \mu)[\rho]$ lies in the derived category of smooth representations of $G(F) \times W_E$ with coefficients in $\overline{\mathbf{Q}}_\ell$, where W_E is the Weil group. Informally, this is the ρ -isotypic component of the cohomology of the tower $\mathrm{Sht}_{G,b,\mu}$.

A recent result of Fargues-Scholze [FS21, Corollary I.7.3] states that if ρ is finite length and admissible, then $R\Gamma(G, b, \mu)[\rho]$ is a complex of finite length admissible representations of $G(F)$ admitting a smooth action of W_E .

Let $\mathrm{Groth}(G_b(F))$ be the Grothendieck group of the category of finite length admissible representations of $G_b(F)$ with $\overline{\mathbf{Q}}_\ell$ coefficients. Also, let $\mathrm{Groth}(G(F) \times W_E)$ be the Grothendieck group of the category of finite length admissible representations of $G(F)$ with $\overline{\mathbf{Q}}_\ell$ coefficients, which come equipped with a smooth action of W_E commuting with the $G(F)$ -action. Following [Shi11] we define a map

$$\mathrm{Mant}_{b,\mu}: \mathrm{Groth}(G_b(F)) \rightarrow \mathrm{Groth}(G(F) \times W_E)$$

(for ‘‘Mantovan’’, referencing [Man04]) sending ρ to the Euler characteristic of $R\Gamma(G, b, \mu)[\rho]$.

The Kottwitz conjecture (appropriately generalized) describes $\text{Mant}_{b,\mu}(\rho)$ in terms of the local Langlands correspondence. The complex dual groups of G and G_b are canonically identified, and we write \widehat{G} for either. Let ${}^L G = \widehat{G} \rtimes W_F$ be the L -group. The basic form of the local Langlands conjecture predicts that the set of isomorphism classes of essentially square-integrable representations of $G(F)$ (resp., $G_b(F)$) is partitioned into L -packets $\Pi_\phi(G)$ (resp., $\Pi_\phi(G_b)$), and that each such packet is indexed by a discrete Langlands parameter $\phi : W_F \times \text{SL}_2(\mathbf{C}) \rightarrow {}^L G$. When ϕ is trivial on $\text{SL}_2(\mathbf{C})$, we say ϕ is supercuspidal; in this case it is expected that the packets $\Pi_\phi(G)$ and $\Pi_\phi(G_b)$ consist entirely of supercuspidal representations.

Our generalized Kottwitz conjecture is conditional on the refined local Langlands correspondence for supercuspidal L -parameters, in the formulation of [Kal16a, Conjecture G]. In particular, it relies crucially on the endoscopic character identities satisfied by L -packets. These are reviewed in Appendix A. At the moment a construction of the refined correspondence for supercuspidal parameters is known when p does not divide the order of the Weyl group of G , cf. [Kal19a], [Kal19b], and the character identities are known for regular supercuspidal parameters under stricter assumptions on p , cf. [FKS19].

Let $S_\phi = \text{Cent}(\phi, \widehat{G})$. For any $\pi \in \Pi_\phi(G)$ and $\rho \in \Pi_\phi(G_b)$ the refined form of the local Langlands conjecture implies the existence of an algebraic representation $\delta_{\pi,\rho}$ of S_ϕ , which can be thought of as measuring the relative position of π and ρ . The conjugacy class of μ determines by duality a conjugacy class of weights of \widehat{G} ; we denote by r_μ the irreducible representation of \widehat{G} of highest weight μ . There is a natural extension of r_μ to ${}^L G_E$, the L -group of the base change of G to E [Kot84a, Lemma 2.1.2]. Write $r_\mu \circ \phi_E$ for the representation of $S_\phi \times W_E$ given by

$$r_\mu \circ \phi_E(s, w) = r_\mu(s \cdot \phi(w)).$$

Conjecture 1.0.1. *Let $\phi : W_F \rightarrow {}^L G$ be a supercuspidal Langlands parameter. Given $\rho \in \Pi_\phi(G_b)$, we have the following equality in $\text{Groth}(G(F) \times W_E)$:*

$$\text{Mant}_{b,\mu}(\rho) = \sum_{\pi \in \Pi_\phi(G)} \pi \boxtimes \text{Hom}_{S_\phi}(\delta_{\pi,\rho}, r_\mu \circ \phi_E). \quad (1.0.1)$$

This conjecture is more general than the formulation of Kottwitz's conjecture in [Rap95] and [RV14], in that two conditions are removed. The first is that we are allowing the cocharacter μ to be non-minuscule – this is what requires passage from the local Shimura varieties $\mathcal{M}_{G,b,\mu}$ to the local shtuka spaces $\text{Sht}_{G,b,\mu}$. The second is that we do not require G to be a B -inner form of its quasi-split inner form G^* . This condition, reviewed in §2.2, has the effect of making the definition of $\delta_{\pi,\rho}$ straightforward. To remove it, we use the formulation of the refined local Langlands correspondence [Kal16a, Conjecture G] based on the cohomology sets $H^1(u \rightarrow W, Z \rightarrow G)$ of [Kal16b]. The definition of $\delta_{\pi,\rho}$ in this setting is a bit more involved and is given in §2.3, see Definition 2.3.2.

We now present our main theorem.

Theorem 1.0.2. *Assume the refined local Langlands correspondence [Kal16a, Conjecture G]. Let $\phi : W_F \times \text{SL}_2 \rightarrow {}^L G$ be a discrete Langlands parameter with coefficients in $\overline{\mathbf{Q}}_\ell$, and let $\rho \in \Pi_\phi(G_b)$ be a member of its L -packet. After ignoring the action of W_E , we have an equality in $\text{Groth}(G(F))$:*

$$\text{Mant}_{b,\mu}(\rho) = \sum_{\pi \in \Pi_\phi(G)} [\dim \text{Hom}_{S_\phi}(\delta_{\pi,\rho}, r_\mu)] \pi + \text{err},$$

where $\text{err} \in \text{Groth}(G(F))$ is a virtual representation whose character vanishes on the locus of elliptic elements of $G(F)$.

If the packet $\Pi_\phi(G)$ consists entirely of supercuspidal representations and the semisimple L -parameter φ_ρ associated with ρ as in [FS21, §I.9.6] is supercuspidal, then in fact $\text{err} = 0$.

Of course we expect that $\varphi_\rho = \phi^{\text{ss}}$, so that if ϕ is supercuspidal, the error term should vanish. In that case we obtain Conjecture 1.0.1 modulo the action of W_E . However, for a discrete but non-supercuspidal ϕ , the error term is often provably nonzero, cf. [Ima].

The shtukas appearing in our work have only one “leg”. Scholze defines moduli spaces of mixed-characteristic shtukas $\text{Sht}_{G,b,\{\mu_i\}}$ with arbitrarily many legs, fibered over a product $\prod_{i=1}^r \text{Spd } \check{E}_i$. It is possible to extend Conjecture 1.0.1 and Theorem 1.0.2 to this setting as well; we leave the details to the interested reader.

Theorem 1.0.2 has an application to the local Langlands correspondence.

Theorem 1.0.3. *Let G be any inner form of GL_n/F , and let π be an irreducible smooth representation of $G(F)$. Then the L -parameter φ_π associated with π as in [FS21, §I.9] agrees with the usual semisimplified L -parameter attached to π .*

1.1 Remarks on the proof, and relation with prior work

Ultimately, Theorem 1.0.2 is proved by an application of a Lefschetz-Verdier trace formula. Let us illustrate the idea in the Lubin-Tate case: say $F = \mathbf{Q}_p$, $G = \text{GL}_n$, $\mu = (1, 0, \dots, 0)$, and b is basic of slope $1/n$. Let H_0 be the p -divisible group over $\overline{\mathbf{F}}_p$ with isocrystal b , so that H_0 has dimension 1 and height n . In this case $G_b(F) = \text{Aut}^0 H_0 = D^\times$, where D/\mathbf{Q}_p is the division algebra of invariant $1/n$. The spaces $\mathcal{M}_K = \mathcal{M}_{G,b,\mu,K}$ are known as the Lubin-Tate tower; we consider these as rigid-analytic spaces over C , where C/\mathbf{Q}_p is a complete algebraically closed field.

Atop the tower sits the infinite-level Lubin-Tate space $\mathcal{M} = \varprojlim_K \mathcal{M}_K$ as described in [SW13]. This is a perfectoid space admitting an action of $G(\mathbf{Q}_p) \times G_b(\overline{\mathbf{Q}}_p)$. The C -points of \mathcal{M} classify equivalence classes of triples (H, α, ι) , where H/\mathcal{O}_C is a p -divisible group, $\alpha: \mathbf{Q}_p^n \rightarrow VH$ is a trivialization of the rational Tate module, and $\iota: H_0 \otimes_{\overline{\mathbf{F}}_p} \mathcal{O}_C/p \rightarrow H \otimes_{\mathcal{O}_C} \mathcal{O}_C/p$ is an isomorphism in the isogeny category. (Equivalence between two such triples is a quasi-isogeny between p -divisible groups which makes both diagrams commute.) Then \mathcal{M} admits an action of $G(\mathbf{Q}_p) \times G_b(\mathbf{Q}_p)$, via composition with α and ι , respectively.

The Hodge-Tate period map exhibits \mathcal{M} as a pro-étale D^\times -torsor over Drinfeld’s upper half-space Ω^{n-1} (the complement in \mathbf{P}^{n-1} of all \mathbf{Q}_p -rational hyperplanes). This map $\mathcal{M} \rightarrow \Omega^{n-1}$ is equivariant for the action of $G(\mathbf{Q}_p)$.

Now suppose $g \in G(\mathbf{Q}_p)$ is a regular elliptic element (that is, an element with irreducible characteristic polynomial). Then g has exactly n fixed points on Ω^{n-1} . For each such fixed point $x \in (\Omega^{n-1})^g$, the element g acts on the fiber \mathcal{M}_x . Because $\mathcal{M} \rightarrow \Omega^{n-1}$ is a $G_b(F)$ -torsor, there must exist $g' \in G_b(\mathbf{Q}_p)$ such that (g, g') acts trivially on \mathcal{M}_x .

Key observation. The elements $g \in G(\mathbf{Q}_p)$ and $g' \in G_b(\mathbf{Q}_p)$ are related, meaning they become conjugate over $\overline{\mathbf{Q}}_p$.

We sketch the proof of this claim. Suppose y corresponds to the triple (H, α, ι) . This means there exists an automorphism γ of H (in the isogeny category) which corresponds to g on the Tate

module and g' on the special fiber, respectively. We verify now that g and g' are related. Let $B_{\text{cris}} = B_{\text{cris}}(C)$ be the crystalline period ring. There are isomorphisms

$$B_{\text{cris}}^n \rightarrow VH \otimes_{\mathbf{Q}_p} B_{\text{cris}} \rightarrow M(H_0) \otimes B_{\text{cris}},$$

where the first map is induced from α , and the second map comes from the comparison isomorphism between étale and crystalline cohomology of H (using ι to identify the latter with $M(H_0)$). The composite map carries the action of g onto that of g' , which is to say that g and g' become conjugate over B_{cris} . This implies that g and g' are related.

Suppose that ρ is an admissible representation of D^\times with coefficients in $\overline{\mathbf{Q}}_\ell$. There is a corresponding $\overline{\mathbf{Q}}_\ell$ -local system \mathcal{L}_ρ on $\Omega_{C,\text{ét}}^{n-1}$.

Let $g \in G(F)$ be elliptic. A naïve form of the Lefschetz trace formula would predict that:

$$\text{tr}(g|R\Gamma_c(\Omega^{n-1}, \mathcal{L}_\rho)) = \sum_{x \in (\Omega^{n-1})^g} \text{tr}(g|\mathcal{L}_{\rho,x}).$$

For each fixed point x , the key observation above gives $\text{tr}(g|\mathcal{L}_{\rho,x}) = \text{tr} \rho(g')$, where g and g' are related. By the Jacquet-Langlands correspondence, there exists a discrete series representation π of $G(\mathbf{Q}_p)$ satisfying $\text{tr} \pi(g) = (-1)^{n-1} \text{tr} \rho(g')$ (here $\text{tr} \pi(g)$ is interpreted as a Harish-Chandra character). Thus the Euler characteristic of $R\Gamma_c(\Omega^{n-1}, \mathcal{L}_\rho)$ equals $(-1)^{n-1} n\pi$ up to a virtual representation with trace zero on the elliptic locus.

In this situation $\mathcal{S}_\mu = \mathbf{Z}_\ell[n-1]$ (up to a Tate twist), and we find that $R\Gamma(G, b, \mu)[\rho]$ is the shift by $n-1$ of the dual of $R\Gamma_c(\Omega^{n-1}, \mathcal{L}_{\rho^\vee})$. Therefore in $\text{Groth}(\text{GL}_n(\mathbf{Q}_p))$ we have

$$\text{Mant}_{b,\mu}(\rho) = n\pi + \text{err},$$

where the character of err vanishes on the locus of elliptic elements. This is in accord with Theorem 1.0.2.

This argument goes back at least to the 1990s, as discussed in [Har15, Chap. 9], and as far as we know first appears in [Fal94]. The present article is our attempt to push this argument as far as it will go. If a suitable Lefschetz formula is valid, then the equality in Theorem 1.0.2 can be reduced to an endoscopic character identity relating representations of $G(F)$ and $G_b(F)$ (Theorem 3.2.9), which we prove in §3.

Therefore the difficulty in Theorem 1.0.2 lies in proving the validity of the Lefschetz formula. Prior work of Strauch and Mieda proved Theorem 1.0.2 in the case of the Lubin-Tate tower [Str05], [Str08], [Mie12], [Mie14a] and also in the case of a basic Rapoport-Zink space for $\text{GSp}(4)$ [Mie].

In applying a Lefschetz formula to a non-proper rigid space, care must be taken to treat the boundary. For instance, if X is the affinoid unit disc $\{|T| \leq 1\}$ in the adic space \mathbf{A}^1 , then the automorphism $T \mapsto T+1$ has Euler characteristic 1 on X , despite having no fixed points. The culprit is that this automorphism fixes the single boundary point in $\overline{X} \setminus X$. Mieda [Mie14b] proves a Lefschetz formula for an operator on a rigid space, under an assumption that the operator has no topological fixed points on a compactification. Now, in all of the above cases, $\mathcal{M}_{G,b,\mu,K}$ admits a *cellular decomposition*. This means (approximately) that $\mathcal{M}_{G,b,\mu,K}$ contains a compact open subset, whose translates by Hecke operators cover all of $\mathcal{M}_{G,b,\mu,K}$. This is enough to establish the “topological fixed point” hypothesis necessary to apply Mieda’s Lefschetz formula. Shen [She14]

constructs a cellular decomposition for a basic Rapoport-Zink space attached to the group $U(1, n - 1)$, which paves the way for a similar proof of Theorem 1.0.2 in this case as well. For general (G, b, μ) , however, the $\mathcal{M}_{G, b, \mu, K}$ do not admit a cellular decomposition, and so there is probably no hope of applying the methods of [Mie14b].

We had no idea how to proceed, until we learned of the shift of perspective offered by Fargues' program on the geometrization of local Langlands [Farb], followed by the work [FS21]. At the center of that program is the stack Bun_G of G -bundles on the Fargues-Fontaine curve. This is a geometrization of Kottwitz' set $B(G)$: There is a bijection $b \mapsto \mathcal{E}^b$ between $B(G)$ and points of the underlying topological space of Bun_G . For basic b there is an open substack $\text{Bun}_G^b \subset \text{Bun}_G$ classifying G -bundles which are everywhere isomorphic to \mathcal{E}^b ; in this situation $\text{Aut } \mathcal{E}^b = G_b(F)$ and so we have an isomorphism $\text{Bun}_G^b \cong [*/G_b(F)]$.

Let μ be a cocharacter of G . As in geometric Langlands, there is a stack $\text{Hecke}_{G, \leq \mu}$ lying over the product $\text{Bun}_G \times \text{Bun}_G$, which parametrizes μ -bounded modifications of G -bundles at one point of the curve. For each μ , one uses $\text{Hecke}_{G, \leq \mu}$ to define a Hecke operator T_μ on a suitable derived category $D(\text{Bun}_G, \mathbf{Z}_\ell)$ of étale \mathbf{Z}_ℓ -sheaves on Bun_G . If $b \in B(G, \mu)$, then the moduli space of local shtukas $\text{Sht}_{G, b, \mu}$ appears as the fiber of $\text{Hecke}_{G, \leq \mu}$ over the point $(\mathcal{E}^b, \mathcal{E}^1)$ of $\text{Bun}_G \times \text{Bun}_G$. Consequently there is an expression for $R\Gamma(G, b, \mu)[\rho]$ in terms of the Hecke operators T_μ , see Proposition 6.4.5.

Heavy use is made in [FS21] of the notion of *universal local acylity* (ULA) as a property of objects $A \in D(X, \mathbf{Z}_\ell)$ for Artin v-stacks X . When $X = [*/G_b(F)]$, a ULA object is an admissible complex of representations of $G_b(F)$. It is proved in [FS21] that the Hecke operators T_μ preserve ULA objects; the admissibility of $R\Gamma(G, b, \mu)[\rho]$ is deduced from this.

We learned from [LZ20] that the ULA condition is precisely the right hypothesis necessary to prove a Lefschetz-Verdier trace formula applicable to the cohomology of A . This explains the counterexample above: $j_! \mathbf{Z}_\ell$ fails to be ULA, where j is the inclusion of the affinoid disc X into its compactification \overline{X} . In fact [LZ20] is written in the context of schemes, but their formalism applies equally well in the context of rigid-analytic spaces and diamonds. Indeed, some interesting new phenomena occur in the diamond context. For instance, if H is a locally profinite group acting continuously on a proper diamond X , and $A \in D(X, \mathbf{Z}_\ell)$ is a ULA object which is H -equivariant, then $R\Gamma(X, A)$ is an admissible H -module. One gets a formula for the *trace distribution* of H acting on $R\Gamma(X, A)$, in terms of local terms living on the fixed-point locus in $H \times X$. We explain the Lefschetz-Verdier trace formula for diamonds in §4.

In §5, we study the Lefschetz-Verdier trace formula as it pertains to the mixed-characteristic affine Grassmannian $\text{Gr}_{G, \leq \mu}$. The object \mathcal{S}_μ is a ULA on $\text{Gr}_{G, \leq \mu}$ and $G(F)$ -equivariant, so it makes sense to ask for its local term $\text{loc}_g(x, A)$ at a fixed point x of a regular element $g \in G(F)$. (Such fixed points are all isolated.) We found quickly that that result we needed for Theorem 1.0.2 would follow if we knew that $\text{loc}_g(x, A)$ agreed with the naïve local term $\text{tr}(g|A_x)$. We asked Varshavsky, who devised a method for proving this agreement in the scheme setting. We show how to deduce the required statement for $\text{Gr}_{G, \leq \mu}$, using the Witt vector affine Grassmannian as a bridge between diamonds and schemes. (We thank the referee for pointing out that an earlier argument we had here was incorrect.)

Finally, in §6 we prove Theorem 1.0.2 by applying our trace formula to the Hecke stack $\text{Hecke}_{G, b, \leq \mu}$. An important step is to show that fixed points of elliptic elements $g \in G(F)$ act-

ing on $\mathrm{Gr}_{G, \leq \mu}$ are admissible, as we observed above in the Lubin-Tate case.

Acknowledgments

We are grateful to Peter Scholze for explaining to us some of the material on inertia stacks that appears in §4. We also thank Jean-François Dat, Laurent Fargues, Martin Olsson, Jack Thorne, and Yakov Varshavsky for many helpful conversations. Finally, DH is grateful to Marie-France Vignéras for sharing a scanned copy of [Vig96].

2 Review of the objects appearing in Kottwitz's conjecture

2.1 Basic notions

Let \check{F} be the completion of the maximal unramified extension of F , and let $\sigma \in \text{Aut } \check{F}$ be the Frobenius automorphism. Let G be a connected reductive group defined over F . Fix a quasi-split group G^* and a $G^*(\overline{F})$ -conjugacy class Ψ of inner twists $G^* \rightarrow G$. Given an element $b \in G(\check{F})$, there is an associated inner form G_b of a Levi subgroup of G^* as described in [Kot97, §3.3, §3.4]. Its group of F -points is given by

$$G_b(F) \cong \left\{ g \in G(\check{F}) \mid \text{Ad}(b)\sigma(g) = g \right\}.$$

Up to isomorphism the group G_b depends only on the σ -conjugacy class $[b]$. It will be convenient to choose b to be decent [RZ96, Definition 1.8]. Then there exists a finite unramified extension F'/F such that $b \in G(F')$. This allows us to replace \check{F} by F' in the above formula. The slope morphism $\nu : \mathbf{D} \rightarrow G_{\check{F}}$ of b , [Kot85, §4], is also defined over F' . The centralizer $G_{F', \nu}$ of ν in $G_{F'}$ is a Levi subgroup of $G_{F'}$. The $G(F')$ -conjugacy class of ν is defined over F , and then so is the $G(F')$ -conjugacy class of $G_{F', \nu}$. There is a Levi subgroup M^* of G^* defined over F and $\psi \in \Psi$ that restricts to an inner twist $\psi : M^* \rightarrow G_b$, see [Kot97, §4.3].

From now on assume that b is basic. This is equivalent to $M^* = G^*$, so that G_b is in fact an inner form of G^* and of G . Furthermore, Ψ is an equivalence class of inner twists $G^* \rightarrow G$ as well as $G^* \rightarrow G_b$. This identifies the dual groups of G^* , G , and G_b , and we write \widehat{G} for either of them.

Let $\phi : W_F \times \text{SL}_2(\mathbf{C}) \rightarrow {}^L G$ be a discrete Langlands parameter and let $S_\phi = \text{Cent}(\phi, \widehat{G})$. For $\lambda \in X^*(Z(\widehat{G})^\Gamma)$ write $\text{Rep}(S_\phi, \lambda)$ for the set of isomorphism classes of algebraic representations of the algebraic group S_ϕ whose restriction to $Z(\widehat{G})^\Gamma$ is λ -isotypic, and write $\text{Irr}(S_\phi, \lambda)$ for the subset of irreducible such representations. The class of b corresponds to a character $\lambda_b : Z(\widehat{G})^\Gamma \rightarrow \mathbf{C}^\times$ via the isomorphism $B(G)_{\text{bas}} \rightarrow X^*(Z(\widehat{G})^\Gamma)$ of [Kot85, Proposition 5.6]. Assuming the validity of the refined local Langlands conjecture [Kal16a, Conjecture G] we will construct in the following two subsections for any $\pi \in \Pi_\phi(G)$ and $\rho \in \Pi_\phi(G_b)$ an element $\delta_{\pi, \rho} \in \text{Rep}(S_\phi, \lambda_b)$ that measures the relative position of π and ρ .

2.2 Construction of $\delta_{\pi, \rho}$ in a special case

The statements of the Kottwitz conjecture given in [Rap95, Conjecture 5.1] and [RV14, Conjecture 7.3] make the assumption that G is a B -inner form of G^* . In that case, the construction of $\delta_{\pi, \rho}$ is straightforward and we shall now recall it.

The assumption on G means that some $\psi \in \Psi$ can be equipped with a decent basic $b^* \in G^*(F^{\text{nr}})$ such that ψ is an isomorphism $G_{F^{\text{nr}}}^* \rightarrow G_{F^{\text{nr}}}$ satisfying $\psi^{-1}\sigma(\psi) = \text{Ad}(b^*)$. In other words, ψ becomes an isomorphism over F from the group $G_{b^*}^*$ to G . Under this assumption, and after choosing a Whittaker datum \mathfrak{w} for G^* , the isocrystal formulation of the refined local Langlands correspondence [Kal16a, Conjecture F], which is implied by the rigid formulation [Kal16a, Conjecture G] according to [Kal18], predicts the existence of bijections

$$\begin{aligned} \Pi_\phi(G) &\cong \text{Irr}(S_\phi, \lambda_{b^*}) \\ \Pi_\phi(G_b) &\cong \text{Irr}(S_\phi, \lambda_{b^*} + \lambda_b) \end{aligned}$$

where we have used the isomorphisms $B(G)_{\text{bas}} \cong X^*(Z(\widehat{G})^\Gamma) \cong B(G^*)_{\text{bas}}$ of [Kot85, Proposition 5.6] to obtain from $[b] \in B(G)_{\text{bas}}$ and $[b^*] \in B(G^*)_{\text{bas}}$ characters λ_b and λ_{b^*} of $Z(\widehat{G})^\Gamma$.

These bijections are uniquely characterized by the endoscopic character identities which are part of [Kal16a, Conjecture F]. Write $\pi \mapsto \tau_{b^*, \mathfrak{w}, \pi}$, $\rho \mapsto \tau_{b^*, \mathfrak{w}, \rho}$ for these bijections and define

$$\delta_{\pi, \rho} := \check{\tau}_{b^*, \mathfrak{w}, \pi} \otimes \tau_{b^*, \mathfrak{w}, \rho}. \quad (2.2.1)$$

While these bijections depend on the choice of Whittaker datum \mathfrak{w} and the choice of b^* , we will argue in Subsection 2.3 that for any pair π and ρ the representation $\delta_{\pi, \rho}$ is independent of these choices. Of course it does depend on b , but this we take as part of the given data.

2.3 Construction of $\delta_{\pi, \rho}$ in the general case

We now drop the assumption that G is a B -inner form of G^* . Because of this, we no longer have the isocrystal formulation of the refined local Langlands correspondence. However, we do have the formulation based on rigid inner twists [Kal16a, Conjecture G]. What this means with regards to the Kottwitz conjecture is that neither π nor ρ correspond to representations of S_ϕ . Rather, they correspond to representations τ_π and τ_ρ of a different group $\pi_0(S_\phi^+)$. Nonetheless it will turn out that $\check{\tau}_\pi \otimes \tau_\rho$ provides in a natural way a representation $\delta_{\pi, \rho}$ of S_ϕ .

In order to make this precise we will need the material of [Kal16b] and [Kal18], some of which is summarized in [Kal16a]. First, we will need the cohomology set $H^1(u \rightarrow W, Z \rightarrow G^*)$ defined in [Kal16b, §3] for any finite central subgroup $Z \subset G^*$ defined over F . As in [Kal18, §3.2] it will be convenient to package these sets for varying Z into the single set

$$H^1(u \rightarrow W, Z(G^*) \rightarrow G^*) := \varinjlim H^1(u \rightarrow W, Z \rightarrow G^*).$$

The transition maps on the right are injective, so the colimit can be seen as an increasing union.

Next, we will need the reinterpretation, given in [Kot], of $B(G)$ as the set of cohomology classes of algebraic 1-cocycles of a certain Galois gerbe $1 \rightarrow \mathbf{D}(\bar{F}) \rightarrow \mathcal{E} \rightarrow \Gamma \rightarrow 1$. This reinterpretation is also reviewed in [Kal18, §3.1]. Let us make it explicit for basic decent elements $b' \in G(F^{\text{nr}})$. There is a uniquely determined 1-cocycle $\mathbf{Z} \cong \langle \sigma \rangle \rightarrow G(F^{\text{nr}})$ whose value at σ is equal to b' . By inflation we obtain a 1-cocycle of W_F in $G(\bar{F})$. Since b' is basic and decent, for some finite Galois extension K/F splitting G the restriction of this 1-cocycle to W_K factors through K^\times and is a homomorphism $K^\times \rightarrow Z(G)(K)$. Moreover, this homomorphism is algebraic and is in fact given by a multiple of the slope morphism $\nu : \mathbf{D} \rightarrow G$. In this way we obtain a 1-cocycle valued in $G(\bar{F})$ of the extension $1 \rightarrow K^\times \rightarrow W_{K/F} \rightarrow \Gamma_{K/F} \rightarrow 1$, which we can pull-back along $\Gamma \rightarrow \Gamma_{K/F}$ and then combine with ν to obtain a 1-cocycle of \mathcal{E} valued in $G(\bar{F})$, that is algebraic in the sense that its restriction to $\mathbf{D}(\bar{F})$ is given by a morphism of algebraic groups, namely ν . The reader is referred to [Kot97, §8 and App B] for further details.

Finally, we will need the comparison map

$$B(G)_{\text{bas}} \rightarrow H^1(u \rightarrow W, Z(G) \rightarrow G)$$

of [Kal18, §3.3]. In fact, this comparison map is already defined on the level of cocycles, via pull-back along the diagram [Kal18, (3.13)], and takes the form

$$G(F^{\text{nr}})_{\text{d, bas}} \rightarrow Z^1(u \rightarrow W, Z(G) \rightarrow G) \quad (2.3.1)$$

where on the left we have the decent basic elements in $G(F^{\text{nr}})$.

After this short review we turn to the construction of $\delta_{\pi, \rho} \in \text{Rep}(S_\phi, \lambda_b)$. Choose any inner twist $\psi \in \Psi$ and let $\bar{z}_\sigma := \psi^{-1}\sigma(\psi) \in G_{\text{ad}}^*(\bar{F})$. Then $\bar{z} \in Z^1(F, G_{\text{ad}}^*)$ and the surjectivity of the natural map $H^1(u \rightarrow W, Z(G^*) \rightarrow G^*) \rightarrow H^1(F, G_{\text{ad}}^*)$ asserted in [Kal16b, Corollary 3.8] allows us to choose $z \in Z^1(u \rightarrow W, Z(G^*) \rightarrow G^*)$ lifting \bar{z} . Then $(\psi, z) : G^* \rightarrow G$ is a rigid inner twist. Let $z_b \in Z^1(u \rightarrow W, Z(G) \rightarrow G)$ be the image of b under (2.3.1). For psychological reasons, let $\xi : G_{F'} \rightarrow G_{b, F'}$ denote the identity map. Then $(\xi \circ \psi, \psi^{-1}(z) \cdot z_b) : G^* \rightarrow G_b$ is also a rigid inner twist.

The L -packets $\Pi_\phi(G)$ and $\Pi_\phi(G_b)$ are now parameterized by representations of a certain cover S_ϕ^+ of S_ϕ . While [Kal16a, Conjecture G] is formulated in terms of a finite cover depending on an auxiliary choice of a finite central subgroup $Z \subset G^*$, we will adopt here the point of view of [Kal18] and work with a canonical infinite cover, namely the preimage of S_ϕ in the universal cover of \widehat{G} . Following [Kal18, §3.3] we can present this universal cover as follows. Let $Z_n \subset Z(G)$ be the subgroup of those elements whose image in $Z(G)/Z(G_{\text{der}})$ is n -torsion, and let $G_n = G/Z_n$. Then G_n has adjoint derived subgroup and connected center. More precisely, $G_n = G_{\text{ad}} \times C_n$, where $C_n = C_1/C_1[n]$ and $C_1 = Z(G)/Z(G_{\text{der}})$. It is convenient to identify $C_n = C_1$ as algebraic tori and take the m/n -power map $C_1 \rightarrow C_1$ as the transition map $C_n \rightarrow C_m$ for $n|m$. The isogeny $G \rightarrow G_n$ dualizes to $\widehat{G}_n \rightarrow \widehat{G}$ and we have $\widehat{G}_n = \widehat{G}_{\text{sc}} \times \widehat{C}_1$. Note that $\widehat{C}_1 = Z(\widehat{G})^\circ$. The transition map $\widehat{G}_m \rightarrow \widehat{G}_n$ is then the identity on \widehat{G}_{sc} and the m/n -power map on \widehat{C}_1 . Set $\widehat{G} = \varprojlim \widehat{G}_n = \widehat{G}_{\text{sc}} \times \widehat{C}_\infty$, where $\widehat{C}_\infty = \varprojlim \widehat{C}_n$. Then \widehat{G} is the universal cover of \widehat{G} . Elements of \widehat{G} can be written as $(a, (b_n)_n)$, where $a \in \widehat{G}_{\text{sc}}$ and $(b_n)_n$ is a sequence of elements $b_n \in \widehat{C}_1$ satisfying $b_n = (b_m)^{\frac{m}{n}}$ for $n|m$. In this presentation, the natural map $\widehat{G} \rightarrow \widehat{G}$ sends $(a, (b_n)_n)$ to $a_{\text{der}} \cdot b_1$, where $a_{\text{der}} \in \widehat{G}_{\text{der}}$ is the image of $a \in \widehat{G}_{\text{sc}}$ under the natural map $\widehat{G}_{\text{sc}} \rightarrow \widehat{G}_{\text{der}}$.

Definition 2.3.1. Let $Z(\widehat{G})^+ \subset S_\phi^+ \subset \widehat{G}$ be the preimages of $Z(\widehat{G})^\Gamma \subset S_\phi \subset \widehat{G}$ under $\widehat{G} \rightarrow \widehat{G}$.

Given a character $\lambda : \pi_0(Z(\widehat{G})^+) \rightarrow \mathbf{C}^\times$ (which we will always assume trivial on the kernel of $Z(\widehat{G})^+ \rightarrow \widehat{G}_n$ for some n) let $\text{Rep}(\pi_0(S_\phi^+), \lambda)$ denote the set of isomorphism classes of representations of $\pi_0(S_\phi^+)$ whose pull-back to $\pi_0(Z(\widehat{G})^+)$ is λ -isotypic, and let $\text{Irr}(\pi_0(S_\phi^+), \lambda)$ be the (finite) subset of irreducible representations. Let λ_z be the character corresponding to the class of z under the Tate-Nakayama isomorphism

$$H^1(u \rightarrow W, Z(G^*) \rightarrow G^*) \rightarrow \pi_0(Z(\widehat{G})^+)^*$$

of [Kal16b, Corollary 5.4], and let λ_{z_b} be the character corresponding to the class of z_b in $H^1(u \rightarrow W, Z(G) \rightarrow G)$. Then according to [Kal16a, Conjecture G], upon fixing a Whittaker datum \mathfrak{w} for G^* there are bijections

$$\begin{aligned} \Pi_\phi(G) &\cong \text{Irr}(\pi_0(S_\phi^+), \lambda_z) \\ \Pi_\phi(G_b) &\cong \text{Irr}(\pi_0(S_\phi^+), \lambda_z + \lambda_{z_b}) \end{aligned}$$

again uniquely determined by the endoscopic character identities. We write $\pi \mapsto \tau_{z, \mathfrak{w}, \pi}$, $\rho \mapsto \tau_{z, \mathfrak{w}, \rho}$ for these bijections, and $\tau \mapsto \pi_{z, \mathfrak{w}, \tau}$, $\tau \mapsto \rho_{z, \mathfrak{w}, \tau}$ for their inverses. We form the representation $\check{\tau}_{z, \mathfrak{w}, \pi} \otimes \tau_{z, \mathfrak{w}, \rho} \in \text{Rep}(\pi_0(S_\phi^+), \lambda_{z_b})$.

Recall the map [Kal18, (4.7)]

$$S_\phi^+ \rightarrow S_\phi, \quad (a, (b_n)) \mapsto \frac{a_{\text{der}} \cdot b_1}{N_{E/F}(b_{[E:F]})}. \quad (2.3.2)$$

Here $a_{\text{der}} \in \widehat{G}_{\text{der}}$ is the image of $a \in \widehat{G}_{\text{sc}}$ under the natural map $\widehat{G}_{\text{sc}} \rightarrow \widehat{G}_{\text{der}}$ and E/F is a sufficiently large finite Galois extension. This map is independent of the choice of E/F . According to [Kal18, Lemma 4.1] pulling back along this map sets up a bijection $\text{Irr}(\pi_0(S_\phi^+), \lambda_{z_b}) \rightarrow \text{Irr}(S_\phi, \lambda_b)$. Note that since ϕ is discrete the group S_ϕ^\natural defined in loc. cit. is equal to S_ϕ . The lemma remains valid, with the same proof, if we remove the requirement of the representations being irreducible, and we obtain the bijection $\text{Rep}(\pi_0(S_\phi^+), \lambda_{z_b}) \rightarrow \text{Rep}(S_\phi, \lambda_b)$.

Definition 2.3.2. Let $\delta_{\pi, \rho}$ be the image of $\check{\tau}_{z, \mathfrak{w}, \pi} \otimes \tau_{z, \mathfrak{w}, \rho}$ under the bijection $\text{Rep}(\pi_0(S_\phi^+), \lambda_{z_b}) \rightarrow \text{Rep}(S_\phi, \lambda_b)$.

In the situation when G is a B -inner form of G^* , this definition of $\delta_{\pi, \rho}$ agrees with the one of Subsection 2.2, because then we can take z to be the image of b^* under (2.3.1) and then $\tau_{z, \mathfrak{w}, \pi}$ and $\tau_{b^*, \mathfrak{w}, \pi}$ are related via (2.3.2), and so are $\tau_{z, \mathfrak{w}, \rho}$ and $\tau_{b^*, \mathfrak{w}, \rho}$, see [Kal18, §4.2].

Lemma 2.3.3. *The representation $\delta_{\pi, \rho}$ is independent of the choices of Whittaker datum \mathfrak{w} and of a rigidifying 1-cocycle $z \in Z^1(u \rightarrow W, Z(G^*) \rightarrow G^*)$.*

Proof. Both of these statements follow from [Kal16a, Conjecture G]. For the independence of Whittaker datum, one can prove that the validity of this conjecture implies that if \mathfrak{w} is replaced by another choice \mathfrak{w}' then there is an explicitly constructed character $(\mathfrak{w}, \mathfrak{w}')$ of $\pi_0(S_\phi/Z(\widehat{G})^\Gamma)$ whose inflation to $\pi_0(S_\phi^+)$ satisfies $\tau_{z, \mathfrak{w}, \sigma} = \tau_{z, \mathfrak{w}', \sigma} \otimes (\mathfrak{w}, \mathfrak{w}')$ for any $\sigma \in \Pi_\phi(G) \cup \Pi_\phi(G_b)$. See §4 and in particular Theorem 4.3 of [Kal13], the proof of which is valid for a general G that satisfies [Kal16a, Conjecture G], bearing in mind that the transfer factor we use here is related to the one used there by $s \mapsto s^{-1}$. The independence of z follows from the same type of argument, but now using [Kal18, Lemma 6.2]. \square

2.4 Spaces of local shtukas and their cohomology

We recall here some material from [SW20] and [Farb] regarding the Fargues-Fontaine curve and moduli spaces of local shtukas.

For any perfectoid space S of characteristic p , we have the Fargues-Fontaine curve X_S [FF], an adic space over \mathbf{Q}_p . If S lies over k , there is a relative version of this curve lying over F . Since F is fixed we use the same notation X_S for this. For $S = \text{Spa}(R, R^+)$ affinoid with pseudouniformizer ϖ , the adic space X_S is defined as follows:

$$\begin{aligned} Y_S &= (\text{Spa } W_{\mathcal{O}_F}(R^+)) \setminus \{p[\varpi] = 0\} \\ X_S &= Y_S / \text{Frob}^{\mathbf{Z}}. \end{aligned}$$

Here Frob is the q th power Frobenius on S .

For an affinoid perfectoid space S lying over the residue field of F , the following sets are in bijection:

1. S -points of $\mathrm{Spd} F$
2. Untilts S^\sharp of S over F ,
3. Cartier divisors of Y_S of degree 1.

Given an untilt S^\sharp , we let $D_{S^\sharp} \subset Y_S$ be the corresponding divisor. If $S^\sharp = \mathrm{Spa}(R^\sharp, R^{\sharp+})$ is affinoid, then the completion of Y_S along D_{S^\sharp} is represented by Fontaine's ring $B_{\mathrm{dR}}^+(R^\sharp)$. The untilt S^\sharp determines a Cartier divisor on X_S , which we still refer to as D_{S^\sharp} .

There is a functor $b \mapsto \mathcal{E}^b$ from the category of isocrystals with G -structure to the category of G -bundles on X_S (for any S). When S is a geometric point this functor induces an isomorphism between the sets of isomorphism classes [Fara].

We now recall Scholze's definition of the local shtuka space. It is a set-valued functor on the pro-étale site of perfectoid spaces over \mathbf{F}_p and is equipped with a morphism to $\mathrm{Spd} C$. Thus it can be described equivalently as a set-valued functor on the pro-étale site of perfectoid spaces over C .

Definition 2.4.1. The local shtuka space $\mathrm{Sht}_{G,b,\mu}$ inputs a perfectoid C -algebra (R, R^+) , and outputs the set of isomorphisms

$$\gamma: \mathcal{E}^1|_{X_{R^b} \setminus D_R} \cong \mathcal{E}^b|_{X_{R^b} \setminus D_R}$$

of G -torsors that are meromorphic along D_R and bounded by μ pointwise on $\mathrm{Spa} R$.

Let us briefly recall the condition of being pointwise bounded by μ . If $\mathrm{Spa}(C, O_C) \rightarrow \mathrm{Spa} R$ is a geometric point we obtain via pull-back $\gamma: \mathcal{E}^1|_{X_{C^b} \setminus \{x_C\}} \rightarrow \mathcal{E}^b|_{X_{C^b} \setminus \{x_C\}}$, where we have written x_C in place of D_C to emphasize that this is a point on X_{C^b} . The completed local ring of X_{C^b} at x_C is Fontaine's ring $B_{\mathrm{dR}}^+(C)$. A trivialization of both bundles \mathcal{E}^1 and \mathcal{E}^b on a formal neighborhood of x_C , together with γ , leads to an element of $G(B_{\mathrm{dR}}(C))$, well defined up to left and right multiplication by elements of $G(B_{\mathrm{dR}}^+(C))$. The corresponding element of the double coset space $G(B_{\mathrm{dR}}^+(C)) \backslash G(B_{\mathrm{dR}}(C)) / G(B_{\mathrm{dR}}^+(C))$ is indexed by a conjugacy class of co-characters of G/C according to the Cartan decomposition, and we demand that this conjugacy class is dominated by μ in the usual order (given by the simple roots of the universal Borel pair).

The space $\mathrm{Sht}_{G,b,\mu}$ is a locally spatial diamond [SW20, §23]. Since the automorphism groups of \mathcal{E}^1 and \mathcal{E}^b are the constant group diamonds $G(\mathbf{Q}_p)$ and $G_b(\mathbf{Q}_p)$ respectively, the space $\mathrm{Sht}_{G,b,\mu}$ is equipped with commuting actions of $G(\mathbf{Q}_p)$ and $G_b(\mathbf{Q}_p)$, acting by pre- and post-composition on γ .

Remark 2.4.2. According to [SW20, Corollary 23.2.2] the above definition recovers the moduli space of local shtukas with one leg and infinite level structure. We have dropped the subscript ∞ used in [SW20] to denote the infinite level structure.

We will use the cohomology theory developed in [Sch17]. For any compact open subgroup $K \subset G(F)$ the quotient $\mathrm{Sht}_{G,b,\mu,K} = \mathrm{Sht}_{G,b,\mu} / K$ is again a locally spatial diamond [SW20, §23]. For each $n = 1, 2, \dots$, let $V_{\mu,n} \in \mathrm{Rep}(\widehat{G}, \mathbf{Z}/\ell^n \mathbf{Z})$ be the Weyl module associated to μ . By the Satake equivalence (Theorem 5.1.1), there is a corresponding object $\mathcal{S}_{\mu,n}$ of $D_{\mathrm{ét}}(\mathrm{Gr}_{G,b,\leq \mu}, \mathbf{Z}/\ell^n \mathbf{Z})$. Define

$$R\Gamma_c(\mathrm{Sht}_{G,b,\mu} / K, \mathcal{S}_\mu) = \varinjlim_U R\Gamma_c(U, \mathcal{S}_\mu),$$

where $U \subset \text{Sht}_{G,b,\mu}/K$ runs over quasicompact open subsets, and where we have put

$$R\Gamma_c(U, \mathcal{S}_\mu) = \varprojlim_n R\Gamma_c(U, \mathcal{S}_{\mu,n}).$$

Then $R\Gamma_c(\text{Sht}_{G,b,\mu}/K, \mathcal{S}_\mu)$ is a complex of \mathbf{Z}_ℓ -modules carrying an action of $G_b(F) \times W_E$.

Definition 2.4.3. Let ρ be an irreducible admissible representation of $G_b(F)$ with coefficients in $\overline{\mathbf{Q}}_\ell$. Then we define

$$R\Gamma(G, b, \mu)[\rho] = \varinjlim_{K \subset G(F)} R\text{Hom}_{G_b(F)}(R\Gamma_c(\text{Sht}_{G,b,\mu}/K, \mathcal{S}_\mu) \otimes \overline{\mathbf{Q}}_\ell, \rho),$$

where K runs over the set of open compact subgroups of $G(F)$.

This defines a finite-length W_E -equivariant object in the derived category of smooth representations of $G(F)$ with coefficients in $\overline{\mathbf{Q}}_\ell$, and we write $\text{Mant}_{b,\mu}(\rho)$ for the image of $R\Gamma(G, b, \mu)[\rho]$ in $\text{Groth}(G(F) \times W_E)$.

Remark 2.4.4. We now discuss the relationship between our definition of $\text{Mant}_{b,\mu}(\rho)$ and the virtual representation $H^*(G, b, \mu)[\rho]$ defined in [RV14].

When μ is minuscule, $\text{Sht}_{G,b,\mu,K}$ is the diamond $\mathcal{M}_{G,b,\mu,K}^\circ$ associated to the local Shimura variety $\mathcal{M}_{G,b,\mu,K}$ [SW20, §24.1]. Moreover, in that case, $\mathcal{S}_\mu = \mathbf{Z}_\ell[d](\frac{d}{2})$ is a shift and twist of the constant sheaf. In [RV14], $H^*(G, b, \mu)[\rho]$ is defined as the alternating sum

$$\sum_{i,j \in \mathbf{Z}} (-1)^{i+j} H^{i,j}(G, b, \mu)[\rho](-d),$$

where

$$H^{i,j}(G, b, \mu)[\rho] = \varinjlim_K \text{Ext}_{G_b(F)}^i(R^j\Gamma_c(\mathcal{M}_{G,b,\mu,K}, \mathbf{Z}_\ell) \otimes \overline{\mathbf{Q}}_\ell, \rho)$$

Note that $H^{i,j}(G, b, \mu)[\rho]$ vanishes for all but finitely many (i, j) and each $H^{i,j}(G, b, \mu)[\rho]$ is an admissible representation of $G_b(F)$, by the analysis in [FS21]. On the other hand, unwinding definitions, we see that there is a spectral sequence $H^{i,j-d}(G, b, \mu)[\rho](-\frac{d}{2}) \implies H^{i+j}(R\Gamma(G, b, \mu)[\rho])$.

Putting these observations together, we get the equality

$$\text{Mant}_{b,\mu}(\rho) = (-1)^d H^*(G, b, \mu)[\rho](\frac{d}{2}).$$

Note that in our formulation, the Tate twist appearing in [RV14, Conjecture 7.3] has been absorbed into the normalization of $\text{Mant}_{b,\mu}$.

3 Transfer of conjugation-invariant functions from $G(F)$ to $G_b(F)$

Throughout, F/\mathbf{Q}_p is a finite extension, and G/F is a connected reductive group.

3.1 The space of strongly regular conjugacy classes in $G(F)$

The following definitions are important for our work.

- $G_{\text{rs}} \subset G$ is the open subvariety of regular semisimple elements, meaning those whose connected centralizer is a maximal torus.
- $G_{\text{sr}} \subset G$ is the open subvariety of strongly regular semi-simple elements, meaning those regular semisimple elements whose centralizer is connected, i.e. a maximal torus;
- $G(F)_{\text{ell}} \subset G(F)$ is the open subset of strongly regular elliptic elements, meaning those strongly regular semisimple elements in $G(F)$ whose centralizer is an elliptic maximal torus.

We put $G(F)_{\text{sr}} = G_{\text{sr}}(F)$ and $G(F)_{\text{rs}} = G_{\text{rs}}(F)$. Note that $G(F)_{\text{ell}} \subset G(F)_{\text{sr}} \subset G(F)_{\text{rs}}$. The inclusion $G(F)_{\text{sr}} \subset G(F)_{\text{rs}}$ is dense.

If g is regular semisimple, then it is necessarily contained in a unique maximal torus T , namely the neutral component $\text{Cent}(g, G)^\circ$, but this is not necessarily all of $\text{Cent}(g, G)$. If G is simply connected, then $\text{Cent}(g, G)$ is connected; thus in such a group, regular semisimple and strongly regular semisimple mean the same thing.

Observe that if g is regular semisimple, then $\alpha(g) \neq 1$ for all roots α relative to the action of T . Indeed, if $\alpha(g) = 1$, then the root subgroup of α would commute with g , and then it would have dimension strictly greater than $\dim T$.

All of the sets $G(F)_{\text{sr}}$, $G(F)_{\text{rs}}$, $G(F)_{\text{ell}}$ are conjugacy-invariant, so we may for instance consider the quotient $G(F)_{\text{sr}} // G(F)$, considered as a topological space.

Lemma 3.1.1. *$G(F)_{\text{sr}} // G(F)$ is locally profinite.*

Proof. Let $T \subset G$ be a F -rational maximal torus, and let $N(T, G)$ be its centralizer in G . The set $H^1(F, N(T, G))$ classifies conjugacy classes of F -rational tori, as follows: given a F -rational torus T' , we must have $T = xT'x^{-1}$ for some $x \in G(\bar{F})$. Then for all $\sigma \in \text{Gal}(\bar{F}/F)$, $x^\sigma x^{-1}$ normalizes T . We associate to T' the class of $\sigma \mapsto x^\sigma x^{-1}$ in $H^1(F, N(T, G))$, and it is a simple matter to see that this defines a bijection as claimed. (In fact $H^1(F, N(T, G))$ is finite.)

There is a map $G(F)_{\text{sr}} // G(F) \rightarrow H^1(F, N(T, G))$, sending the conjugacy class of $g \in G(F)_{\text{sr}}$ to the conjugacy class of the unique F -rational torus containing it, namely $\text{Cent}(g, G)^\circ$. We claim that this map is locally constant.

To prove the claim, we consider

$$\varphi : G(F) \times T_{\text{sr}}(F) \rightarrow G_{\text{sr}}(F), \quad (g, t) \mapsto gtg^{-1},$$

a morphism of p -adic analytic varieties. We would like to show that φ is open. To do this, we will compute its differential at the point (g, t) , by means of a change of variable. Consider the map

$$\psi = L_{gtg^{-1}}^{-1} \circ \varphi \circ (L_g \times L_t).$$

Explicitly, for $(z, w) \in G(F) \times T(F)$ we have $\psi(z, w) = gt^{-1}ztwz^{-1}g^{-1}$.

Let $\mathfrak{g} = \text{Lie } G$, $\mathfrak{t} = \text{Lie } T$. The derivative $d\psi(1, 1) : \mathfrak{g} \times \mathfrak{t} \rightarrow \mathfrak{g}$ is given by the formula

$$d\psi(1, 1)(Z, W) = \text{Ad}(g)[(\text{Ad}(t^{-1}) - \text{id})Z + W].$$

We would like to check that $d\psi(1, 1)$ is surjective. We may decompose $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{t}^\perp$, where \mathfrak{t}^\perp is the descent to F of the direct sum of all root subspaces of $\mathfrak{g}_{\overline{F}}$ for the action of T .

The element t is strongly regular. We have observed that $\alpha(t) \neq 1$ for all roots of \mathfrak{g} for the action of T . Therefore $\text{Ad}(t^{-1}) - \text{id} : \mathfrak{g}/\mathfrak{t} \rightarrow \mathfrak{t}^\perp$ is an isomorphism. It follows that $d\psi$ is surjective. The derivative of φ at (g, t) is

$$d\varphi(g, t) = dL_{gtg^{-1}}(gtg^{-1}) \circ d\psi(1, 1) \circ (dL_g(1) \times dL_t(1)).$$

All terms dL are isomorphisms, so $d\varphi(g, t)$ is also surjective. Thus φ is a submersion in the sense of Bourbaki VAR §5.9.1, hence it is open by loc. cit. §5.9.4.

Therefore if $g \in T(F)_{\text{sr}}$ and g' is sufficiently close to g in $G(F)$, then g' is conjugate in $G(F)$ to an element of $T(F)$, which proves the claim about the local constancy of $G(F)_{\text{sr}} // G(F) \rightarrow H^1(F, N(T, G))$.

The fiber of this map over T' is $T'(F)_{\text{sr}}$ modulo the action of the finite group $N(T', G)(F)/T'(F)$. This group acts freely on $T'(F)_{\text{sr}}$, since by definition $\text{Cent}(g, G) = T'$ for $g \in T'(F)_{\text{sr}}$. Since $T'(F)$ is locally profinite, so is its quotient by the free action of a finite group. \square

3.2 Hecke correspondences

Suppose that $b \in G(\check{F})$ is basic. The goal of this section is to define a family of explicit Hecke operators, which input a conjugation-invariant function on $G(F)_{\text{rs}}$ and output a conjugation-invariant function on $G_b(F)_{\text{rs}}$. Given a sufficiently strong version of the local Langlands conjectures, we will show that the Hecke operators act predictably on the trace characters attached to irreducible admissible representations.

We begin by recalling the concept of related elements and the definition of their invariant in the isocrystal setting from [Kal14].

Lemma 3.2.1. *Suppose $g \in G(F)$ and $g' \in G_b(F)$ are strongly regular elements which are conjugate over an algebraic closure of \check{F} . Then they are conjugate over \check{F} .*

Proof. Let K be an algebraic closure of \check{F} . Say $g' = zgz^{-1}$ with $z \in G(K)$. Let $T = \text{Cent}(g, G)$; then for all τ in the inertia group $\text{Gal}(\overline{F}/F^{\text{nr}})$, $z^{-\tau}z$ commutes with g and therefore lies in $T(K)$. Then $\tau \mapsto z^{-\tau}z$ is a cocycle in $H^1(\check{F}, T)$. Since T is a connected algebraic group, $H^1(\check{F}, T) = 0$ [Ste65, Theorem 1.9]. If $x \in T(K)$ splits the cocycle, then $y = zx^{-1} \in G(\check{F})$, and $g' = ygy^{-1}$, so that g and g' are related. \square

It is customary to call elements g, g' as in the above lemma *stably conjugate*, or *related*. Suppose we have strongly regular elements $g \in G(F)_{\text{rs}}$ and $g' \in G_b(F)_{\text{rs}}$ which are related. Let $T = \text{Cent}(g, G)$, and suppose $y \in G(\check{F})$ with $g' = ygy^{-1}$. The rationality of g means that $g^\sigma = g$, whereas the rationality of g' in G_b means that $(g')^\sigma = b^{-1}g'/b$. Combining these statements shows that $b_0 := y^{-1}by^\sigma$ commutes with g and therefore lies in $T(\check{F})$.

Definition 3.2.2. For strongly regular related elements $g \in G(F)_{\text{rs}}$ and $g' \in G_b(F)_{\text{rs}}$, the invariant $\text{inv}[b](g, g')$ is the class of $y^{-1}by^\sigma$ in $B(T)$, where $y \in G(F)$ satisfies $g' = ygy^{-1}$.

Fact 3.2.3. The invariant $\text{inv}[b](g, g') \in B(T)$ only depends on b , g , and g' and not on the element y which conjugates g into g' . It depends on the rational conjugacy classes of g and g' as follows:

- For $z \in G(F)$ we have $\text{inv}[b](\text{ad } z)(g), g') = (\text{ad } z)(\text{inv}[b](g, g'))$, a class in $B(\text{ad } z)(T)$.
- For $z \in G_b(F)$ we have $\text{inv}[b](g, \text{ad } z)(g') = \text{inv}[b](g, g')$.

The image of $\text{inv}[b](g, g')$ under the composition of $B(T) \rightarrow B(G)$ and $\kappa: B(G) \rightarrow \pi_1(G)_\Gamma$ equals $\kappa(b)$.

Definition 3.2.4. We define a diagram of topological spaces

$$\begin{array}{ccc} & \text{Rel}_b & \\ & \swarrow \quad \searrow & \\ G(F)_{\text{sr}} // G(F) & & G_b(F)_{\text{sr}} // G_b(F). \end{array} \tag{3.2.1}$$

as follows. The space Rel_b is the set of conjugacy classes of triples (g, g', λ) , where $g \in G(F)_{\text{sr}}$ and $g' \in G_b(F)_{\text{sr}}$ are related, and $\lambda \in X_*(T)$, where $T = \text{Cent}(g, G)$. It is required that $\kappa(\text{inv}[b](g, g'))$ agrees with the image of λ in $X_*(T)_\Gamma$. We consider (g, g', λ) conjugate to $((\text{ad } z)(g), (\text{ad } z')(g'), (\text{ad } z)(\lambda))$ whenever $z \in G(F)$ and $z' \in G_b(F)$. We give $\text{Rel}_b \subset (G(F) \times G_b(F) \times X_*(G))/(G(F) \times G_b(F))$ the subspace topology, where $X_*(G)$ is taken to be discrete.

Remark 3.2.5. Given $g \in G(F)_{\text{sr}}$ and λ a cocharacter of its torus, there is at most one conjugacy class of $g' \in G_b(F)$ with $(g, g', \lambda) \in \text{Rel}_b$. In other words, g and $\text{inv}[b](g, g')$ determine the conjugacy class of g' . Indeed, suppose (g, g', λ) and (g, g'', λ) are both in Rel_b . Then $g' = ygy^{-1}$ and $g'' = zgz^{-1}$ for some $y, z \in G(F^{\text{nr}})$, and $y^{-1}by^\sigma$ and $z^{-1}bz^\sigma$ are σ -conjugate in $T(F)$. This means there exists $t \in T(F)$ such that $y^{-1}by^\sigma = (zt)^{-1}b(zt)^\sigma$. We see that $x = zty^{-1} \in G_b(F)$, and that x conjugates g' onto g'' .

Lemma 3.2.6. The map $\text{Rel}_b \rightarrow G(F)_{\text{sr}} // G(F)$ is a homeomorphism locally on the source. Its image consists of those classes that transfer to G_b . In particular, the image is open and closed.

The analogous statement is true for $\text{Rel}_b \rightarrow G_b(F)_{\text{sr}} // G_b(F)$.

Proof. The proof of Lemma 3.1.1 shows that $G(F)_{\text{sr}} // G(F)$ is the disjoint union of spaces $T(F)_{\text{sr}}/W_T$, as $T \subset G$ runs through the finitely many conjugacy classes of F -rational maximal tori, and $W_T = N(T, G)(F)/T(F)$ is a finite group. By the above remark, Rel_b injects into the disjoint union of the spaces $T(F)_{\text{sr}}/W_T \times X_*(T)$, with the map to $G(F)_{\text{sr}} // G(F)$ corresponding to the projection $T(F)_{\text{sr}}/W_T \times X_*(T) \rightarrow T(F)_{\text{sr}}/W_T$. Since $X_*(T)$ is discrete, this map is a homeomorphism locally on the source. The other statements are evident from the definitions. \square

The definition of Rel_b already suggests a means for transferring functions from $G(F)_{\text{sr}} // G(F)$ to $G_b(F)_{\text{sr}} // G_b(F)$, namely, by pulling back from $G(F)_{\text{sr}} // G(F)$ to Rel_b , multiplying by a compactly supported kernel function, and then pushing forward to $G_b(F)_{\text{sr}} // G_b(F)$. We will define one such kernel function for each conjugacy class of \bar{F} -rational cocharacters $\mu: \mathbf{G}_m \rightarrow G$.

Let \widehat{G} be the Langlands dual group. Given a cocharacter μ , we obtain a triple $(\widehat{T}, \widehat{B}, \widehat{\mu})$ up to \widehat{G} -conjugacy, where $\widehat{T} \subset \widehat{G}$ is a maximal torus, $\widehat{B} \subset \widehat{G}$ is a Borel subgroup containing \widehat{T} , and $\widehat{\mu} : \widehat{T} \rightarrow \mathbf{G}_m$ is a character that is \widehat{B} -dominant. Let r_μ be the unique irreducible rational representation of the dual group \widehat{G} whose highest weight with respect to $(\widehat{T}, \widehat{B})$ is $\widehat{\mu}$.

A cocharacter $\lambda \in X_*(T)$ corresponds to a character $\widehat{\lambda} \in X^*(\widehat{T})$. Let $r_\mu[\lambda]$ be the $\widehat{\lambda}$ -weight space of r_μ . The quantity $\dim r_\mu[\lambda]$ will give us our kernel function.

We now fix a commutative ring Λ in which p is invertible. For a topological space X , we let $C(X, \Lambda)$ be the space of continuous Λ -valued functions on X , where Λ is given the discrete topology.

Definition 3.2.7. We define the Hecke operator

$$T_{b,\mu}^{G \rightarrow G_b} : C(G(F)_{\text{sr}} // G(F), \Lambda) \rightarrow C(G_b(F)_{\text{sr}} // G_b(F), \Lambda)$$

by

$$[T_{b,\mu}^{G \rightarrow G_b} f](g') = \sum_{(g,g',\lambda) \in \text{Rel}_b} f(g) \dim r_\mu[\lambda].$$

Analogously, we define

$$T_{b,\mu}^{G_b \rightarrow G} : C(G_b(F)_{\text{sr}} // G_b(F), \Lambda) \rightarrow C(G(F)_{\text{sr}} // G(F), \Lambda)$$

by

$$[T_{b,\mu}^{G_b \rightarrow G} f'](g) = \sum_{(g,g',\lambda) \in \text{Rel}_b} f'(g') \dim r_\mu[\lambda].$$

Since r_μ is finite-dimensional, the sum is finite.

Lemma 3.2.8. *The operator $T_{b,\mu}^{G \rightarrow G_b}$ is the zero operator unless $[b]$ is the unique basic class in $B(G, \mu)$.*

Proof. Suppose there exists an F -rational maximal torus $T \subset G$ and a cocharacter $\lambda \in X_*(T)$ such that $r_\mu[\lambda] \neq 0$. Then $\widehat{\mu}$ and $\widehat{\lambda}$ must agree when restricted to the center $Z(\widehat{G})$, which is to say that $\widehat{\mu}$ and $\widehat{\lambda}$ have the same image in $X^*(Z(\widehat{G}))$. Equivalently, if we conjugate μ so as to assume it is a cocharacter of T , then μ and λ have the same image under $X_*(T) \cong \pi_1(T) \rightarrow \pi_1(G)$. By Fact 3.2.3 and the functoriality of κ the image of λ in $\pi_1(G)_\Gamma$ equals $\kappa(b)$. We conclude that $\kappa([b])$ equals the image of μ in $\pi_1(G)_\Gamma$. This means that $[b]$ is the unique basic class in $B(G, \mu)$. \square

Assume therefore that $[b]$ is the unique basic class in $B(G, \mu)$. We may define a ‘‘truncation’’ $\text{Rel}_{b,\mu} \subset \text{Rel}_b$, consisting of conjugacy classes of triples (g, g', λ) for which $\lambda \leq \mu$. Then the kernel function $(g, g', \lambda) \mapsto \dim r_\mu[\lambda]$ is supported on $\text{Rel}_{b,\mu}$. In the diagram

$$\begin{array}{ccc} & \text{Rel}_{b,\mu} & \\ & \swarrow \quad \searrow & \\ G(F)_{\text{rs}} // G(F) & & G_b(F)_{\text{rs}} // G_b(F), \end{array} \tag{3.2.2}$$

both maps are finite étale over their respective images.

The following theorem is proved in §3.4. It relates the operator $T_{b,\mu}^{G \rightarrow G_b}$ to the local Jacquet-Langlands correspondence for G .

Theorem 3.2.9. *Assume that $b \in B(G, \mu)$ is basic, and that Λ is an algebraically closed field of characteristic 0. Let $\phi: W_F \rightarrow {}^L G$ be a discrete Langlands parameter with coefficients in Λ , and let $\rho \in \Pi_\phi(G_b)$. Let $\Theta_\rho \in C(G_b(F)_{\text{sr}} // G_b(F), \Lambda)$ be its Harish-Chandra character. Finally, let $d = \langle 2\rho_G, \mu \rangle$. Then for any $g \in G(F)_{\text{sr}}$ that transfers to $G_b(F)$, we have*

$$\left[T_{b, \mu}^{G_b \rightarrow G} \Theta_\rho \right] (g) = (-1)^d \sum_{\pi \in \Pi_\phi(G)} \dim \text{Hom}_{S_\phi}(\delta_{\pi, \rho}, r_\mu) \Theta_\pi(g). \quad (3.2.3)$$

Remark 3.2.10. When g does not transfer to $G_b(F)$, the left-hand side is zero by definition, but the right-hand side need not be zero, as the following example shows.

Example 3.2.11. Let $G = \text{GL}_2$, and let $\mu: \mathbf{G}_m \rightarrow \mathbf{G}$ the co-character sending x to the diagonal matrix with entries $(x, 1)$. We have $\pi_1(G) = \mathbf{Z}$ as a trivial Γ -module. Let $b \in B(G, \mu)$ be the basic class. Then b corresponds to the isocrystal of slope $1/2$, and $G_b(F)$ is the multiplicative group of the nonsplit quaternion algebra over F . Let $\phi: W_F \rightarrow {}^L G = \text{GL}_2(\mathbf{C})$ be an irreducible representation. The L -packets $\Pi_\phi(G) = \{\pi\}$ and $\Pi_\phi(G_b) = \{\rho\}$ are singletons, with π and ρ supercuspidal. We have $S_\phi = Z(\widehat{G}) = \mathbf{C}^\times$, and $\delta_{\pi, \rho}$ is the identity character of S_ϕ . The representation r_μ is the standard representation of $\widehat{G} = \text{GL}_2(\mathbf{C})$, and $\dim \text{Hom}_{S_\phi}(\delta_{\pi, \rho}, r_\mu) = 2$. Finally, $d = 1$. Therefore the right-hand side of (3.2.3) equals $-2\Theta_\pi(g)$.

Let $w\mu$ be the cocharacter sending x to the diagonal matrix with entries $(1, x)$. The map $\lambda \mapsto r_\mu[\lambda]$ sends μ and $w\mu$ to 1 and all other co-characters to 0. For any strongly regular $g' \in G_b(F)$ there is a unique $G(F)$ -conjugacy class of strongly regular $g \in G(F)$ related to g' . Let $S \subset G$ be the centralizer of one such g . Then $X_*(S) \cong \mathbf{Z}[\Gamma_{E/F}]$ for a quadratic extension E/F and the map $\pi_1(S)_\Gamma \rightarrow \pi_1(G)_\Gamma$ is an isomorphism. There are two elements $\lambda, w\lambda \in X_*(S)$ that map to $\text{inv}[b](g, g')$. Therefore $T_{b, \mu}^{G \rightarrow G_b} f(g') = 2f(g)$. Setting $f = \Theta_\rho$, we find that Theorem 3.2.9 reduces to the Jacquet-Langlands character identity

$$\Theta_\rho(g') = -\Theta_\pi(g).$$

In this situation the functions $T_{b, \mu}^{G \rightarrow G_b} \Theta_\pi$ and $-2\Theta_\rho$ are equal.

We can switch the roles of the two groups and let G be the non-trivial inner form of GL_2 . We can still choose the same μ and b . Then G_b is isomorphic to GL_2 . Now, not all $g' \in G_b(F)$ transfer to $G(F)$; only the elliptic ones do. Therefore the functions $T_{b, \mu}^{G \rightarrow G_b} \Theta_\pi$ and $-2\Theta_\rho$ are not equal, because the former is zero on all non-elliptic g' , while the latter is not.

3.3 Interaction with orbital integrals

Let Λ be an algebraically closed field of characteristic zero. For a topological space X , we let $C_c(X, \Lambda)$ be the space of compactly supported locally constant Λ -valued functions. The space of distributions $\text{Dist}(G(F), \Lambda)$ is the Λ -linear dual of $C_c(G(F), \Lambda)$. The subspace of invariant distributions $\text{Dist}(G(F), \Lambda)^{G(F)}$ is the linear dual of the space of coinvariants $C_c(G(F), \Lambda)_{G(F)}$.

Given a Λ -valued Haar measure dx on $G(F)$, integration against a function $f \in C(G(F)_{\text{sr}} // G(F), \Lambda)$ is a $G(F)$ -invariant distribution on $G(F)$. Due to the functions in $C_c(G(F), \Lambda)$ being locally constant and having compact support, the “integral” is in reality a finite sum. A “stable”

variant on the Weyl integration formula can be used to compute this distribution in terms of orbital integrals. Before we can explain this, we need to discuss choices of measures.

Choose a Λ -valued Haar measure on F . Then a choice of an element $\eta \in \bigwedge^{\dim(G)}(\mathrm{Lie}(G)(F)^*) = \bigwedge^{\dim(G)}(\mathrm{Lie}(G)^*)(F)$ leads to a Λ -valued Haar measure dx_η on $G(F)$. We would like to be able to assign, more generally, a Λ -valued Haar measure on $G(F)$ to any element of $\bigwedge^{\dim(G)}(\mathrm{Lie}(G)^*)(\overline{F})$. To do this, we fix a compatible system $p_n \in \Lambda^\times$, $n = 1, 2, 3, \dots$, of roots of p , i.e. $p_1 = p$ and $p_n^{(n/m)} = p_m$ for all $m|n$. This produces a map $\mathbf{Q} \rightarrow \Lambda$, $r/s \mapsto p_s^r$, which we denote by $t \mapsto p^t$. For any $x \in \overline{F}^\times$ we then define $|x|_\Lambda \in \Lambda$ by $|x|_\Lambda = q^{-\mathrm{ord}_F(x)}$, where $\mathrm{ord}_F : \overline{F}^\times \rightarrow \mathbf{Q}$ is the unique extension of the normalized valuation $F^\times \rightarrow \mathbf{Z}$, and q is the size of the residue field of F , an integral power of p . Now, given $\eta \in \bigwedge^{\dim(G)}(\mathrm{Lie}(G)^*)(\overline{F})$, there exists $a \in \overline{F}^\times$ such that $a\eta \in \bigwedge^{\dim(G)}(\mathrm{Lie}(G)^*)(F)$. The Haar measure $dx_\eta := |a|_\Lambda^{-1} dx_{a\eta}$ of $G(F)$ depends only on η , and not on a .

This procedure allows us to choose Haar measures compatibly in the following two situations. First, consider the inner forms G and G_b . They are canonically identified over \overline{F} , which gives an identification $\bigwedge^{\dim(G)}(\mathrm{Lie}(G)^*)(\overline{F}) = \bigwedge^{\dim(G)}(\mathrm{Lie}(G_b)^*)(\overline{F})$. Haar measures on $G(F)$ and $G_b(F)$ corresponding to the same η will be called compatible. Second, consider two maximal tori $T \subset G$ and $T_b \subset G_b$. Over \overline{F} , the torus T is conjugate to T_b in $G_{\overline{F}} = G_{b,\overline{F}}$. The resulting isomorphism $T_{\overline{F}} \rightarrow T_{b,\overline{F}}$ is well-defined up to the action of the Weyl group. The action of the Weyl group on $\bigwedge^{\dim(G)}(\mathrm{Lie}(T)^*)(\overline{F})$ is through an $\mathcal{O}_{\overline{F}}^\times$ -valued character. Therefore, an element of $\bigwedge^{\dim(G)}(\mathrm{Lie}(T)^*)(\overline{F})$ produces the same Haar measure on $T(F)$ as any of its Weyl conjugates. Since the Weyl orbits in $\bigwedge^{\dim(G)}(\mathrm{Lie}(T)^*)(\overline{F})$ are naturally identified with those in $\bigwedge^{\dim(G)}(\mathrm{Lie}(T_b)^*)(\overline{F})$, we can speak of compatibility of measures on $T(F)$ and $T_b(F)$. In fact, choosing a Haar measure on one maximal torus of G provides compatible choices on all maximal tori of all inner forms.

From now on we assume that the Haar measures on $G(F)$ and $G_b(F)$ have been chosen compatibly, and the Haar measures on all tori of G and G_b have been chosen compatibly.

We now return to the discussion of distributions. For $\phi \in C_c(G(F), \Lambda)$, let $\phi_G \in C(G(F)_{\mathrm{sr}} // G(F), \Lambda)$ be the orbital integral function,

$$\phi_G(y) = \int_{x \in G(F)/G(F)_y} \phi(xy x^{-1}) dx.$$

Our formula for the distribution attached to f takes the form

$$\int_{G(F)} f(x) \phi(x) dx = \langle f, \phi_G \rangle_G. \quad (3.3.1)$$

We explain now the notation $\langle f, \phi_G \rangle_G$. For a function $g \in C(G(F)_{\mathrm{sr}} // G(F), \Lambda)$, we define

$$\langle f, g \rangle_G = \sum_T |W(T, G)(F)|^{-1} \int_{t \in T(F)_{\mathrm{sr}}} |D(t)| \sum_{t_0 \sim t} f(t_0) g(t_0) dt,$$

where

- T runs over a set of representatives for the stable classes of maximal tori,

- $W(T, G) = N(T, G)/T$ is the absolute Weyl group,
- $D(t) = \det \left(\text{Ad}(t) - 1 \mid \text{Lie } G / \text{Lie } T \right)$ is the usual Weyl discriminant, and
- t_0 runs over the $G(F)$ -conjugacy classes inside of the stable class of t .

Note that the integral over T need not exist, i.e. the resulting sum need not be finite. It will do so in the case $g = \phi_G$ for $\phi \in C_c(G(F), \Lambda)$.

The following lemma shows that the operators $T_{b,\mu}^{G \rightarrow G_b}$ and $T_{b,\mu}^{G_b \rightarrow G}$ are adjoint with respect to the pairing $\langle \cdot, \cdot \rangle_G$ and its analogue $\langle \cdot, \cdot \rangle_{G_b}$, defined similarly.

Lemma 3.3.1. *Given $f' \in C(G_b(F)_{\text{sr}} // G_b(F), \Lambda)$ and $g \in C(G(F)_{\text{sr}} // G(F), \Lambda)$ we have*

$$\langle T_{b,\mu}^{G_b \rightarrow G} f', g \rangle_G = \langle f', T_{b,\mu}^{G \rightarrow G_b} g \rangle_{G_b},$$

provided one of the two sides is finite.

Proof. By definition $\langle T_{b,\mu}^{G_b \rightarrow G} f', g \rangle_G$ equals

$$\sum_T |W(T, G)(F)|^{-1} \int_{t \in T(F)_{\text{sr}}} |D(t)| \sum_{t_0 \sim t} \sum_{(t_0, t'_0, \lambda)} r_{\mu, \lambda} f'(t'_0) g(t_0) dt. \quad (3.3.2)$$

The first sum runs over a set of representatives for the stable classes of maximal tori in G . The second sum runs over the set t_0 of $G(F)$ -conjugacy classes of elements that are stably conjugate to t . Let T_{t_0} denote the centralizer of t_0 . The third sum runs over triples (t_0, t'_0, λ) , where t'_0 is a $G_b(F)$ -conjugacy class that is stably conjugate to t_0 , and $\lambda \in X_*(T_{t_0})$ maps to $\text{inv}(t_0, t'_0) \in X_*(T_{t_0})_\Gamma$. Note that if T does not transfer to G_b then it does not contribute to the sum, because the sum over (t_0, t'_0, λ) is empty. Let \mathcal{X} be a set of representatives for those stable classes of maximal tori in G that transfer to G_b . The above expression becomes

$$\sum_{T \in \mathcal{X}} |W(T, G)(F)|^{-1} \int_{t \in T(F)_{\text{sr}}} |D(t)| \sum_{(t_0, t'_0, \lambda)} r_{\mu, \lambda} f'(t'_0) g(t_0) dt,$$

where now the second sum runs over triples (t_0, t'_0, λ) with t_0 a $G(F)$ -conjugacy class and t'_0 a $G_b(F)$ -conjugacy class, both stably conjugate to t , and $\lambda \in X_*(T_{t_0})$ mapping to $\text{inv}(t_0, t'_0) \in X_*(T_{t_0})_\Gamma$.

Let \mathcal{X}' be a set of representatives for those stable classes of maximal tori of G_b that transfer to G . We have a bijection $\mathcal{X} \leftrightarrow \mathcal{X}'$. Fix arbitrarily an admissible isomorphism $T \rightarrow T'$ for any $T \in \mathcal{X}$ and $T' \in \mathcal{X}'$ that correspond under this bijection. It induces an isomorphism $W(T, G) \rightarrow W(T', G_b)$ of finite algebraic groups. Given $t \in T(F)_{\text{reg}}$ let $t' \in T'(F)_{\text{reg}}$ be its image under the admissible isomorphism. Then $|D(t)| = |D(t')|$, and (3.3.2) becomes

$$\sum_{T' \in \mathcal{X}'} |W(T', G_b)(F)|^{-1} \int_{t' \in T'_{\text{reg}}(F)} |D(t')| \sum_{(t_0, t'_0, \lambda)} r_{\mu, \lambda} f'(t'_0) g(t_0) dt, \quad (3.3.3)$$

where now the second sum runs over triples (t_0, t'_0, λ) , where t_0 is a $G(F)$ -conjugacy class, t'_0 is a $G_b(F)$ -conjugacy class, both are stably conjugate to t' , and $\lambda \in X_*(T_{t_0})$ maps to $\text{inv}(t_0, t'_0) \in X_*(T_{t_0})_\Gamma$. Reversing the arguments from the beginning of this proof we see that this expression equals $\langle f', T_{b,\mu}^{G \rightarrow G_b} g \rangle_{G_b}$. \square

This lemma hints at the possibility of extending the operator $T_{b,\mu}^{G \rightarrow G_b}$ from the space of invariant functions on $G(F)$ to the space of invariant distributions on $G(F)$. Given an invariant distribution d on $G(F)$, we would like to define $T_{b,\mu}^{G \rightarrow G_b} d$ by the formula

$$\langle T_{b,\mu}^{G \rightarrow G_b} d, \phi_{G_b} \rangle = \langle d, T_{b,\mu}^{G_b \rightarrow G} \phi_{G_b} \rangle \quad (3.3.4)$$

for any test function $\phi \in C_c(G_b(F), \Lambda)$, where now $\langle -, - \rangle$ denotes the pairing of canonical duality between $\text{Dist}(G(F), \Lambda)^{G(F)}$ and $C_c(G(F), \Lambda)_{G(F)}$, and the corresponding pairing for $G_b(F)$. But at the moment there is no guarantee that $T_{b,\mu}^{G_b \rightarrow G} \phi_{G_b} \in C(G(F)_{\text{sr}} // G(F), \Lambda)$ will be the orbital integral function for a compactly supported test function on $G(F)$. This problem is reminiscent of the problem of endoscopic transfer of functions, where the (stable) orbital integrals of the transfer function are determined explicitly in terms of the transfer factors, but proving the existence of a test function with the prescribed orbital integrals is a highly non-trivial matter. We shall discuss this problem in §6, where we prove that $T_{b,\mu}^{G_b \rightarrow G} \phi_G$ is indeed the orbital integral of a compactly supported test function. Then we can use (3.3.4) to define an operator

$$T_{b,\mu}^{G \rightarrow G_b} : \text{Dist}(G(F), \Lambda)^{G(F)} \rightarrow \text{Dist}(G_b(F), \Lambda)^{G_b(F)}$$

which extends the operator

$$T_{b,\mu}^{G \rightarrow G_b} : C(G(F)_{\text{sr}} // G(F), \Lambda) \rightarrow C(G_b(F)_{\text{sr}} // G_b(F), \Lambda)$$

defined in Definition 3.2.7.

3.4 Proof of Theorem 3.2.9

We now give the proof of Theorem 3.2.9. We will use the notation and results of §A.1.

We are given a discrete Weil parameter ϕ , a representation $\rho \in \Pi_\phi(G_b)$ in its L -packet, and an element $g \in G(F)_{\text{sr}}$. We assume that g is related to an element of $G_b(F)$. This means there exists a triple in Rel_b of the form (g, g', λ) . For the moment we fix such a triple (g, g', λ) .

Let $s \in S_\phi$ be a semi-simple element, and let $\dot{s} \in S_\phi^+$ be a lift of it. Then we have the refined endoscopic datum $\dot{\mathfrak{e}} = (H, \mathcal{H}, \dot{s}, \eta)$ defined in (A.1.1); we choose as in that section a z -pair $\mathfrak{z} = (H_1, \eta_1)$. Then

$$\begin{aligned} & e(G_b) \sum_{\rho' \in \Pi_\phi(G_b)} \text{tr } \tau_{z, \mathfrak{w}, \rho'}(\dot{s}) \Theta_{\rho'}(g') \\ \stackrel{(A.1.2)}{=} & \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g') S_{\Theta_{\phi^s}}(h_1) \\ \stackrel{(A.1.4)}{=} & \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g) \left\langle \text{inv}[b](g, g'), s_{h,g}^\natural \right\rangle S_{\Theta_{\phi^s}}(h_1) \\ = & \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g) \lambda(s_{h,g}^\natural) S_{\Theta_{\phi^s}}(h_1). \end{aligned}$$

We now multiply this expression by the kernel function $\dim r_\mu[\lambda]$, and then sum over all $G_b(F)$ -conjugacy classes of elements $g' \in G_b(F)$ and all $\lambda \in X_*(T_g)$ such that (g, g', λ) lies in Rel_b . We obtain

$$\begin{aligned}
& e(G_b) \sum_{(g', \lambda)} \sum_{\rho' \in \Pi_\phi(G_b)} \text{tr } \tau_{z, \mathfrak{w}, \rho'}(\dot{s}) \Theta_{\rho'}(g') \dim r_\mu[\lambda] \\
&= \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g) S \Theta_{\phi^s}(h_1) \sum_{(g', \lambda)} \lambda(s_{h, g}^\natural) \dim r_\mu[\lambda] \\
&\stackrel{(*)}{=} \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g) S \Theta_{\phi^s}(h_1) \text{tr } r_\mu(s_{h, g}^\natural) \\
&\stackrel{(**)}{=} \text{tr } r_\mu(s^\natural) \sum_{h_1 \in H_1(F)/\text{st}} \Delta(h_1, g) S \Theta_{\phi^s}(h_1) \\
&\stackrel{\text{(A.1.2)}}{=} \text{tr } r_\mu(s^\natural) e(G) \sum_{\pi \in \Pi_\phi(G)} \text{tr } \tau_{z, \mathfrak{w}, \pi}(\dot{s}) \Theta_\pi(g).
\end{aligned}$$

We justify (**): Let $T \subset G$ be the centralizer of g . The image of $s_{h, g}^\natural$ under any admissible embedding $\widehat{T} \rightarrow \widehat{G}$ is conjugate to s^\natural in \widehat{G} and $\text{tr } r_\mu$ is conjugation-invariant. Recall here that $s^\natural \in S_\phi$ is the image of \dot{s} under (2.3.2).

We justify (*): $\lambda \in X_*(T)$ determines the $G_b(F)$ -conjugacy class of g' , since $\text{inv}[b](g, g') \in B(T)$ determines it. Therefore the sum over (g', λ) is in reality a sum only over λ . There exists $g' \in G_b(F)$ with $\kappa(\text{inv}[b](g, g'))$ being the image of λ in $X_*(T)_\Gamma$ if and only if the image of λ under $X_*(T) \rightarrow X_*(T)_\Gamma \rightarrow \pi_1(G)_\Gamma$ equals $\kappa(b)$. Since the image of μ in $\pi_1(G)_\Gamma$ also equals $\kappa(b)$, the sum over (g', λ) is in fact the sum over $\lambda \in X_*(T)$ having the same image as μ in $\pi_1(G)_\Gamma$. In terms of the dual torus \widehat{T} this is the sum over $\lambda \in X^*(\widehat{T})$ whose restriction to $Z(\widehat{G})^\Gamma$ equals that of μ . Since for λ not satisfying this condition the number $\dim r_\mu[\lambda]$ is zero, we may extend the sum to be over all $\lambda \in X^*(T) = X^*(\widehat{T})$.

We now continue with the equation. Multiply both sides of the above equation by $\text{tr } \check{\tau}_{z, \mathfrak{w}, \rho}(\dot{s})$. As functions of $\dot{s} \in S_\phi^+$, both sides then become invariant under $Z(\widehat{G})^+$ and thus become functions of the finite quotient $\bar{S}_\phi = S_\phi^+/Z(\widehat{G})^+ = S_\phi/Z(\widehat{G})^\Gamma$. Now apply $|\bar{S}_\phi|^{-1} \sum_{\bar{s} \in \bar{S}_\phi}$ to both sides to obtain an equality between

$$|\bar{S}_\phi|^{-1} e(G_b) \sum_{\bar{s} \in \bar{S}_\phi} \sum_{(g', \lambda)} \sum_{\rho' \in \Pi_\phi(G_b)} \text{tr } \check{\tau}_{z, \mathfrak{w}, \rho}(\dot{s}) \text{tr } \tau_{z, \mathfrak{w}, \rho'}(\dot{s}) \Theta_{\rho'}(g') \dim r_\mu[\lambda] \quad (3.4.1)$$

and

$$|\bar{S}_\phi|^{-1} e(G) \sum_{\bar{s} \in \bar{S}_\phi} \text{tr } r_\mu(s^\natural) \sum_{\pi \in \Pi_\phi(G)} \text{tr } \check{\tau}_{z, \mathfrak{w}, \rho}(\dot{s}) \text{tr } \tau_{z, \mathfrak{w}, \pi}(\dot{s}) \Theta_\pi(g), \quad (3.4.2)$$

where in both formulas \dot{s} is an arbitrary lift of \bar{s} and $s^\natural \in S_\phi$ is the image of \dot{s} under (2.3.2). Executing the sum over \bar{s} in (3.4.1) gives

$$e(G_b) \sum_{(g', \lambda)} \Theta_\rho(g') \dim r_\mu[\lambda] = e(G_b) [T_{b, \mu}^{G_b \rightarrow G} \Theta_\rho](g).$$

To treat (3.4.2) note that $\check{\tau}_{z,w,\rho} \otimes \tau_{z,w,\pi}(\dot{s}) = \check{\delta}_{\pi,\rho}(s^\natural)$. Furthermore, the composition of the map (2.3.2) with the natural projection $S_\phi \rightarrow S_\phi/Z(\widehat{G})^\Gamma$ is equal to the natural projection $S_\phi^+ \rightarrow S_\phi^+/Z(\widehat{G})^+ = S_\phi/Z(\widehat{G})^\Gamma = \bar{S}_\phi$. Thus s^\natural is simply a lift of \bar{s} to S_ϕ . We find that (3.4.2) equals

$$e(G) |\bar{S}_\phi|^{-1} \sum_{\bar{s} \in \bar{S}_\phi} \text{tr } r_\mu(s^\natural) \text{tr } \check{\delta}_{\pi,\rho}(s^\natural) = e(G) \dim \text{Hom}_{S_\phi}(\delta_{\pi,\rho}, r_\mu).$$

We have now reduced Theorem 3.2.9 to the identity

$$e(G)e(G_b) = (-1)^{\langle 2\rho_G, \mu \rangle}, \quad (3.4.3)$$

where ρ_G is the sum of the positive roots. Recall that G^* is a quasi-split inner form of G . Let $\mu_1, \mu_2 \in X^*(Z(\widehat{G}_{\text{sc}})^\Gamma)$ be the elements corresponding to the inner twists $G^* \rightarrow G$ and $G^* \rightarrow G_b$ by Kottwitz's homomorphism [Kot86, Theorem 1.2]. By Lemma A.2.1 we have $e(G_b)e(G) = (-1)^{\langle 2\rho, \mu_2 - \mu_1 \rangle}$. But since G_b is obtained from G by twisting by b , the difference $\mu_2 - \mu_1$ is equal to the image of $\kappa(b) \in X^*(Z(\widehat{G})^\Gamma)$ under the map $X^*(Z(\widehat{G})^\Gamma) \rightarrow X^*(Z(\widehat{G}_{\text{sc}})^\Gamma)$ dual to the natural map $Z(\widehat{G}_{\text{sc}}) \rightarrow Z(\widehat{G})$. Since $b \in B(G, \mu)$ we see that $\mu_2 - \mu_1 = \mu$ and (3.4.3) follows. The proof of Theorem 3.2.9 is complete.

4 The Lefschetz-Verdier trace formula for v-stacks

The goal of this section is to build up some machinery related to the Lefschetz-Verdier trace formula.

We briefly review the setup in the context of a separated finite type morphism of schemes $p: X \rightarrow \text{Spec } k$, where k is an algebraically closed field. Let ℓ be a prime unequal to the characteristic of k , and let A be an object of $D(X_{\text{ét}}, \overline{\mathbf{Q}}_\ell)$, the derived category of étale $\overline{\mathbf{Q}}_\ell$ -sheaves on X . Suppose $f: X \rightarrow X$ is a k -linear endomorphism. If we are given the additional datum of a morphism¹ $Rf_! A \rightarrow A$, we obtain an operator $Rp_! A \cong Rp_! Rf_! A \rightarrow Rp_! A$ on the compactly supported cohomology $Rp_! A = R\Gamma_c(X, A)$.

In the special case that X is proper over k , so that $R\Gamma_c(X, A) = R\Gamma(X, A)$, the Lefschetz-Verdier trace formula [SGA77], [Var07] expresses $\text{tr } f | R\Gamma(X, A)$ in terms of data living on the fixed point locus $\text{Fix}(f)$ of f . In particular, at isolated fixed points $x \in \text{Fix}(f)$, there are local terms $\text{loc}_x(f, A)$, and if all fixed points are isolated, then $\text{tr } f | R\Gamma(X, A)$ is the sum of the $\text{loc}_x(f, A)$.

In order to apply the Lefschetz-Verdier trace formula, we need to assume that A satisfies a suitable finiteness hypothesis (constructible of finite Tor dimension). Under this hypothesis one establishes an isomorphism [SGA77, Exposé III, (3.1.1)]

$$\mathbf{D}A \boxtimes_k^{\mathbf{L}} A \cong \underline{\text{RHom}}(\text{pr}_1^* A, \text{pr}_2^! A) \quad (4.0.1)$$

in $D((X \times_k X)_{\text{ét}}, \overline{\mathbf{Q}}_\ell)$, where $\text{pr}_1, \text{pr}_2: X \times_k X \rightarrow X$ are the projection maps, and \mathbf{D} is Verdier duality relative to k . Once (4.0.1) is established, the definition of local terms and the validity of the Lefschetz-Verdier trace formula can be derived by applying Grothendieck's six functor formalism. The special case where $X = \text{Spec } k$ is instructive; the finiteness condition on A is that it be a perfect complex of $\overline{\mathbf{Q}}_\ell$ -vector spaces, and then (4.0.1) reduces to the fact that $A^\vee \otimes^{\mathbf{L}} A \rightarrow \underline{\text{RHom}}(A, A)$ is an isomorphism. This allows us to express the trace of an endomorphism $f \in \text{End } A$ as the image of f under the evaluation map $A^\vee \otimes^{\mathbf{L}} A \rightarrow \overline{\mathbf{Q}}_\ell$.

In this section we extend the Lefschetz-Verdier trace formula to the setting of perfectoid spaces, diamonds, and v-stacks. The main result is Theorem 4.3.7 and its corollary (4.3.5). We very closely follow the formalism of [LZ20] concerning the symmetric monoidal 2-category of cohomological correspondences. In both the schematic and perfectoid settings, the finiteness condition required of the object A can be stated in terms of the property of *universal local acyclicity* (ULA); as noted in [FS21, Theorem IV.2.24], this is precisely the hypothesis necessary to obtain the isomorphism in (4.0.1).

The statement of Lefschetz-Verdier is formally identical between the scheme and perfectoid settings. But in the perfectoid setting, there arises the possibility that the fixed point locus $\text{Fix}(f)$ has the structure of a locally profinite set, in which case the local terms appearing in Lefschetz-Verdier are not a function on $\text{Fix}(f)$, but rather a *distribution* on $\text{Fix}(f)$. This observation is critical to our applications.

For our applications, we have included two additional theorems concerning local terms in the perfectoid setting, which could also have been stated in the schematic setting and may be of

¹Equivalently, a morphism $A \rightarrow Rf^! A$. A special case occurs when f is an automorphism, A is an honest sheaf on $X_{\text{ét}}$, and $A \rightarrow f^* A$ is a morphism, such as the identity morphism on the constant sheaf $A = \overline{\mathbf{Q}}_\ell$. More generally, we may replace f with an algebraic correspondence $c = (c_1, c_2): C \rightarrow X \times_k X$. In that setting, the required extra datum is a morphism $c_1^* \mathcal{F} \rightarrow c_2^* \mathcal{F}$: this is the notion of a *cohomological correspondence* lying over c .

independent interest. Theorem 4.5.3 is a sort of Künneth isomorphism for local terms on a fiber product of stacks. Theorem 4.6.1 states that in the situation of a smooth group G acting on a diamond X , the local terms corresponding to individual elements $g \in G$ agree with local terms computed on the quotient stack $[X/G]$.

4.1 Small v-stacks and the six-functor formalism

We recall here some material from [Sch17] on the main classes of geometric objects we deal with: perfectoid spaces, diamonds, and v-stacks. Let Perf be the category of perfectoid spaces in characteristic p . There are four topologies we consider on Perf , which we list from coarsest to finest: the analytic topology, the étale topology, the pro-étale topology, and the v-topology.

The v-topology is a rough analogy for the fpqc topology on schemes. All representable presheaves on Perf are sheaves for the v-topology [Sch17, Theorem 1.2].

A *diamond* is a pro-étale sheaf on Perf of the form X/R , where X is a perfectoid space and $R \subset X \times X$ is a pro-étale equivalence relation. Diamonds are automatically v-sheaves [Sch17, Proposition 11.9]. A particularly well-behaved class of diamonds are the locally spatial diamonds [Sch17, Definition 1.4]. There is a natural functor $X \mapsto X^\diamond$ from analytic adic spaces over \mathbf{Z}_p to locally spatial diamonds.

A v-sheaf Y on Perf is *small* if there exists a surjective map $X \rightarrow Y$ from a perfectoid space² X . A *small v-stack* [Sch17, Definition 12.4] is a v-stack Y on Perf such that there exists a surjective map $X \rightarrow Y$ from a perfectoid space, such that $X \times_Y X$ is a small v-sheaf.

Let Λ be a ring which is n -torsion for some n prime to p . For every v-stack X there is a triangulated category $D_{\text{ét}}(X, \Lambda)$ [Sch17, Definition 1.7]. If X is a locally spatial diamond, then $D_{\text{ét}}(X, \Lambda)$ is equivalent to the left-completion of the derived category of sheaves of Λ -modules on the étale topology of X [Sch17, Proposition 14.15].

The familiar six functors of Grothendieck have analogues in the world of small v-stacks [Sch17, Definition 1.7]. There is a derived tensor product $\otimes_{\Lambda}^{\mathbf{L}}$ and a derived internal hom $\underline{\text{RHom}}_{\Lambda}$. For any morphism $f: Y \rightarrow X$ of small v-stacks, there is a pair of adjoint functors f^* and Rf_* .

We propose for convenience the following definition.

Definition 4.1.1. A morphism $f: Y \rightarrow X$ is *representable in nice diamonds* or simply *nice* if it is compactifiable [Sch17, Definition 22.2], representable in locally spatial diamonds [Sch17, Definition 13.3], and if f has finite geometric transcendence degree [Sch17, Definition 22.2].

If $f: Y \rightarrow X$ is representable in nice diamonds, then there are an adjoint pair of functors $Rf_!$ and $Rf^!$ [Sch17, Sections 22 and 23].

Theorem 4.1.2 ([Sch17, Theorem 1.8]). *The six operations \otimes_{Λ} , $\underline{\text{RHom}}_{\Lambda}$, f^* , Rf_* , $Rf_!$, and $Rf^!$ obey the rules:*

$$(P1.) \quad f^* A \otimes_{\Lambda}^{\mathbf{L}} f^* B \cong f^*(A \otimes_{\Lambda}^{\mathbf{L}} B),$$

$$(P2.) \quad Rf_* \underline{\text{RHom}}_{\Lambda}(f^* A, B) \cong \underline{\text{RHom}}_{\Lambda}(A, Rf_* B),$$

$$(P3.) \quad Rf_!(A \otimes_{\Lambda}^{\mathbf{L}} f^* B) \cong Rf_! A \otimes_{\Lambda}^{\mathbf{L}} B \quad (\text{the projection formula}),$$

²There is also a smallness condition on X relative to a cutoff cardinal. We will be ignoring set-theoretic issues.

(P4.) $\underline{\mathrm{RHom}}_{\Lambda}(Rf_!A, B) \cong Rf_*\underline{\mathrm{RHom}}_{\Lambda}(A, Rf^!B)$ (local Verdier duality),

(P5.) $Rf^!\underline{\mathrm{RHom}}_{\Lambda}(A, B) \cong \underline{\mathrm{RHom}}_{\Lambda}(f^*A, Rf^!B)$

We also need the following base change results.

Theorem 4.1.3 ([Sch17, Theorem 1.9]). *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{\tilde{g}} & Y \\ f' \downarrow & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array} \quad (4.1.1)$$

be a cartesian diagram of small v -stacks.

(BC1.) *If f is representable in nice diamonds, then $g^*Rf_! \cong Rf'_!\tilde{g}^*$.*

(BC2.) *If g is representable in nice diamonds, then $Rg^!Rf_* \cong Rf'_*R\tilde{g}^!$.*

There is a notion of *cohomological smoothness* [Sch17, Definition 23.8] for morphisms between v -stacks which are representable in nice diamonds. Let Λ be an n -torsion ring for some n not divisible by p . For a morphism $f: Y \rightarrow X$ of small v -stacks which is representable in nice diamonds, there is a natural map of functors

$$Rf^!\Lambda_X \otimes_{\Lambda}^{\mathbf{L}} f^* \rightarrow Rf^!, \quad (4.1.2)$$

adjoint to

$$Rf_!(Rf^!\Lambda_X \otimes_{\Lambda}^{\mathbf{L}} f^* A) \xrightarrow{(P3)} Rf_!Rf^!\Lambda_X \otimes_{\Lambda}^{\mathbf{L}} A \xrightarrow{\mathrm{counit}} A.$$

If f is cohomologically smooth, then (4.1.2) is an equivalence [Sch17, Theorem 1.10]. Furthermore, the object $Rf^!\Lambda_X$ is invertible in the monoidal category $D_{\acute{e}t}(Y, \Lambda)$. (For X a locally spatial diamond, an object A of $D_{\acute{e}t}(X, \Lambda)$ is invertible if and only if v -locally on X there is an isomorphism $A \cong \Lambda_X[n]$.)

There is also the following base change theorem involving cohomologically smooth morphisms.

Theorem 4.1.4 ([Sch17, Theorem 1.10]). *In the cartesian diagram (4.1.1), assume that f is cohomologically smooth. Then $\tilde{g}^*Rf_! \cong R(f')^!g^*$ and $(f')^*Rg^! \cong R\tilde{g}^!f^*$.*

Finally, we need the notion of the dualizing complex for v -stacks.

Definition 4.1.5 (The dualizing complex). Let $f: X \rightarrow S$ be a morphism of small v -stacks which is representable in nice diamonds. We define $K_{X/S} = Rf^!\Lambda$, an object in $D_{\acute{e}t}(X, \Lambda)$.

Suppose we are in the special case that $S = \mathrm{Spd} C$ for an algebraically closed perfectoid field C , and that f is proper. Then $Rf_* = Rf_!$, and so there is a morphism $Rf_*K_{X/S} = Rf_!Rf^!\Lambda \xrightarrow{\mathrm{counit}} \Lambda$, which induces a morphism on the level of global sections

$$H^0(X, K_{X/S}) = H^0(S, Rf_*K_{X/S}) \rightarrow H^0(S, \Lambda) = \Lambda,$$

which we notate as $\omega \mapsto \int_X \omega$.

We record the following lemmas for convenience.

Lemma 4.1.6. *Let $f: Y \rightarrow X$ be a morphism of S -Artin v-stacks, and let $A, I \in D_{\text{ét}}(X, \Lambda)$ with I invertible. The natural map $Rf^!A \otimes_{\Lambda}^{\mathbf{L}} f^*I \rightarrow Rf^!(A \otimes I)$ of (4.1.2) is an isomorphism.*

Proof. Let $w_{A,I}$ be this morphism. We also get a map $w_{A \otimes I, I^{-1}}: Rf^!(A \otimes_{\Lambda}^{\mathbf{L}} I) \otimes f^*I^{-1} \rightarrow Rf^!A$, which induces $Rf^!(A \otimes I) \rightarrow Rf^!A \otimes f^*I$. This is the inverse to $w_{A,I}$. \square

Lemma 4.1.7. *Let $f: Y \rightarrow X$ be a morphism of S -Artin v-stacks which is cohomologically smooth. There is a canonical isomorphism*

$$R\Delta_{Y/X}^! \Lambda_{Y \times_X Y} \otimes_{\Lambda}^{\mathbf{L}} Rf^! \Lambda_X \cong \Lambda_Y, \quad (4.1.3)$$

so that $K_{Y/Y \times_X Y} \cong K_{Y/X}^{-1}$ is invertible.

Proof. Let $\text{pr}_1, \text{pr}_2: Y \times_X Y \rightarrow Y$ be the projection morphisms; each is cohomologically smooth. We have

$$\Lambda_Y \cong \text{Rid}_Y^! \Lambda_Y \cong R\Delta_{Y/X}^! R\text{pr}_1^! \Lambda_Y \cong R\Delta_{Y/X}^! \Lambda_{Y \times_X Y} \otimes_{\Lambda}^{\mathbf{L}} \Delta_{Y/X}^* R\text{pr}_1^! \Lambda_Y,$$

where in the last isomorphism we used Lemma 4.1.6 combined with the cohomological smoothness of $R\text{pr}_1$. Now use Theorem 4.1.4 to obtain an isomorphism $\Delta_{Y/X}^* R\text{pr}_1^! f^* \Lambda_X \cong \Delta_{Y/X}^* \text{pr}_2^* Rf^! \Lambda_X \cong Rf^! \Lambda_X$. \square

For the remainder of the section, we fix Λ , an n -torsion ring for some n not divisible by p . We will now start writing $f_!$ for $Rf_!$ and \otimes for $\otimes_{\Lambda}^{\mathbf{L}}$, etc.

4.2 Artin v-stacks

In our applications, we will encounter stacky maps $f: Y \rightarrow X$ between v-stacks, for which $f_!$ and $f^!$ are not available using the material in [Sch17]. In [GHW], the full six-functor formalism is extended to a class of stacky maps between Artin v-stacks, using the ∞ -categorical machinery of [LZ].

Definition 4.2.1 ([FS21, Definition IV.1.1]). An *Artin v-stack* is a small v-stack X such that the diagonal $\Delta_X: X \rightarrow X$ is representable in locally spatial diamonds, and such that there is some surjective map $g: U \rightarrow X$ from a locally spatial diamond U such that g is separated and cohomologically smooth.

A morphism of v-stacks $f: Y \rightarrow X$ is *smooth-locally nice* if there exists a surjective cohomologically smooth map $g: U \rightarrow Y$ from a locally spatial diamond U such that $f \circ g$ is nice.

Similarly we define for a fixed small v-stack S the notion of an S -Artin v-stack $\mathfrak{X} \rightarrow S$, by requiring that the relative diagonal $\Delta_{\mathfrak{X}/S}: \mathfrak{X} \rightarrow \mathfrak{X} \times_S \mathfrak{X}$ is representable in locally spatial diamonds.

Theorem 4.2.2 ([GHW]). *If $f: Y \rightarrow X$ is any smooth-locally nice map of Artin v-stacks, there exist functors $f_!$ and $f^!$ satisfying the properties listed in Theorems 4.1.2 and 4.1.3, agreeing with the constructions in [Sch17] when f is nice.*

Definition 4.2.3 ([FS21, Definition IV.1.10]). Let $f: Y \rightarrow X$ be a map of Artin v-stacks. Then f is *cohomologically smooth* if there exists a separated, representable in locally spatial diamonds, and cohomologically smooth surjection $g: U \rightarrow Y$ from a locally spatial diamond such that $f \circ g: U \rightarrow X$ is nice and cohomologically smooth.

4.3 The category of cohomological correspondences

The Lefschetz-Verdier trace formula was expressed elegantly by Lu and Zheng [LZ20] in the language of symmetric monoidal 2-categories. [LZ20] works with schemes, but our treatment of Artin v-stacks is virtually identical. First we recall some definitions and constructions concerning the categorical trace.

Definition 4.3.1 ([LZ20, Definition 1.1, Construction 1.6]). An object X of a symmetric monoidal category $(\mathcal{C}, \otimes, 1_{\mathcal{C}})$ is *dualizable* if there exists an object X^{\vee} together with morphisms $\text{ev}_X: X^{\vee} \otimes X \rightarrow 1_{\mathcal{C}}$ and $\text{coev}_X: 1_{\mathcal{C}} \rightarrow X \otimes X^{\vee}$, such that the compositions

$$X \xrightarrow{\text{coev}_X \otimes \text{id}_X} X \otimes X^{\vee} \otimes X \xrightarrow{\text{id}_X \otimes \text{ev}_X} X$$

and

$$X^{\vee} \xrightarrow{\text{id}_{X^{\vee}} \otimes \text{coev}_X} X^{\vee} \otimes X \otimes X^{\vee} \xrightarrow{\text{ev}_X \otimes \text{id}_X} X^{\vee}$$

are isomorphic to the identities on X and X^{\vee} , respectively. Equivalently, the functor $Y \mapsto X \otimes Y$ has right adjoint $Y \mapsto X^{\vee} \otimes Y$. If X is dualizable, then $X^{\vee} \otimes Y$ serves as an internal mapping object $\underline{\text{Hom}}(X, Y)$.

Let $\Omega_{\mathcal{C}} = \text{End}(1_{\mathcal{C}})$ be the (1-)category of endomorphisms of the unit object of \mathcal{C} . Let $f \in \text{End } X$ be an endomorphism of a dualizable object X . Define the categorical trace $\text{tr}(f)$ as the composite:

$$1_{\mathcal{C}} \xrightarrow{\text{coev}_X} X \otimes X^{\vee} \xrightarrow{f \otimes \text{id}_{X^{\vee}}} X \otimes X^{\vee} \xrightarrow{\text{ev}_X} 1_{\mathcal{C}}$$

so that $\text{tr}(f)$ is an object of $\Omega_{\mathcal{C}}$.

Definition 4.3.2 (The category of correspondences). We define a symmetric monoidal 2-category Corr_S . The objects of Corr_S are S -Artin v-stacks, which are smooth-locally nice over S . Given objects X and X' , the category $\text{Hom}_{\text{Corr}_S}(X, X')$ has for its objects the correspondences:

$$\begin{array}{ccc} & C & \\ c_1 \swarrow & & \searrow c_2 \\ X & & X' \end{array} \quad (4.3.1)$$

with each c_i a morphism of S -Artin v-stacks, with c_2 assumed to be smooth-locally nice. The composition of $(c_1, c_2): C \rightarrow X \times_S X'$ with $(d_1, d_2): D \rightarrow X' \times_S X''$ is the correspondence $(c_1 d'_1, d_2 c'_2)$ defined by the diagram:

$$\begin{array}{ccccc} & & C \times_{X'} D & & \\ & & d'_1 \swarrow & \searrow c'_2 & \\ & C & & & D \\ c_1 \swarrow & & & & \searrow d_2 \\ X & & & & X'' \\ & & c_2 \searrow & \swarrow d_1 & \\ & & X' & & \end{array} \quad (4.3.2)$$

If $c = (c_1, c_2): C \rightarrow X \times_S X'$ and $c' = (c'_1, c'_2): C' \rightarrow X \times_S X'$ represent two objects in $\text{Hom}_{\text{Corr}_S}(X_1, X_2)$, a 2-morphism $c \implies c'$ is a *proper* morphism $p: C \rightarrow C'$ such that the following diagram commutes:

$$\begin{array}{ccc}
 & C & \\
 c_1 \swarrow & & \searrow c_2 \\
 X & & X' \\
 c'_1 \swarrow & p \downarrow & \searrow c'_2 \\
 & D &
 \end{array} \tag{4.3.3}$$

The monoidal structure is defined by $X_1 \otimes X_2 = X_1 \times_S X_2$, with unit object S . Given $c = (c_1, c_2): C \rightarrow X_1 \times_S X'_1$ and $d = (d_1, d_2): D \rightarrow X_2 \times_S X'_2$ representing objects in $\text{Hom}_{\text{Corr}_S}(X_1, X'_1)$ and $\text{Hom}_{\text{Corr}_S}(X_2, X'_2)$ respectively, we define the object $c \otimes d$ of $\text{Hom}_{\text{Corr}_S}(X_1 \times_S X_2, X'_1 \times_S X'_2)$ as the correspondence:

$$\begin{array}{ccc}
 & C \times_S D & \\
 c_1 \times_S d_1 \swarrow & & \searrow c_2 \times_S d_2 \\
 X_1 \times_S X_2 & & X'_1 \times_S X'_2
 \end{array} \tag{4.3.4}$$

Definition 4.3.3 (The category of cohomological correspondences). We define a symmetric monoidal 2-category CoCorr_S , which comes equipped with a functor to Corr_S . An object of CoCorr_S is a pair (X, A) , where X is an S -Artin v-stack which is smooth-locally nice over S , and $A \in D_{\text{ét}}(X, \Lambda)$. Given objects $\mathfrak{X} = (X, A)$ and $\mathfrak{X}' = (X', A')$ of CoCorr_S , the category $\text{Hom}_{\text{CoCorr}_S}(\mathfrak{X}, \mathfrak{X}')$ consists of pairs (c, u) , where $c = (c_1, c_2)$ is a correspondence as in (4.3.1), and $u: c_1^* A \rightarrow c_2^* A'$ is a morphism in $D_{\text{ét}}(C, \Lambda)$. The composition of $(c, u): (X, A) \rightarrow (X', A')$ with $(d, v): (X', A') \rightarrow (X'', A'')$ is (e, w) , where $e: C \times_{X'} D \rightarrow X \times_S X''$ is the correspondence in (4.3.2), and w is the composition

$$(d'_1)^* c_1^* A \xrightarrow{u} (d'_1)^* c_2^* A' \xrightarrow{\alpha} (c'_2)^! d_1^* A' \xrightarrow{v} (c'_2)^! d_2^* A''$$

where α is adjoint to the base change isomorphism $(c'_2)_!(d'_1)^* \cong d_1^*(c_2)_!$. If $\mathfrak{f} = (c, u)$ and $\mathfrak{f}' = (c', u')$ are two objects in $\text{Hom}_{\text{CoCorr}_S}(\mathfrak{X}, \mathfrak{X}')$, a 2-morphism $\mathfrak{f} \implies \mathfrak{f}'$ is a proper morphism p as in (4.3.3), such that u' is the composition

$$\begin{aligned}
 (c'_1)^* A &\xrightarrow{\text{unit}} p_* p^* (c'_1)^* A \cong p_* c_1^* A \xrightarrow{u} p_* c_2^* A' \\
 &\cong p_* p^! (c'_2)^! A' \cong p_! p^! (c'_2)^! A' \xrightarrow{\text{counit}} (c'_2)^! A'.
 \end{aligned}$$

The symmetric monoidal structure on Corr_S is given by $(X, A) \otimes (X', A') = (X \times_S X', A \boxtimes_S A')$. The unit object is $1_{\text{CoCorr}_S} = (S, \Lambda_S)$. Finally, given $(c, u): (X_1, A_1) \rightarrow (X'_1, A'_1)$ and $(d, v): (X_2, A_2) \rightarrow (X'_2, A'_2)$, the tensor product $(c, u) \otimes (d, v)$ is $(c \otimes d, w)$, where $c \otimes d$ is the correspondence in (4.3.4), and w is the composition

$$(c_1 \times_S d_1)^* (A_1 \boxtimes_S A_2) \cong c_1^* A_1 \boxtimes d_1^* A_2 \xrightarrow{u \boxtimes_S v} c_2^* A'_1 \boxtimes_S d_2^* A'_2 \xrightarrow{\kappa} (c_2 \times_S d_2)^! (A_2 \boxtimes_S A'_2),$$

where κ is adjoint to the Künneth isomorphism $(c_2 \times_S d_2)_!(B_1 \boxtimes_S B_2) \cong (c_2)_! B_1 \boxtimes_S (d_2)_! B_2$.

The category CoCorr_S has internal mapping objects: if $\mathfrak{X}_1 = (X_1, A_1)$ and $\mathfrak{X}_2 = (X_2, A_2)$, then $\underline{\mathrm{Hom}}(\mathfrak{X}_1, \mathfrak{X}_2) = (X_1 \times_S X_2, \underline{\mathrm{RHom}}(\mathrm{pr}_1^* A_1, \mathrm{pr}_2^! A_2))$, where $\mathrm{pr}_i: X_1 \times_S X_2 \rightarrow X_i$ is the projection.

We have the following characterization of dualizable objects in CoCorr_S .

Proposition 4.3.4. *Let X be an S -Artin v-stack, with smooth-locally nice structure map $\pi: X \rightarrow S$, and let $A \in D_{\acute{e}t}(X, \Lambda)$. The following are equivalent.*

1. *The object (X, A) is dualizable in CoCorr_S .*
2. *The natural map $m: \mathbf{D}_{X/S} A \boxtimes_S A \rightarrow \underline{\mathrm{RHom}}(\mathrm{pr}_1^* A, \mathrm{pr}_2^! A)$ (see proof for construction) is an isomorphism.*

In this situation, the dual of (X, A) is $(X, \mathbf{D}_{X/S} A)$.

Proof. Let $\mathfrak{X} = (X, A)$, and let $\mathfrak{X}' = (X, \mathbf{D}_{X/S} A)$. There is a morphism $\mathbf{e}: \mathfrak{X}' \otimes \mathfrak{X} \rightarrow 1_{\mathrm{CoCorr}_S}$, defined as the pair (c, u) , where $c = (\Delta_{X/S}, \pi)$ and u is the composition

$$\Delta_{X/S}^*(\mathbf{D}_{X/S} A \boxtimes_S A) \xrightarrow{\sim} \mathbf{D}_{X/S} A \otimes A \rightarrow K_{X/S} = \pi^! \Lambda_S.$$

Then \mathbf{e} induces a morphism $\mathfrak{X}' \otimes \mathfrak{Y} \rightarrow \underline{\mathrm{Hom}}(\mathfrak{X}, \mathfrak{Y})$ for any object \mathfrak{Y} of CoCorr_S . For $\mathfrak{Y} = \mathfrak{X}$, the map u becomes the map m in (2).

Suppose \mathfrak{X} is dualizable, with witnesses \mathfrak{X}^\vee , $\mathrm{ev}_{\mathfrak{X}}$, and $\mathrm{coev}_{\mathfrak{X}}$. Then $\mathfrak{X}^\vee \otimes \mathfrak{Y} \rightarrow \underline{\mathrm{Hom}}(\mathfrak{X}, \mathfrak{Y})$ is an isomorphism for all objects \mathfrak{Y} . Setting $\mathfrak{Y} = 1_{\mathrm{CoCorr}_S}$, we find an isomorphism $\mathfrak{X}^\vee \cong \mathfrak{X}'$ which identifies $\mathrm{ev}_{\mathfrak{X}}$ with \mathbf{e} . Setting $\mathfrak{Y} = \mathfrak{X}$, we find that m is an isomorphism.

Conversely, if m is an isomorphism, let $\mathrm{coev}_{\mathfrak{X}} = (d, w)$, where $d = (\pi, \Delta_{X/S})$ and w is the composition

$$\pi^* \Lambda_S \cong \Lambda_X \xrightarrow{\epsilon} \underline{\mathrm{RHom}}_\Lambda(A, A) \xrightarrow{(P5)} \Delta_{X/S}^! \underline{\mathrm{RHom}}_\Lambda(\mathrm{pr}_1^* A, \mathrm{pr}_2^! A)$$

followed by $m^{-1}: \Delta_{X/S}^! \underline{\mathrm{RHom}}_\Lambda(\mathrm{pr}_1^* A, \mathrm{pr}_2^! A) \rightarrow \Delta_{X/S}^!(A \boxtimes_S \mathbf{D}_{X/S} A)$. Here ϵ is adjoint to $\mathrm{id}_A: A \rightarrow A$. A diagram chase now shows that $\mathrm{coev}_{\mathfrak{X}}$ and $\mathrm{ev}_{\mathfrak{X}}$ witness the dualizability of \mathfrak{X} . \square

In the scheme setting, a pair (X, A) is dualizable if and only if A is locally acyclic over S [LZ20, Theorem 2.16], under some mild assumptions. Similarly, if $X \rightarrow S$ is a nice morphism of v-stacks, then (X, A) is dualizable if and only if A is universally locally acyclic over S [FS21, Theorem IV.2.24]. That result extends to the situation of stacky morphisms. If $X \rightarrow S$ is smooth-locally nice, with uniformization $U \rightarrow X$, then (X, A) is dualizable if and only if $A|_U$ is ULA over S .

If \mathfrak{X} is a dualizable object of CoCorr_S , and $f: \mathfrak{X} \rightarrow \mathfrak{X}$ is an endomorphism, we may define the categorical trace $\mathrm{tr}(f)$, an object of $\Omega \mathrm{CoCorr}_S = \mathrm{End} 1_{\mathrm{CoCorr}_S}$. Let us make this explicit. The category $\Omega \mathrm{CoCorr}_S$ has objects (X, ω) , where X is an S -Artin v-stack and $\omega \in H^0(X, K_{X/S})$. A morphism $(X, \omega) \rightarrow (X', \omega')$ is a diagram

$$\begin{array}{ccc} X & \xrightarrow{p} & X' \\ & \searrow \pi & \swarrow \pi' \\ & & S \end{array}$$

with p proper, such that $\omega' = p_*\omega$. Here

$$p_* : H^0(X, K_{X/S}) \rightarrow H^0(X', K_{X'/S})$$

is induced from $\pi_* K_{X/S} \cong \pi'_* p_* p^! K_{X'/S} \cong \pi'_* p_! p^! K_{X'/S} \xrightarrow{\text{counit}} \pi'_* K_{X'/S}$.

Now let $\mathfrak{X} = (X, A)$ be a dualizable object and let $\mathfrak{f} = (c, u) : \mathfrak{X} \rightarrow \mathfrak{X}$ be an endomorphism, with $c = (c_1, c_2) : C \rightarrow X \times_S X$ and $u : c_1^* A \rightarrow c_2^* A$. By definition, $\text{tr}(\mathfrak{f}) = \text{ev}_{\mathfrak{X}} \circ (f \otimes \text{id}_{X^\vee}) \circ \text{coev}_{\mathfrak{X}}$. Let us make explicit $\text{tr}(\mathfrak{f})$. It is represented by a pair $(\text{tr}(c), \text{tr}(u))$. Here $\text{tr}(c) \in \Omega_{1_{\text{Corr}_S}}$ is the correspondence $\text{Fix}(c) \rightarrow S \times_S S = S$, where $\text{Fix}(c)$ is the fixed-point locus of the correspondence c , as in the cartesian diagram:

$$\begin{array}{ccc} \text{Fix}(c) & \xrightarrow{c'} & X \\ \Delta'_{X/S} \downarrow & & \downarrow \Delta_{X/S} \\ C & \xrightarrow{c} & X \times_S X \end{array}$$

For its part, the element $\text{tr}(u)$ is an element of $H^0(\text{Fix}(c), K_{\text{Fix}(c)/S})$. It is the image of $u \in \text{Hom}(c_1^* A, c_2^* A)$ under

$$\begin{aligned} H^0(C, \underline{\text{RHom}}(c_1^* A, c_2^* A)) & \stackrel{(P5)}{\sim} H^0(C, c^! \underline{\text{RHom}}(\text{pr}_1^* A, \text{pr}_2^* A)) \\ & \stackrel{(4.0.1)}{\sim} H^0(C, c^! (\mathbf{D}_{X/S} A \boxtimes_S A)) \\ & \xrightarrow{\alpha} H^0(C, c^! (\Delta_{X/S})_* (\mathbf{D}_{X/S} A \otimes A)) \\ & \xrightarrow{\text{ev}_A} H^0(C, c^! (\Delta_{X/S})_* K_{X/S}) \\ & \stackrel{(BC2)}{\sim} H^0(C, (\Delta'_{X/S})_* (c')^! K_{X/S}) \\ & \cong H^0(\text{Fix}(c), K_{\text{Fix}(c)/S}). \end{aligned}$$

Here, the map labeled α is adjoint to $(\Delta_{X/S})^* (\mathbf{D}_{X/S} A \boxtimes_S A) \xrightarrow{\sim} \mathbf{D}_{X/S} A \otimes A$.

Definition 4.3.5 (Inertia stack, characteristic class). In the special case that $\mathfrak{f} = \text{id}_{\mathfrak{X}} = (\Delta_{X/S}, \text{id}_A)$, the object $\text{tr}(\Delta_{X/S}) = X \times_{X \times_S X} X$ is the *inertia stack* of X , which we notate as $\text{In}_S(X)$. Its objects are pairs (x, g) , where $x \in X$ and $g \in \text{Aut } x$. Then $\text{tr}(\text{id}_A)$ is an element of $H^0(\text{In}_S(X), K_{\text{In}_S(X)})$, which we call the *characteristic class* of \mathfrak{X} . We notate this as $\text{cc}_{X/S}(A)$.

We record one lemma here for later reference.

Lemma 4.3.6. *Let $i : U \rightarrow X$ be an open immersion of S -Artin v -stacks. Then $\text{In}_S(i) : \text{In}_S(U) \rightarrow \text{In}_S(X)$ is also an open immersion. If $A \in D_{\text{ét}}(X, \Lambda)$ is ULA over S , then so is $i^* A$, and then*

$$\text{cc}_{U/S}(i^* A) = \text{In}_S(i)^* \text{cc}_{X/S}(A).$$

Proof. All constructions are local on X . □

Theorem 4.3.7 (Relative Lefschetz-Verdier trace formula). *Let $\mathfrak{X} = (X, A)$ be a dualizable object, and let $\mathfrak{f} \in \text{End } \mathfrak{X}$ lie over the correspondence $c: C \rightarrow X \times_S X$. Suppose we are given a diagram*

$$\begin{array}{ccccc} X & \longleftarrow & C & \longrightarrow & X \\ q \downarrow & & p \downarrow & & q \downarrow \\ X' & \longleftarrow & C' & \longrightarrow & X' \end{array}$$

with p, q proper. Then $\mathfrak{X}' = (X', q_ A)$ is also dualizable. Let $\mathfrak{q}: \mathfrak{X} \rightarrow \mathfrak{X}'$ be the evident 1-morphism lying over q . There is a unique morphism $\mathfrak{f}' \in \text{End } \mathfrak{X}'$ lying over $C' \rightarrow X' \times_S X'$, such that p defines a 2-morphism \tilde{p} filling in the square*

$$\begin{array}{ccc} \mathfrak{X} & \xrightarrow{\mathfrak{f}} & \mathfrak{X} \\ q \downarrow & & \downarrow q \\ \mathfrak{X}' & \xrightarrow{\mathfrak{f}'} & \mathfrak{X}' \end{array}$$

Finally, there exists a (necessarily unique) 2-morphism $\text{tr}(\tilde{p}): \text{tr}(\mathfrak{f}) \implies \text{tr}(\mathfrak{f}')$ lying over $\text{tr}(p): \text{tr}(c) \implies \text{tr}(c')$.

Proof. This is formally the same as the proof of [LZ20, Theorem], so we give a brief sketch. (In [LZ20] one gets a statement about the more general Lefschetz-Verdier pairing, which we also could have established.) The existence and uniqueness of $\mathfrak{f}' \in \text{End } \mathfrak{X}'$ follows from the definition of 2-morphisms in CoCorr_S . The dualizability of \mathfrak{X}' is [LZ20, Proposition 2.23]. The dual of \mathfrak{X}' is $(\mathfrak{X}')^\vee = (X', q_* \mathbf{D}_{X/S} A)$; there is another natural map $\mathfrak{q}^\vee: \mathfrak{X}^\vee \rightarrow (\mathfrak{X}')^\vee$ defined similarly to \mathfrak{q} .

Consider the diagram:

$$\begin{array}{ccccc} 1_{\text{CoCorr}_S} & \xrightarrow{\text{coev}_{\mathfrak{X}}} & \mathfrak{X} \otimes \mathfrak{X}^\vee & \xrightarrow{\mathfrak{f} \otimes \text{id}_{\mathfrak{X}^\vee}} & \mathfrak{X} \otimes \mathfrak{X}^\vee & \xrightarrow{\text{ev}_{\mathfrak{X}}} & 1_{\text{CoCorr}_S} \\ & \searrow \text{coev}_{\mathfrak{X}'} & \downarrow q \otimes q^\vee & & \downarrow q \otimes q^\vee & & \\ & & \mathfrak{X}' \otimes (\mathfrak{X}')^\vee & \xrightarrow{\mathfrak{f}' \otimes \text{id}_{(\mathfrak{X}')^\vee}} & \mathfrak{X}' \otimes (\mathfrak{X}')^\vee & \xrightarrow{\text{ev}_{\mathfrak{X}'}} & 1_{\text{CoCorr}_S} \end{array}$$

The two outer triangles can be filled in with a 2-morphism, as can the inner square (via $p \otimes \text{id}_{q^\vee}$). See [LZ20, Construction 1.7] for details. Composing, we find the required 2-morphism $\text{tr}(\tilde{p}): \text{tr}(\mathfrak{f}) \implies \text{tr}(\mathfrak{f}')$. \square

Corollary 4.3.8. *Let (X, A) be a dualizable object, and let $p: X \rightarrow X'$ be proper. Then $(X', p_! A)$ is dualizable. The morphism $\text{In}_S(p): \text{In}_S(X) \rightarrow \text{In}_S(X')$ is proper, and*

$$\text{In}_S(p)_* \text{cc}_{X/S}(A) = \text{cc}_{X'/S}(p_! A).$$

We can immediately deduce a familiar-looking trace formula from Theorem 4.3.7 in the case of proper diamonds. Suppose $S = \text{Spd } C$ for an algebraically closed perfectoid field C , and suppose $q: X \rightarrow S$ is a nice diamond. Let $A \in D_{\text{ét}}(X, \Lambda)$ be ULA over S . Then $\mathfrak{X} = (X, A)$ is a dualizable object of CoCorr_S . Let $\mathfrak{f} = (c, u)$ be an endomorphism of \mathfrak{X} lying over a correspondence $c: Y \rightarrow X \times_S X$. The categorical trace $\text{tr}(\mathfrak{f})$ is an endomorphism of 1_{CoCorr_S} consisting of the pair $(\text{Fix}(c), \omega)$,

where $\text{Fix}(c) = Y \times_{c, X \times_S X, \Delta_{X/S}} X$ is the fixed point locus of the correspondence, and ω is a global section of $K_{\text{Fix}(c)/S}$.

Now suppose that X and Y are both proper over S . In the setting of Theorem 4.3.7, we put $\mathfrak{X}' = (S, q_*A)$ and $C' = S$. The dualizability of \mathfrak{X}' means that $q_*A = R\Gamma(X, A)$ is a perfect complex. The morphism $f': \mathfrak{X}' \rightarrow \mathfrak{X}'$ supplied by Theorem 4.3.7 is the endomorphism $q_*(f): q_*A \rightarrow q_*A$. Finally, the existence of $\text{tr}(\tilde{p}): \text{tr}(f) \implies \text{tr}(f')$ lying over $\text{tr}(p): \text{tr}(c) \implies \text{tr}(\text{id}_S)$ implies that

$$\text{tr} \left(q_*(f) \mid R\Gamma(X, A) \right) = \int_{\text{Fix}(c)} \omega. \quad (4.3.5)$$

Definition 4.3.9. Let $x \in \text{Fix}(c)$ be an isolated point, such that $x \rightarrow S$ an isomorphism. The *local term* $\text{loc}_x(f)$ is the restriction of ω to x , considered as an element of Λ .

If f arises from an automorphism $g: X \rightarrow X$ along with a morphism $u: g^*A \rightarrow A$, we write the local term as $\text{loc}_x(g, A)$ (the dependence on u being implicit).

In the latter situation, if it so happens that $\text{Fix}(g)$ consists of finitely many isolated S -points x_1, \dots, x_n , then (4.3.5) reduces to

$$\text{tr} \left(q_*(g) \mid R\Gamma(X, A) \right) = \sum_{i=1}^n \text{loc}_{x_i}(g, A).$$

4.4 Examples

In the following, assume $S = \text{Spd } C$ for an algebraically closed perfectoid field C , or else $S = \text{Spd } k$ for an algebraically closed discrete field of characteristic p . In both cases, we freely identify $D_{\text{ét}}(S, \Lambda)$ with the derived category of Λ -modules.

Example 4.4.1. An object $A \in D_{\text{ét}}(S, \Lambda)$ is ULA over S if and only if (when considered as a complex of Λ -modules) A is perfect. (See Lemma B.1.2 in the Appendix for a proof of this claim and related conditions.)

Example 4.4.2. Let T be a locally profinite set, and let $T_S = \underline{T} \times S$ be the associated constant diamond over S . Then $f: T_S \rightarrow S$ is nice. Let $C(T, \Lambda)$ be the ring of continuous functions $T \rightarrow \Lambda$, for the discrete topology on Λ . We may naturally identify $D_{\text{ét}}(T_S, \Lambda)$ with the derived category of the abelian category of smooth $C(T, \Lambda)$ -modules in the sense of §B.2. Indeed, $D_{\text{ét}}(T_S, \Lambda) \cong D(T_{S, \text{ét}}, \Lambda)$ since T_S locally has cohomological dimension zero, so it is enough to identify $\text{Sh}(T_{S, \text{ét}}, \Lambda)$ with the category of smooth $C(T, \Lambda)$ -modules, which is Lemma B.2.5. This identification matches the constant sheaf Λ with the smooth $C(T, \Lambda)$ -module $C_c(T, \Lambda)$. Under this identification, we have concrete descriptions of the four operations associated with f (where we freely use some language and notation from §B.2):

- f^* sends a Λ -module M to the smooth $C(T, \Lambda)$ -module $C_c(T, \Lambda) \otimes_{\Lambda} M$,
- f_* sends a smooth $C(T, \Lambda)$ -module M to M^c regarded as a Λ -module.
- $f_!$ sends a smooth $C(T, \Lambda)$ -module M to its underlying Λ -module.

- $f^!$ sends a smooth Λ -module M to $\mathrm{RHom}_\Lambda(C_c(T, \Lambda), M)^s$. In particular $H^0(T_S, f^! \Lambda_S) \cong \mathrm{Dist}(T, \Lambda)$, the module of Λ -valued distributions on T .

It will not generally be true that Λ_{T_S} is ULA over S , unless T is discrete.

Example 4.4.3. Suppose G is a locally pro- p group. Let $[S/G_S]$ be the classifying v-stack of G_S -torsors. We assume that there is a free action of G_S on a cohomologically smooth diamond $X \rightarrow S$. In that case $[S/G_S]$ has a cohomologically smooth uniformization by the cohomologically smooth diamond X/G_S ; thus $[S/G_S]$ is a smooth S -Artin v-stack. Then $K_{[S/G_S]/S}$ is the invertible sheaf associated with the module $\mathrm{Haar}(G, \Lambda)$ of Λ -valued Haar measures on G .

We may identify $D_{\text{ét}}([S/G_S], \Lambda)$ with the derived category of Λ -modules with a smooth G -action [FS21, Theorem V.1.1]. If π is a smooth representation of G on a Λ -module, let $[\pi]$ denote the corresponding object of $D_{\text{ét}}([S/G_S], \Lambda)$. The object $[\pi]$ is ULA over S if and only if π is admissible, by which we mean the condition that the derived K -invariants of π are a perfect complex for all compact open $K \subset G$. In that case, $\mathbf{D}_{[S/G_S]/S}[\pi] \cong [\pi^\vee] \otimes K_{[S/G_S]/S}$ is the smooth dual twisted by the module of Haar measures.

Example 4.4.4. Let G be a locally pro- p group as in Example 4.4.3. Let T be a locally profinite set equipped with a continuous action of G , and let T_S be the constant diamond over S . The stacky quotient $[T_S/G_S]$ is an S -Artin v-stack; it is uniformized by $(X \times_S T_S)/G_S$. It is not in general cohomologically smooth over S . The category $D_{\text{ét}}([T_S/G_S], \Lambda)$ may be identified with the derived category of G -equivariant smooth $C(T, \Lambda)$ -modules. Using this identification, we get a natural isomorphism $H^0([T_S/G_S], K_{[T_S/G_S]/S}) \cong \mathrm{Dist}(T, \mathrm{Haar}(G, \Lambda))^G$, the module of G -invariant distributions on T with values in $\mathrm{Haar}(G, \Lambda)$. If G is unimodular, and if we choose a Haar measure on G , then this is isomorphic to $\mathrm{Dist}(T, \Lambda)^G$.

A stack of this form appears as the inertia stack of $[S/G_S]$. Indeed, $\mathrm{In}_S([S/G_S]) \cong [G_S // G_S]$ is the stack of conjugacy classes of G . (Reasoning: For a perfectoid space $Y \rightarrow S$, a Y -point of $\mathrm{In}_S([S/G_S])$ is a G_S -torsor $\tilde{Y} \rightarrow Y$ together with a G_S -equivariant automorphism $i: \tilde{Y} \rightarrow \tilde{Y}$. Such an automorphism arises as $i(y) = f(y).y$, where $f: \tilde{Y} \rightarrow G_S$ is a morphism satisfying $f(gy) = gf(y)g^{-1}$. The pair (\tilde{Y}, f) then constitutes a Y -point of $[G_S // G_S]$.)

Suppose now that G is unimodular, and suppose we have chosen a Haar measure on it. Let π be a smooth admissible representation of G on a Λ -module. Then $\mathrm{cc}_{[S/G_S]/S}([\pi])$ is an element of

$$H^0([G_S // G_S], K_{[G_S // G_S]/S}) \cong \mathrm{Dist}(G, \Lambda)^G,$$

the module of conjugacy-invariant Λ -valued distributions on G . This element is none other than the trace distribution of π .

4.5 A Künneth theorem for characteristic classes

The goal of this section is to prove the compatibility of the categorical trace with fiber products, to get an analogue of the relation $\mathrm{tr}(A \otimes B) = \mathrm{tr}(A) \mathrm{tr}(B)$ for square matrices.

As an example of what we will do, suppose $X_1, X_2 \rightarrow S$ are two smooth-locally nice morphisms of S -Artin v-stacks, and suppose $A_i \in D_{\text{ét}}(X_i, \Lambda)$ is ULA over S for $i = 1, 2$. Then $A_1 \boxtimes_S A_2$ is ULA over S , so we may define the characteristic class $\mathrm{cc}_{X_1 \times_S X_2/S}(A_1 \boxtimes_S A_2)$ in $H^0(\mathrm{In}_S(X_1 \times_S X_2))$.

X_2), $K_{\text{In}_S(X_1 \times_S X_2)}$). We have an isomorphism $\text{In}_S(X_1 \times_S X_2) \cong \text{In}_S(X_1) \times_S \text{In}_S(X_2)$, and therefore a Künneth map

$$\kappa_S: K_{\text{In}_S(X_1)/S} \otimes K_{\text{In}_S(X_2)/S} \rightarrow K_{\text{In}_S(X_1 \times_S X_2)/S}$$

which we notate as $\mu_1 \otimes \mu_2 \mapsto \mu_1 \boxtimes_S \mu_2$ for global sections μ_i of $K_{\text{In}_S(X_i)/S}$. Then it is straightforward to show (and a corollary of Theorem 4.5.3 below) that

$$\text{cc}_{X_1/S}(A_1) \boxtimes_S \text{cc}_{X_2/S}(A_2) = \text{cc}_{X_1 \times_S X_2/S}(A_1 \boxtimes_S A_2).$$

For our applications we need a more general result involving fiber products over bases other than S . First, we need a modification of the above Künneth map in a general setting.

Definition 4.5.1 (Modified Künneth map). Let $U \rightarrow T$ be a cohomologically smooth morphism of S -Artin v-stacks. Suppose we are given a commutative diagram of S -Artin v-stacks

$$\begin{array}{ccc} Y_i & \xrightarrow{f_i} & X_i \\ g_i \downarrow & & \downarrow \\ U & \longrightarrow & T \end{array}$$

for $i = 1, 2$. Let $f: Y_1 \times_U Y_2 \rightarrow X_1 \times_T X_2$ and $g: Y_1 \times_U Y_2 \rightarrow U$ be the induced product maps. Let A_i be an object of $D_{\text{ét}}(X_i, \Lambda)$ for $i = 1, 2$. We define a map

$$\kappa_{U/T}: f_1^! A_1 \boxtimes_U f_2^! A_2 \rightarrow f^!(A_1 \boxtimes_T A_2) \otimes g^* K_{U/T}$$

as follows. There is a cartesian diagram

$$\begin{array}{ccc} Y_1 \times_U Y_2 & \xrightarrow{\Delta'_{U/T}} & Y_1 \times_T Y_2 \\ g \downarrow & & \downarrow g_1 \times_T g_2 \\ U & \xrightarrow{\Delta_{U/T}} & U \times_T U \end{array}$$

The map $\kappa_{U/T}$ is defined as the composition

$$\begin{aligned} & f_1^! A_1 \boxtimes_U f_2^! A_2 \\ \cong & (\Delta'_{U/T})^*(f_1^! A_1 \boxtimes_T f_2^! A_2) \\ \rightarrow & (\Delta'_{U/T})^*(f_1 \times_T f_2)^!(A_1 \boxtimes_T A_2) \\ \stackrel{(4.1.3)}{\cong} & (\Delta'_{U/T})^*(f_1 \times_T f_2)^!(A_1 \boxtimes_T A_2) \otimes g^*(\Delta_{U/T})^! \Lambda_T \otimes g^* K_{U/T} \\ \rightarrow & (\Delta'_{U/T})^*(f_1 \times_T f_2)^!(A_1 \boxtimes_T A_2) \otimes (\Delta'_{U/T})^! \Lambda_{Y_1 \times_T Y_2} \otimes g^* K_{U/T} \\ \rightarrow & (\Delta'_{U/T})^!(f_1 \times_T f_2)^!(A_1 \boxtimes_T A_2) \otimes g^* K_{U/T} \\ \cong & f^!(A_1 \boxtimes_T A_2) \otimes g^* K_{U/T}. \end{aligned}$$

In particular the case $X_1 = X_2 = T = S$ yields a map

$$\kappa_{U/S}: K_{Y_1/S} \boxtimes_U K_{Y_2/S} \rightarrow K_{Y_1 \times_U Y_2/S} \otimes g^* K_{U/S}. \quad (4.5.1)$$

We can now introduce the setup of the main theorem of this section. We consider bases $T \rightarrow S$ satisfying two hypotheses: (1) $T \rightarrow S$ is cohomologically smooth, and (2) $\Delta_{T/S}: T \rightarrow T \times_S T$ is cohomologically smooth. These are satisfied for instance when $T = [S/G]$, where G is a cohomologically smooth group diamond over S . Considering the diagram

$$\begin{array}{ccccc} \mathrm{In}_S(T) & \longrightarrow & T & & \\ \downarrow & & \downarrow \Delta_{T/S} & & \\ T & \xrightarrow{\Delta_{T/S}} & T \times_S T & \longrightarrow & T \\ & & \downarrow & & \downarrow \\ & & T & \longrightarrow & S \end{array}$$

in which both squares are cartesian, we see that $\mathrm{In}_S(T) \rightarrow S$ is also cohomologically smooth. Furthermore we can trivialize the dualizing complex $K_{\mathrm{In}_S(T)/S}$.

Lemma 4.5.2. *Let T be an S -Artin v -stack such that the structure map $\pi: T \rightarrow S$ and diagonal $\Delta_{T/S}: T \rightarrow T \times_S T$ are both cohomologically smooth. Then the object $\mathfrak{T} = (T, \Lambda_T) \in \mathrm{CoCorr}_S$ is dualizable, and its characteristic class $\mathrm{cc}_{T/S}(\Lambda_T)$, considered as a morphism $\Lambda_{\mathrm{In}_S(T)} \rightarrow K_{\mathrm{In}_S(T)/S}$, is an isomorphism.*

In the case $T = [S/G]$, where G is a cohomologically smooth group diamond over S , the inertia stack is $\mathrm{In}_S(T) = [G // G]$, the stack of conjugacy classes of G . This is a cohomologically smooth stack of dimension 0, so perhaps it is unsurprising that it has trivial dualizing complex; the lemma states that the trivialization is in fact canonical.

Proof. Let $\mathrm{pr}_1, \mathrm{pr}_2: T \times_S T \rightarrow T$ be the projection morphisms. The morphism in $D_{\mathrm{ét}}(T, \Lambda)$ associated with the cohomological correspondence $\mathrm{coev}_{\mathfrak{T}}: (S, \Lambda_S) \rightarrow (T \times_S T, \mathrm{pr}_2^* K_{T/S})$ is an isomorphism:

$$\pi^* \Lambda_S \cong \Delta_{T/S}^! \mathrm{pr}_1^! \pi^* \Lambda_S \stackrel{\alpha}{\cong} \Delta_{T/S}^! \mathrm{pr}_2^* \pi^! \Lambda_S = \Delta_{T/S}^! \mathrm{pr}_2^* K_{T/S}.$$

Here we use cohomological smoothness of π to get the isomorphism α , as in Theorem 4.1.4. The morphism $\Delta_{T/S}^* \mathrm{pr}_2^* K_{T/S} \rightarrow \pi^! \Lambda_S$ associated with $\mathrm{ev}_{\mathfrak{T}}: (T \times_S T, \mathrm{pr}_2^* K_{T/S}) \rightarrow (S, \Lambda_S)$ is also an isomorphism (this is true without any hypotheses on π or $\Delta_{T/S}$). Referring to the diagram

$$\begin{array}{ccccc} & & \mathrm{In}_S(T) & & \\ & & \swarrow h & \searrow h & \\ & T & & & T \\ \swarrow \pi & & \Delta_{T/S} & \Delta_{T/S} & \searrow \pi \\ S & & T \times_S T & & S \end{array}$$

we may describe the characteristic class $\text{cc}_{T/S}(\Lambda_T)$ as the composition

$$\begin{aligned} h^* \pi^* \Lambda_S &\cong h^* \Delta_{T/S}^! \text{pr}_2^* K_{T/S} \\ &\stackrel{\beta}{\cong} h^! \Delta_{T/S}^* \text{pr}_2^* K_{T/S} \cong h^! K_{T/S} \cong K_{\text{In}_S(T)/S}, \end{aligned}$$

which is an isomorphism. Here we once again applied Theorem 4.1.4, using the cohomological smoothness of $\Delta_{T/S}$. \square

Now suppose $p_i: X_i \rightarrow T$ ($i = 1, 2$) is a morphism of S -Artin v-stacks, with fiber product $p: X_1 \times_T X_2 \rightarrow T$. There is an isomorphism

$$\text{In}_S(X_1 \times_T X_2) \cong \text{In}_S(X_1) \times_{\text{In}_S(T)} \text{In}_S(X_2),$$

and therefore a modified Künneth map

$$\kappa_{\text{In}_S(T)/S}: K_{\text{In}_S(X_1)/S} \boxtimes_{\text{In}_S(T)} K_{\text{In}_S(X_2)/S} \rightarrow K_{\text{In}_S(X_1 \times_T X_2)/S} \otimes \text{In}(p)^* K_{\text{In}_S(T)/S}.$$

Using the trivialization $\Lambda_{\text{In}_S(T)/S} \cong K_{\text{In}_S(T)/S}$ from Lemma 4.5.2, we obtain a map

$$K_{\text{In}_S(X_1)/S} \boxtimes_{\text{In}_S(T)} K_{\text{In}_S(X_2)/S} \rightarrow K_{\text{In}_S(X_1 \times_T X_2)/S}, \quad (4.5.2)$$

which on global sections we notate as $\mu_1 \otimes \mu_2 \mapsto \mu_1 \boxtimes_{\text{In}_S(T)} \mu_2$.

Finally we can state the main theorem of the section.

Theorem 4.5.3. *Let $X_1, X_2 \rightarrow T$ be two morphisms of S -Artin v-stacks, and let $A_i \in D_{\text{ét}}(X_i, \Lambda)$ be ULA over S . Assume that the structure map $\pi: T \rightarrow S$ and the diagonal $\Delta_{T/S}: T \rightarrow T \times_S T$ are both cohomologically smooth. Then $A_1 \boxtimes_T A_2$ is ULA over S , and*

$$\text{cc}_{X_1/S}(A_1) \boxtimes_{\text{In}_S(T)} \text{cc}_{X_2/S}(A_2) = \text{cc}_{X_1 \boxtimes_T X_2/S}(A_1 \boxtimes_T A_2).$$

In order to prove Theorem 4.5.3, we need to enhance the category CoCorr_S to include data coming from a smooth base v-stack, which we allow to vary. Such a category would have objects $(X \rightarrow T, A)$, where $T \rightarrow S$ is cohomologically smooth, $X \rightarrow T$ is a morphism of v-stacks, and $A \in D_{\text{ét}}(X, \Lambda)$. The morphisms $(X \rightarrow T, A) \rightarrow (X' \rightarrow T', A')$ would be pairs (q^\natural, α) , where $q^\natural = (p, q, p')$ is a morphism of correspondences:

$$\begin{array}{ccccc} X & \xleftarrow{c} & C & \xrightarrow{c} & X' \\ p \downarrow & & q \downarrow & & \downarrow p' \\ T & \xleftarrow{u} & U & \xrightarrow{u'} & T' \end{array} \quad (4.5.3)$$

and $\alpha: c^* A \rightarrow (c')^! A'$ is a morphism. We assume that u' is cohomologically smooth.

Given pairs $\mathfrak{X}_1 = (X_1 \rightarrow T, A_1)$ and $\mathfrak{X}_2 = (X_2 \rightarrow T, A_2)$ with common base T , we could then define $\mathfrak{X}_1 \boxtimes_T \mathfrak{X}_2 = (X_1 \times_T X_2 \rightarrow T, A_1 \boxtimes_T A_2)$. However, it turns out that this definition is not functorial in \mathfrak{X}_1 and \mathfrak{X}_2 . That is: given a morphism $e: T \rightarrow T'$ in Corr_S and morphisms $f_i: \mathfrak{X}_i \rightarrow \mathfrak{X}'_i$ lying over e , one cannot in general define a map $f_1 \otimes_e f_2: \mathfrak{X}_1 \boxtimes_T \mathfrak{X}_2 \rightarrow \mathfrak{X}'_1 \boxtimes_{T'} \mathfrak{X}'_2$. Essentially, this is due to the appearance of the invertible sheaf $K_{U/S}$ appearing in the modified Künneth map (4.5.1).

To obtain a functorial definition of $\mathfrak{X}_1 \boxtimes_T \mathfrak{X}_2$, we need to define an enhancement of the category which keeps track of an invertible sheaf living on the base.

Definition 4.5.4 (The category of based cohomological correspondences). We define a symmetric monoidal 2-category BCoCorr_S . The objects of BCoCorr_S are triples $(X \rightarrow T, A, B)$, where $X \rightarrow T$ is a morphism of S -Artin v-stacks, both smooth-locally nice over S , where A is an object of $D_{\text{ét}}(X, \Lambda)$, and where B is an *invertible* object of $D_{\text{ét}}(T, \Lambda)$. We assume that the structure map $T \rightarrow S$ be cohomologically smooth. Given objects $\mathfrak{X} = (X \rightarrow T, A, B)$ and $\mathfrak{X}' = (X' \rightarrow T', A', B')$, an object of the category $\text{Hom}_{\text{BCoCorr}_S}(\mathfrak{X}, \mathfrak{X}')$ is a triple $(q^\natural, \alpha, \beta)$. The first element in the triple is a morphism $q^\natural = (p, q, p')$ between correspondences as in (4.5.3), where u' is cohomologically smooth. The second element is a morphism $\alpha: c^*A \rightarrow (c')^!A'$, and the third is an *isomorphism* $\beta: u^*B \xrightarrow{\sim} (u')^*B'$. Compositions of morphisms are defined similarly as in Definition 4.3.3. (Note that the cohomological smoothness of u' is preserved under composition of correspondences.) Given objects $(X \rightarrow T, A, B)$ and $(X' \rightarrow T', B')$, and morphisms between them represented by $C \rightarrow X \times_U X'$ (lying over $U \rightarrow T \times_S T'$) and $D \rightarrow X \times_V X'$ (lying over $V \rightarrow T \times_S T'$), a 2-morphism is a pair of proper morphisms $C \rightarrow D$ and $U \rightarrow V$ making the relevant diagrams commute.

The monoidal structure on BCoCorr_S is defined by

$$(X \rightarrow T, A, B) \otimes (X' \rightarrow T', A', B') = (X \times_S X' \rightarrow T \times_S T', A \boxtimes_S A', B \boxtimes_S B').$$

The unit object is (S, S, Λ_S) .

Finally, we introduce the obvious monoidal functors $\mathcal{B}, \mathcal{S}: \text{BCoCorr}_S \rightarrow \text{CoCorr}_S$, with $\mathcal{B}(X \rightarrow T, A, B) = (T, B)$ (the *base*) and $\mathcal{S}(X \rightarrow T, A, B) = (X, A)$ (the *source*).

So far the objects A and B in Definition 4.5.4 have nothing to do with each other. They only begin to interact when we talk about fiber products of objects of BCoCorr_S over common bases. Given objects $\mathfrak{X}_i = (X_i \xrightarrow{p_i} T, A_i, B)$ for $i = 1, 2$ with common base $\mathfrak{T} = (T, B)$, we define

$$\mathfrak{X}_1 \boxtimes_{\mathfrak{T}} \mathfrak{X}_2 = (X_1 \times_T X_2 \xrightarrow{p} T, (A_1 \boxtimes_T A_2) \otimes p^*B^{-1}).$$

We claim that $\boxtimes_{\mathfrak{T}}$ defines a monoidal functor

$$\text{BCoCorr}_S \times_{\mathcal{B}, \text{CoCorr}_S} \text{BCoCorr}_S \rightarrow \text{BCoCorr}_S. \quad (4.5.4)$$

A morphism in the category $\text{BCoCorr}_S \times_{\mathcal{B}, \text{CoCorr}_S} \text{BCoCorr}_S$ is a morphism $e: \mathfrak{T} \rightarrow \mathfrak{T}'$ in CoCorr_S together with a pair of morphisms $f_i: \mathfrak{X}_i \rightarrow \mathfrak{X}'_i$ ($i = 1, 2$) lying over e . We may represent this state of affairs with a diagram

$$\begin{array}{ccccc} X_i & \xleftarrow{c_i} & C_i & \xrightarrow{c'_i} & X'_i \\ p_i \downarrow & & q_i \downarrow & & \downarrow p'_i \\ T & \xleftarrow{u} & U & \xrightarrow{u'} & T' \end{array} \quad (4.5.5)$$

for $i = 1, 2$, with u' smooth, together with morphisms $\alpha_i: c_i^*A_i \rightarrow (c'_i)^!A'_i$ for $i = 1, 2$ and an isomorphism $\beta: u^*B \xrightarrow{\sim} (u')^*B'$. Taking fiber products over the base correspondence, we obtain a morphism of correspondences $q^\natural = (p, q, p')$ fitting into a diagram

$$\begin{array}{ccccc} X_1 \times_T X_2 & \xleftarrow{c} & C_1 \times_U C_2 & \xrightarrow{c'} & X'_1 \times_{T'} X'_2 \\ p \downarrow & & q \downarrow & & \downarrow p' \\ T & \xleftarrow{u} & U & \xrightarrow{u'} & T' \end{array} \quad (4.5.6)$$

The required morphism

$$c^* ((A_1 \boxtimes_T A_2) \otimes p^* B^{-1}) \rightarrow (c')^! ((A'_1 \boxtimes_{T'} A'_2) \otimes (p')^*(B')^{-1}) \quad (4.5.7)$$

is defined as the composition

$$\begin{aligned} c^* ((A_1 \boxtimes_T A_2) \otimes p^* B^{-1}) &\cong (c_1^* A_1 \boxtimes_U c_2^* A_2) \otimes q^* u^* B^{-1} \\ &\xrightarrow{(\alpha_1 \boxtimes_U \alpha_2) \otimes \beta} (c'_1)^! A'_1 \boxtimes_U (c'_2)^! A'_2 \otimes q^* ((u')^! B')^{-1} \\ &\xrightarrow{\kappa_{U/T'}} (c')^! (A'_1 \boxtimes_{T'} A'_2) \otimes q^* (K_{U/T'} \otimes ((u')^! B')^{-1}) \\ &\cong (c')^! (A'_1 \boxtimes_{T'} A'_2) \otimes q^* (u')^* (B')^{-1} \\ &\cong (c')^! ((A'_1 \boxtimes_{T'} A'_2) \otimes (p')^* (B')^{-1}), \end{aligned}$$

where in the last step we used Lemma 4.1.6.

To completely justify that (4.5.4) is a functor, one must also produce a 2-isomorphism

$$(f'_1 \boxtimes_{e'} f'_2) \circ (f_1 \boxtimes_e f_2) \cong (f'_1 \circ f_1) \boxtimes_{e' \circ e} (f'_2 \circ f_2) \quad (4.5.8)$$

whenever all compositions are defined. Furthermore one must also show that (4.5.4) is a monoidal functor; that is, we have an isomorphism

$$(\mathfrak{X}_1 \boxtimes_{\mathfrak{T}} \mathfrak{X}_2) \otimes (\mathfrak{X}'_1 \boxtimes_{\mathfrak{T}'} \mathfrak{X}'_2) \cong (\mathfrak{X}_1 \otimes \mathfrak{X}'_1) \boxtimes_{\mathfrak{T} \otimes \mathfrak{T}'} (\mathfrak{X}_2 \otimes \mathfrak{X}'_2). \quad (4.5.9)$$

The details are straightforward but tedious.

We can now prove Theorem 4.5.3. Let $X_1, X_2 \rightarrow T$ be two morphisms of S -Artin v -stacks, and let $A_i \in D_{\text{ét}}(X_i, \Lambda)$ be ULA over S . Assume that the structure map $\pi: T \rightarrow S$ and the diagonal $\Delta_{T/S}$ are both cohomologically smooth.

Let $\mathfrak{X}_i = (X_i \rightarrow T, A_i, \Lambda_T) \in \text{BCoCorr}_S$ for $i = 1, 2$, so that $\mathcal{B}(\mathfrak{X}_1) = \mathcal{B}(\mathfrak{X}_2) = \mathfrak{T} = (T, \Lambda_T)$. Then \mathfrak{X}_i is dualizable, with dual $\mathfrak{X}_i^\vee = (X_i \rightarrow T, \mathbf{D}_{X_i/S} A_i, K_{T/S})$, as witnessed by $\text{coev}_{\mathfrak{X}_i}: 1_{\text{BCoCorr}_S} \rightarrow \mathfrak{X}_i \otimes \mathfrak{X}_i^\vee$ and $\text{ev}_{\mathfrak{X}_i}: \mathfrak{X}_i^\vee \otimes \mathfrak{X}_i \rightarrow 1_{\text{BCoCorr}_S}$. Note that $\mathcal{B}(\text{coev}_{\mathfrak{X}_i}) = \text{coev}_{\mathfrak{T}}$ and $\mathcal{S}(\text{coev}_{\mathfrak{X}_i}) = \text{coev}_{\mathcal{S}(\mathfrak{X}_i)}$, and similarly for ev . Then the categorical trace of $1_{\mathfrak{X}_i}$ is $\text{tr}(1_{\mathfrak{X}_i}) = \text{ev}_{\mathfrak{X}_i} \circ \text{coev}_{\mathfrak{X}_i}$, so that $\mathcal{S}(\text{tr}(1_{\mathfrak{X}_i})) = \text{tr}(1_{\mathcal{S}(\mathfrak{X}_i)}) = \text{cc}_{X_i/S}(A_i)$.

Now consider $\mathfrak{X} = \mathfrak{X}_1 \boxtimes_{\mathfrak{T}} \mathfrak{X}_2 = (X_1 \boxtimes_T X_2 \rightarrow T, A_1 \boxtimes_T A_2, \Lambda_T)$. Define an object $\mathfrak{X}^\vee = \mathfrak{X}_1^\vee \boxtimes_{\mathfrak{T}^\vee} \mathfrak{X}_2^\vee$, and define morphisms $\text{coev}_{\mathfrak{X}}$ and $\text{ev}_{\mathfrak{X}}$ via the diagrams

$$\begin{array}{ccc} 1_{\text{BCoCorr}_S} & \xrightarrow{\text{coev}_{\mathfrak{X}_1} \boxtimes_{\text{coev}_{\mathfrak{T}}} \text{coev}_{\mathfrak{X}_2}} & (\mathfrak{X}_1 \otimes \mathfrak{X}_1^\vee) \boxtimes_{\mathfrak{T} \otimes \mathfrak{T}^\vee} (\mathfrak{X}_2 \otimes \mathfrak{X}_2^\vee) \\ & \searrow \text{coev}_{\mathfrak{X}} & \downarrow \cong (4.5.9) \\ & & \mathfrak{X} \otimes \mathfrak{X}^\vee \end{array}$$

and

$$\begin{array}{ccc} \mathfrak{X} \otimes \mathfrak{X}^\vee & & \\ \downarrow (4.5.9) \cong & \searrow \text{ev}_{\mathfrak{X}} & \\ (\mathfrak{X}_1^\vee \otimes \mathfrak{X}_1) \boxtimes_{\mathfrak{T} \otimes \mathfrak{T}^\vee} (\mathfrak{X}_2^\vee \otimes \mathfrak{X}_2) & \xrightarrow{\text{ev}_{\mathfrak{X}_1} \boxtimes_{\text{ev}_{\mathfrak{T}}} \text{ev}_{\mathfrak{X}_2}} & 1_{\text{BCoCorr}_S} \end{array}$$

Then \mathfrak{X}^\vee , $\text{coev}_{\mathfrak{X}}$ and $\text{ev}_{\mathfrak{X}}$ witness the dualizability of \mathfrak{X} . It follows that $\mathcal{S}(\mathfrak{X}) = (X_1 \times_T X_2, A_1 \boxtimes_T A_2)$ is dualizable, so that $A_1 \boxtimes_T A_2$ is ULA over S . Now consider $\text{tr}(1_{\mathfrak{X}}) = \text{ev}_{\mathfrak{X}} \circ \text{coev}_{\mathfrak{X}}$, an endomorphism of 1_{BCoCorr_S} . On the one hand, $\mathcal{S}(\text{tr}(1_{\mathfrak{X}})) = \text{tr}(1_{\mathcal{S}(\mathfrak{X})}) \in \text{End } 1_{\text{CoCorr}_S}$ is the data of the inertia stack $\text{In}_S(X_1 \times_T X_2)$ together with the characteristic class $\text{cc}_{X_1 \times_T X_2/S}(A) \in H^0(\text{In}_S(X_1 \times_T X_2), K_{\text{In}_S(X_1 \times_T X_2)/S})$. On the other hand, (4.5.8) gives a 2-isomorphism

$$\text{tr}(1_{\mathfrak{X}}) \cong (\text{ev}_{\mathfrak{X}_1} \boxtimes_{\text{ev}_{\mathfrak{X}_2}} \text{ev}_{\mathfrak{X}_2}) \circ (\text{coev}_{\mathfrak{X}_1} \boxtimes_{\text{coev}_{\mathfrak{X}_2}} \text{coev}_{\mathfrak{X}_2}) \cong \text{tr}(1_{\mathfrak{X}_1}) \boxtimes_{\text{tr}(1_{\mathfrak{X}_2})} \text{tr}(1_{\mathfrak{X}_2}).$$

The source of this morphism is the correspondence $\text{In}_S(X_1) \times_{\text{In}_S(T_1)} \text{In}_S(X_2) \rightarrow S \times_S S \cong S$ together with a global section of $K_{\text{In}_S(X_1 \times_T X_2)/S}$. Reviewing the definition of \boxtimes for morphisms in BCoCorr_S as in (4.5.7), we see that this section is the image of $\text{cc}_{X_1/S}(A_1) \otimes \text{cc}_{X_2/S}(A_2)$ under

$$\begin{aligned} K_{\text{In}_S(X_1)/S} \boxtimes_{\text{In}_S(T)} K_{\text{In}_S(X_2)/S} &\xrightarrow{\kappa_{\text{In}_S(T)/S}} K_{\text{In}_S(X_1 \times_T X_2)/S} \otimes \text{In}(p)^* K_{\text{In}_S(T)} \\ &\cong K_{\text{In}_S(X_1 \times_T X_2)/S}, \end{aligned}$$

where the last isomorphism is induced from the inverse to $\text{cc}_{T/S}(\Lambda_T): \Lambda_{\text{In}_S(T)} \rightarrow K_{\text{In}_S(T)/S}$. The result is exactly $\text{cc}_{X_1/S}(A_1) \boxtimes_{\text{In}_S(T)} \text{cc}_{X_2/S}(A_2)$ as defined in Theorem 4.5.3.

4.6 The case of $[X/G]$ for G smooth

Let X be a nice diamond over $S = \text{Spd}C$ which is equipped with an action of a cohomologically smooth S -group diamond G . Let $\alpha: X \times_S G \rightarrow X$ be the action map $(x, g) \mapsto g(x)$. Let $Y = [X/G]$ be the stacky quotient; this is an S -Artin v-stack which is locally nice over S .

Let $A \in D_{\text{ét}}(Y, \Lambda)$ be ULA over S . Then the pair (Y, A) is dualizable in CoCorr_S , and we obtain a characteristic class

$$\text{cc}_{Y/S}(A) \in H^0(\text{In}_S(Y), K_{\text{In}_S(Y)/S}).$$

On the other hand, the pullback A_X of A along $X \rightarrow Y$ is also ULA over S (because $G \rightarrow S$ is cohomologically smooth). For each element $g \in G(S)$, we have an isomorphism $u_g: A_X \rightarrow g^* A_X$ lying over $g: X \rightarrow X$. The pair (g, u_g) constitutes an endomorphism of the dualizable object (X, A_X) in CoCorr_S , so we may define the categorical trace $\text{tr}(g, u_g) \in H^0(\text{Fix}(g), K_{\text{Fix}(g)/S})$. Here $\text{Fix}(g) = X \times_{g, X \times_S X, \Delta_{X/S}} X$ is the fixed-point locus of g on X . The object of this section is to show how the $\text{tr}(g, u_g)$ can be derived from $\text{cc}_{Y/S}(A)$.

First we give a concrete presentation of $\text{In}_S(Y)$. Define a correspondence c on X by

$$\begin{aligned} c = \text{pr}_X \times_S \alpha: X \times_S G &\rightarrow X \times_S X \\ (x, g) &\mapsto (x, g(x)). \end{aligned}$$

Then the fixed-point locus $\text{Fix}(c) \subset X \times_S G$ is G -stable for the G -action on $X \times_S G$ given by $h(x, g) = (h(x), hgh^{-1})$, and then $\text{In}_S(Y) \cong [\text{Fix}(c)/G]$. With respect to this isomorphism, the canonical map $p: \text{In}_S(Y) \rightarrow \text{In}_S([S/G]) \cong [G // G]$ is the quotient by G of the projection map $\text{Fix}(c) \rightarrow G$.

The G -equivariance of $A|_X$ may be expressed as an isomorphism $u: A|_{X \times_S G} \rightarrow \alpha^* A|_X$. This is not a cohomological correspondence in general (as $\alpha^* \neq \alpha^!$). To obtain a cohomological correspondence

on non-stacky objects, we work over the base G . Let $X_G = X \times_S G$, and consider the correspondence \tilde{c} defined by the diagram of diamonds over G :

$$\begin{array}{ccc} & X_G & \\ \text{id}_{X_G} \swarrow & & \searrow \tilde{\alpha}: (x,g) \mapsto (g(x),g) \\ X_G & & X_G \end{array}$$

By design, the fiber of this correspondence over $g \in G(C)$ is the graph of the automorphism $g: X \rightarrow X$. Moreover, there is a natural isomorphism $\text{Fix}(c) \cong \text{Fix}(\tilde{c})$, and the fiber of $\text{Fix}(\tilde{c})$ over any $g \in G(C)$ is exactly $\text{Fix}(g)$. The G -equivariance of $A|_X$ is encoded by an isomorphism $\tilde{u}: A|_{X_G} \rightarrow \tilde{\alpha}^* A|_{X_G}$. Since $\tilde{\alpha}$ is an isomorphism, we have $\tilde{\alpha}^* \cong \tilde{\alpha}^!$, and therefore the pair (\tilde{c}, \tilde{u}) constitutes an endomorphism of the dualizable object $(X_G, A|_{X_G})$ of CoCorr_G . The categorical trace of (\tilde{c}, \tilde{u}) is an element

$$\text{tr}(\tilde{c}, \tilde{u}) \in H^0(\text{Fix}(\tilde{c}), K_{\text{Fix}(\tilde{c})/G}).$$

This is the ‘‘universal local term’’ for the action of G on X , in the sense that for any $g \in G(C)$, the restriction map

$$H^0(\text{Fix}(\tilde{c}), K_{\text{Fix}(\tilde{c})/G}) \rightarrow H^0(\text{Fix}(g), K_{\text{Fix}(g)/S})$$

carries $\text{tr}(\tilde{c}, \tilde{u})$ onto $\text{tr}(g, u_g)$.

We want to compare the characteristic class $\text{cc}_{Y/S}(A)$ with the universal local term $\text{tr}(\tilde{c}, \tilde{u})$. To do this, we first observe that from the Cartesian square

$$\begin{array}{ccc} \text{Fix}(\tilde{c}) & \longrightarrow & G \\ q \downarrow & & \downarrow \\ \text{In}_S(Y) & \longrightarrow & [G // G] \end{array}$$

we obtain a canonical map $q^* K_{\text{In}_S(Y)/[G//G]} \rightarrow K_{\text{Fix}(\tilde{c})/G}$, and thus a canonical pullback map

$$q^*: H^0(\text{In}_S(Y), K_{\text{In}_S(Y)/[G//G]}) \rightarrow H^0(\text{Fix}(\tilde{c}), K_{\text{Fix}(\tilde{c})/G}). \quad (4.6.1)$$

Next, Lemma 4.5.2 applied to $T = [S/G]$ shows that $\text{cc}_{T/S}(\Lambda_T)$ is an isomorphism $\Lambda_{[G//G]} \xrightarrow{\sim} K_{[G//G]/S}$. This induces an isomorphism

$$K_{\text{In}_S(Y)/[G//G]} \cong p^! \Lambda_{[G//G]} \xrightarrow{\sim} p^! K_{[G//G]/S} \cong K_{\text{In}_S(Y)/S}. \quad (4.6.2)$$

Combining (4.6.1) and (4.6.2), we obtain a canonical map

$$\iota: H^0(\text{In}_S(Y), K_{\text{In}_S(Y)/S}) \rightarrow H^0(\text{Fix}(\tilde{c}), K_{\text{Fix}(\tilde{c})/G}). \quad (4.6.3)$$

The main result of this section is the following theorem.

Theorem 4.6.1. *Notation and assumptions as above, we have an equality*

$$\iota(\text{cc}_{Y/S}(A)) = \text{tr}_{\tilde{c}}(\tilde{u}, A|_{X_G}).$$

Proof. We restate the theorem in the language of based cohomological correspondences. The main players are

- $T = ([S/G], \Lambda_{[S/G]})$, a dualizable object of CoCorr_S .
- $\mathfrak{Y} = (Y \rightarrow [S/G], A, \Lambda_{[S/G]})$, a dualizable object of BCoCorr_S with base T .
- $\mathfrak{X}_G = (X_G, A_{X_G})$, a dualizable object of CoCorr_G with base 1_{CoCorr_G} .
- $\alpha \in \text{End } \mathfrak{X}_G$, the endomorphism described by the pair (\tilde{c}, \tilde{u}) .

We would like to relate $\text{tr}(\text{id}_{\mathfrak{Y}})$ to $\text{tr}(\alpha)$. The idea is to promote α to an endomorphism of based cohomological correspondences which lies over id_T . To this end we introduce some more objects in BCoCorr_S :

- $\mathfrak{G} = (G \rightarrow S, \Lambda_G, \Lambda_S)$ with base 1_{CoCorr_S} ,
- $\mathfrak{G}' = (G \rightarrow [S/G] \times_S [S/G], \Lambda_G, \Lambda_{[S/G]} \boxtimes_S K_{[S/G]/S})$, with base $T \otimes T^\vee$, where the morphism $G \rightarrow [S/G] \times_S [S/G]$ is defined as the trivial $G \times_S G$ -torsor $G_{G \times_S G} = G \times_S G \times_S G$.

We also define morphisms in BCoCorr_S :

- $\text{coev}_G: \mathfrak{G} \rightarrow \mathfrak{G}'$, which has base coev_T and source $1_{(G, \Lambda_G)}$,
- $\text{ev}_G: \mathfrak{G}' \rightarrow \mathfrak{G}$, which has base ev_T and source $1_{(G, \Lambda_G)}$.
- An endomorphism $\alpha_0: \mathfrak{G}' \rightarrow \mathfrak{G}'$ lying over $\text{id}_{T \otimes T^\vee}$, corresponding³ to the automorphism $G_{G \times_S G} \rightarrow G_{G \times_S G}$ defined by $(x, g, h) \mapsto (x, gx, h)$.

Now observe that $\text{tr}_G(\alpha_0) := \text{ev}_G \circ \alpha_0 \circ \text{coev}_G$ is an endomorphism of \mathfrak{G} with base $\text{tr}(\text{id}_T)$ and source $\text{id}_S(\mathfrak{G})$, whose underlying based correspondence is shown in the diagram:

$$\begin{array}{ccccc}
 G & \xleftarrow{\text{id}_G} & G & \xrightarrow{\text{id}_G} & G \\
 \downarrow & & \downarrow & & \downarrow \\
 S & \longleftarrow & [G // G] & \longrightarrow & S
 \end{array}$$

where the central horizontal arrow sends g to its own conjugacy class.

Recall that $\mathcal{S}: \text{BCoCorr}_S \rightarrow \text{CoCorr}_S$ takes a based cohomological correspondce onto its source. Let $\mathcal{F}: \text{CoCorr}_G \rightarrow \text{CoCorr}_S$ be the functor which forgets the base G . We have a diagram in

³Strictly speaking, v-stacks form a 2-category, and we are describing here a 2-morphism from $G \rightarrow [S/G]^2$ to itself.

CoCorr $_S$:

$$\begin{array}{ccc}
\mathcal{S}(1_{\text{BCoCorr}_S} \boxtimes_{1_{\text{CoCorr}_S}} \mathfrak{G}) & \xrightarrow{\sim} & \mathcal{F}(1_{\text{CoCorr}_G}) \\
\mathcal{S}(\text{coev}_{\mathfrak{Y}} \boxtimes_{\text{coev}_T} \text{coev}_G) \downarrow & & \downarrow \mathcal{F}(\text{coev}_{\mathfrak{X}_G}) \\
\mathcal{S}((\mathfrak{Y} \otimes \mathfrak{Y}^\vee) \boxtimes_{T \otimes T^\vee} \mathfrak{G}') & \xrightarrow{\sim} & \mathcal{F}(\mathfrak{X}_G \otimes \mathfrak{X}_G^\vee) \\
\mathcal{S}(\text{id}_{\mathfrak{Y} \otimes \mathfrak{Y}^\vee} \boxtimes_{\text{id}_{T \otimes T^\vee}} \alpha_0) \downarrow & & \downarrow \mathcal{F}(\alpha) \\
\mathcal{S}((\mathfrak{Y} \otimes \mathfrak{Y}^\vee) \boxtimes_{T \otimes T^\vee} \mathfrak{G}') & \xrightarrow{\sim} & \mathcal{F}(\mathfrak{X}_G \otimes \mathfrak{X}_G^\vee) \\
\mathcal{S}(\text{ev}_{\mathfrak{Y}} \boxtimes_{\text{ev}_T} \text{ev}_G) \downarrow & & \downarrow \mathcal{F}(\text{ev}_{\mathfrak{X}_G}) \\
\mathcal{S}(1_{\text{BCoCorr}_S} \boxtimes_{1_{\text{CoCorr}_S}} \mathfrak{G}) & \xrightarrow{\sim} & \mathcal{F}(1_{\text{CoCorr}_G})
\end{array}$$

where all squares are filled in with 2-isomorphisms. The functoriality property of \boxtimes from (4.5.8) now gives a 2-isomorphism

$$\mathcal{S} \left(\text{cc}_{Y/S}(A) \boxtimes_{\text{cc}_{[S/G]/S}(\Lambda_{[S/G]})} \text{tr}_G(\alpha_0) \right) \xrightarrow{\sim} \mathcal{F}(\text{tr}(\alpha)). \quad (4.6.4)$$

The isomorphism of v-stacks implicit in (4.6.4) is expressed by the fact that we have a cartesian diagram:

$$\begin{array}{ccc}
\text{Fix}(\tilde{c}) & \longrightarrow & G \\
\downarrow & & \downarrow \\
\text{In}_S(Y) & \longrightarrow & [G // G]
\end{array}$$

On the level of cohomology classes, (4.6.4) tells us that $\text{tr}(\alpha)$, considered as an element of $H^0(\text{Fix}(\tilde{c}), K_{\text{Fix}(\tilde{c})/S})$ can be derived from $\text{cc}_{Y/S}(A)$ in the manner described by the theorem. \square

We conclude the section with a remark about isolated fixed points. Assume there exists a conjugacy-invariant open subset $U \subset G$ whose elements act on X with only isolated fixed points. (This is the case for the action of the positive loop group on the affine Grassmannian. We study that scenario in the next section.) Write $\text{In}_S([X/G])_U$ for the pullback of $[U // G]$ under $\text{In}_S([X/G]) \rightarrow [G // G]$. Then $\text{In}_S([X/G])_U \rightarrow [U // G]$ is étale over $[U // G]$; as such we have a canonical trivialization $K_{\text{In}_S([X/G])_U/S} \cong \Lambda_{\text{In}_S([X/G])_U}$. Therefore the restriction over U of the characteristic class of A is an element

$$\text{cc}_{[X/G]/S}(A)^U \in H^0(\text{In}_S([X/G])_U, \Lambda);$$

that is, it is a continuous function on the space of pairs $(x, g) \in X \times_S U$ with $g(x) = x$. Theorem 4.6.1 implies that this function is $(x, g) \mapsto \text{loc}_x(g, A)$.

5 Local terms on the B_{dR} -affine Grassmannian

The goal of this chapter is to explicitly compute certain local terms on the B_{dR} -affine Grassmannian, in terms of the geometric Satake equivalence.

5.1 The main result

The goal of this chapter is to explicitly compute certain local terms on the B_{dR} -affine Grassmannian, in terms of the geometric Satake equivalence.

To explain the main result, let us fix some notation. Let F/\mathbf{Q}_p be a finite extension with residue field \mathbf{F}_q . Let C be the completion of an algebraic closure of F . Let G/F be a split reductive group, and let $\text{Gr}_G = LG/L^+G$ be the associated B_{dR} -affine Grassmannian over $\text{Spd } C$. We explain the notation: for a perfectoid C -algebra R , we have the loop group $LG(R) = B_{\text{dR}}(R)$ and its positive subgroup $L^+G(R) = B_{\text{dR}}^+(R)$. Then Gr_G is an ind-spatial diamond admitting an action of L^+G and in particular its subgroup $G(F)$. For a cocharacter μ of $G_{\overline{F}}$, we let $\text{Gr}_{G, \leq \mu}$ be the corresponding closed Schubert cell; this is a proper diamond. Finally, define the *local Hecke stack* by

$$\text{Hecke}_G^{\text{loc}} = [L^+G \backslash \text{Gr}_G] = [L^+G \backslash LG/L^+G].$$

We remark that there are versions of these objects living over $\text{Spd } F$, but we will not need these for our results.

Fix a coefficient ring $\Lambda \in \{\mathbf{Z}/\ell^n\mathbf{Z}, \mathbf{Z}_\ell\}$. The *Satake category*

$$\text{Sat}_G(\Lambda) \subset D_{\text{ét}}(\text{Hecke}_{G,C}^{\text{loc}}, \Lambda)$$

is the subcategory of objects which are perverse, Λ -flat, and ULA over $\text{Spd } C$ [FS21, Definition I.6.2]. It is a symmetric monoidal category under the convolution product.

Theorem 5.1.1 ([FS21, Theorem I.6.3]). *There is an equivalence of symmetric monoidal categories:*

$$\begin{array}{ccc} \text{Rep}_{\widehat{G}}(\Lambda) & \xrightarrow{\sim} & \text{Sat}_G(\Lambda) \\ V & \mapsto & \mathcal{S}_V \end{array}$$

where \widehat{G} is the Langlands dual group (considered over Λ), and $\text{Rep}_{\widehat{G}}(\Lambda)$ is the category of representations of \widehat{G} on finite projective Λ -modules.

We continue to write \mathcal{S}_V for the pullback of this object along the quotient $\text{Gr}_G \rightarrow [L^+G \backslash \text{Gr}_G]$.

Our next order of business is to determine, for $g \in G(F)_{\text{sr}}$, the fixed point locus Gr_G^g . The answer is the same regardless of which sort of affine Grassmannian we consider (classical, Witt vector, B_{dR}), as the following proposition shows.

Proposition 5.1.2. *Let K^+ be a discrete valuation ring with algebraically closed residue field k and fraction field K . Let G be a reductive group over K^+ . Let $g \in G(K^+)$ be an element whose image in $G(k)$ is strongly regular, and let $T = \text{Cent}(g, G)$. The inclusion $T \subset G$ induces a bijection*

$$T(K)/T(K^+) \cong (G(K)/G(K^+))^g,$$

so that the fixed point locus of g may be identified with $X_*(T)$.

Consequently if Gr_G is any incarnation of the affine Grassmannian, then $\text{Gr}_{G, \leq \mu}^g$ is finite over its base with underlying set $X_*(T)_{\leq \mu}$.

Proof. Let \mathcal{B} be the (reduced) Bruhat-Tits building of the split reductive group G_K over the discretely valued field K . Thus \mathcal{B} is a locally finite simplicial complex admitting an action of $LG = G(K)$. We will identify the LG -set Gr_G with a piece of this building.

By [BT84, 5.1.40] there exists a hyperspecial point $\bar{o} \in \mathcal{B}$ corresponding to $L^+G = G(K^+)$. The point \bar{o} can be characterized by [BT84, 4.6.29] as the unique fixed point of L^+G . Let \mathcal{B}^{ext} be the extended Bruhat-Tits building of G_K . Recall that $\mathcal{B}^{\text{ext}} = \mathcal{B} \times X_*(A_G)_{\mathbf{R}}$, where A_G is the connected center of G . The group LG acts on $X_*(A_G)_{\mathbf{R}}$ via the isomorphism $X_*(A_G)_{\mathbf{R}} \rightarrow X_*(A'_G)_{\mathbf{R}}$, where A'_G is the maximal abelian quotient of G . Let $o = (\bar{o}, z)$ be any point in \mathcal{B}^{ext} lying over \bar{o} . Then L^+G can be characterized as the full stabilizer of o in $G(K)$: It is clear that L^+G stabilizes o , and the reverse inclusion follows from the Cartan decomposition $LG = L^+G \cdot X_*(T) \cdot L^+G$ (which relies on \bar{o} being hyperspecial) and the fact that $X_*(T)$ acts on the apartment of T in \mathcal{B}^{ext} by translations. It follows that the action of LG on \mathcal{B}^{ext} provides an LG -equivariant bijection from Gr_G to the orbit of LG through o .

Now suppose $x \in \text{Gr}_G$ is fixed by a strongly regular element $g \in L^+T_{\text{sr}}$. Then its image in \mathcal{B}^{ext} is a g -fixed point belonging to the orbit of o , and we can write $x = ho$ for some $h \in LG$. For every root $\alpha: T \rightarrow \mathbf{G}_m$, the element $\alpha(g)$ does not lie in the kernel of $L^+\mathbf{G}_m \rightarrow \mathbf{G}_m$. According to [Tit79, 3.6.1] the image of x in \mathcal{B} belongs to the apartment \mathcal{A} of T . At the same time, $g \in L^+G$ also fixes \bar{o} , so for the same reason $\bar{o} \in \mathcal{A}$. Thus \bar{o} belongs to both apartments \mathcal{A} and $h^{-1}\mathcal{A}$. Since L^+G acts transitively on the apartments containing \bar{o} [BT84, 4.6.28], we can multiply h on the right by an element of L^+G to ensure that $h^{-1}\mathcal{A} = \mathcal{A}$. By [BT72, 7.4.10] we then have $h \in L^+N(T, G)$. Since \bar{o} is hyperspecial, every Weyl reflection is realized in L^+G and hence we may again modify h on the right to achieve $h \in LT$. We see now that $x = ho$ is fixed by all of LT and that furthermore the coset $x = hL^+G$ is the image of the coset hL^+T . \square

Proposition 5.1.2 shows that if we fix a split maximal torus $\widehat{T} \subset \widehat{G}$, there is a natural finite-to-one map

$$\begin{aligned} \text{Gr}_G^g &\rightarrow X_+^*(\widehat{T}) \\ x &\mapsto \nu_x. \end{aligned}$$

Note that ν_x simply records which open Schubert cell of Gr_G contains the point x .

Now, for any $V \in \text{Rep}(\widehat{G})$ and any $x \in \text{Gr}_G^g$, there is an associated local term $\text{loc}_x(g, \mathcal{S}_V) \in \Lambda$. The main result of this chapter is the following theorem, giving an explicit computation of these local terms.

Theorem 5.1.3. *Let $V \in \text{Rep}(\widehat{G})$ be an object of the Satake category, and let $g \in G(F)_{\text{sr}}$ be a strongly regular semisimple element. Then for any $x \in \text{Gr}_G^g$, there is an equality in Λ :*

$$\text{loc}_x(g, \mathcal{S}_V) = (-1)^{\langle 2\rho, \nu_x \rangle} \text{rank}_{\Lambda} V[\nu_x].$$

Note that since V is (by hypothesis) a finite projective Λ -module, and tori are reductive in the strongest sense, the weight space $V[\nu_x]$ is a finite projective Λ -module, so the right-hand side of this equality is well-defined.

Due to the highly inexplicit nature of local terms, the proof of Theorem 5.1.3 is rather indirect. Indeed, we would be able to give a simple proof of Theorem 5.1.3 *if* we knew the equality between “true” and “naive” local terms on Gr_G . Unfortunately, this equality seems to be a very difficult problem. Even for schemes, the problem of comparing true and naive local terms was only settled very recently by Varshavsky. Instead, our strategy reduces the computation of the local terms in Theorem 5.1.3 to an analogous computation on the Witt vector affine Grassmannian, where a global-to-local argument can be pushed through. The key theme in the proof is the idea that *local terms are constant in families*.

For our applications, the following restatement of the main results of this section in terms of characteristic classes on the quotient $[\mathrm{Gr}_{G, \leq \mu} / L_m^+ G]$ will be useful.

Theorem 5.1.4. *Let V be such that \mathcal{S}_V is supported on some Schubert cell $\mathrm{Gr}_{G, \leq \mu}$. Choose some large m such that the L^+G -action on this cell factors through the quotient $L_m^+ G$, and set $X = [\mathrm{Gr}_{G, \leq \mu} / L_m^+ G]$.*

Then the set of connected components of $\mathrm{In}_S(X)_{\mathrm{sr}}$ may be identified with $X_^+(T)_{\leq \mu} / W$, and the dualizing complex of $\mathrm{In}_S(X)_{\mathrm{sr}}$ has a canonical trivialization. With respect to those identifications, the restriction of $\mathrm{cc}_{X/S}(\mathcal{S}_V)$ to $\mathrm{In}_S(X)_{\mathrm{sr}}$ is the function sending $\lambda \in X_*(T)$ to $(-1)^{(2\rho_G, \lambda)} \mathrm{rank}_\Lambda V[\lambda]$.*

Proof. The first claim is proved in §5.4 below. Since $L_m^+ G$ is a cohomologically smooth group diamond, Theorem 4.6.1 applies to the quotient $X = [\mathrm{Gr}_{G, \leq \mu} / L_m^+ G]$. The remark following the proof of that theorem applies to the locus $L_m^+ G_{\mathrm{sr}}$, so that we may relate $\mathrm{cc}_{X/S}(\mathcal{S}_V)$ to the local terms $\mathrm{loc}_x(g, \mathcal{S}_V)$. The latter have been computed by Theorem 5.1.3. \square

5.2 Strategy of proof

In this section, we reduce Theorem 5.1.3 to four auxiliary propositions stated below. The proofs of these propositions will occupy the remainder of this chapter.

First, we show that the local terms appearing in Theorem 5.1.3 are essentially independent of g .

Proposition 5.2.1. *In the notation and setup of Theorem 5.1.3, $\mathrm{loc}_x(g, \mathcal{S}_V)$ depends on g and x only through the cocharacter ν_x . More precisely, if $g, g' \in G(F)_{\mathrm{sr}}$ are two strongly regular semisimple elements and $x \in \mathrm{Gr}_G^g$ resp. $x' \in \mathrm{Gr}_G^{g'}$ are fixed points such that $\nu_x = \nu_{x'}$, then*

$$\mathrm{loc}_x(g, \mathcal{S}_V) = \mathrm{loc}_{x'}(g', \mathcal{S}_V).$$

Next, we are going to degenerate from characteristic zero into characteristic p . For this, fix a split reductive model $\mathcal{G}/\mathcal{O}_F$ of G , and let $\mathrm{Gr}_{\mathcal{G}}$ be the associated Beilinson-Drinfeld affine Grassmannian over $S = \mathrm{Spd} \mathcal{O}_C$. Recall that this is a small v -sheaf which interpolates between the B_{dR} -affine Grassmannian Gr_G and the Witt vector affine Grassmannian Gr_G^W , in the sense that we have a

commutative diagram

$$\begin{array}{ccccc}
(\mathrm{Gr}_G^W)^\diamond & \xrightarrow{i} & \mathrm{Gr}_G & \xleftarrow{j} & \mathrm{Gr}_G \\
\downarrow & & \downarrow & & \downarrow \\
s = \mathrm{Spd} \overline{\mathbf{F}}_q & \longrightarrow & S & \longleftarrow & \eta = \mathrm{Spd} C
\end{array}$$

with cartesian squares. We will crucially use the fact that all of these Grassmannians satisfy compatible forms of geometric Satake, in the sense that there are natural monoidal functors

$$\begin{array}{ccc}
& D_{\text{ét}}(\mathrm{Gr}_G, \Lambda) & \\
& \nearrow & \uparrow j^* \\
\mathrm{Rep}(\widehat{G}) & \longrightarrow D_{\text{ét}}(\mathrm{Gr}_G, \Lambda) & \\
& \searrow & \downarrow i^* \\
& D_{\text{ét}}((\mathrm{Gr}_G^W)^\diamond, \Lambda) &
\end{array}$$

such that the vertical arrows are equivalences of categories on the essential images of $\mathrm{Rep}(\widehat{G})$.

Proposition 5.2.2. *Let $g \in \mathcal{G}(\mathcal{O}_F)$ be an element such that $\bar{g} \in \mathcal{G}(\mathbf{F}_q)$ is strongly regular semisimple. Then $\mathcal{T} = \mathrm{Cent}(g, \mathcal{G})$ is a maximal torus, and there is a natural isomorphism*

$$\mathrm{Gr}_G^g \cong \mathrm{Gr}_{\mathcal{T}} \cong X_*(\mathcal{T})_S.$$

In particular, if $\beta \simeq S \subset \mathrm{Gr}_G^g$ is any connected component, then β_η and β_s are isolated fixed points for the g -action in the generic and special fiber, respectively.

Proposition 5.2.3. *Let $g \in \mathcal{G}(\mathcal{O}_F)$ be an element such that $\bar{g} \in \mathcal{G}(\mathbf{F}_q)$ is strongly regular semisimple. Then for any $V \in \mathrm{Rep}(\widehat{G})$ and any connected component $\beta \subset \mathrm{Gr}_G^g$, we have the equality*

$$\mathrm{loc}_{\beta_\eta}(g, j^* \mathcal{S}_V) = \mathrm{loc}_{\beta_s}(g, i^* \mathcal{S}_V)$$

of local terms.

Finally, we compute the local terms on the Witt vector affine Grassmannian by a direct argument.

Proposition 5.2.4. *Let $g \in \mathcal{G}(\mathcal{O}_F)$ be an element with finite prime-to- p order such that $\bar{g} \in \mathcal{G}(\mathbf{F}_q)$ is strongly regular semisimple. Then for any $x \in \mathrm{Gr}_G^{W, g}$ and any $V \in \mathrm{Rep}(\widehat{G})$,*

$$\mathrm{loc}_x(g, \mathcal{S}_V) = (-1)^{\langle 2\rho, \nu_x \rangle} \mathrm{rank}_\Lambda V[\nu_x]$$

where $\nu_x \in X_+^*(\widehat{T})$ is as before.

5.3 Local terms and base change

In this section we prove two key technical results, namely that formation of local terms commutes with any base change, and with passage from perfect schemes to v-sheaves.

In order to fix notation, we briefly recall the key definitions concerning local terms; we apologize for the overlap with Chapter 4. Let S be a small v-sheaf, which will be our base object. Let $f : X \rightarrow S$ be a map of v-sheaves representable in nice diamonds. Consider a correspondence $c = (c_1, c_2) : C \rightarrow X \times_S X$ given by a map of v-sheaves representable in nice diamonds. This gives rise to a cartesian diagram

$$\begin{array}{ccc} \mathrm{Fix}(c) & \xrightarrow{\Delta'} & C \\ c' \downarrow & & \downarrow c=(c_1, c_2) \\ X & \xrightarrow{\Delta} & X \times_S X \end{array}$$

of small v-sheaves. We will sometimes assume that c_1 is proper and that $\mathrm{Fix}(c)$ is a disjoint union of open-closed subspaces which are proper over S . These conditions will hold e.g. if f and c are proper.

Let $\mathcal{F} \in D_{\text{ét}}(X, \Lambda)$ be an f -ULA object. Recall that a cohomological correspondence over c is a map $u : Rc_2!c_1^*\mathcal{F} \rightarrow \mathcal{F}$, i.e. an element $u \in \mathrm{Hom}(c_1^*\mathcal{F}, Rc_2!\mathcal{F})$. If c_1 is proper, then applying $Rf_!$ induces an endomorphism

$$\begin{aligned} Rf_!u : Rf_!\mathcal{F} &\rightarrow Rf_!Rc_1*c_1^*\mathcal{F} = Rf_!Rc_1!c_1^*\mathcal{F} \\ &\cong Rf_!Rc_2!c_1^*\mathcal{F} \xrightarrow{u} Rf_!\mathcal{F} \end{aligned}$$

of $Rf_!\mathcal{F}$. On the other hand, there is a natural map

$$\mathrm{Hom}(c_1^*\mathcal{F}, Rc_2!\mathcal{F}) \rightarrow H^0(\mathrm{Fix}(c), K_{\mathrm{Fix}(c)/S}),$$

cf. the discussion immediately before Definition 4.3.5, and we write $\mathrm{tr}_c(u, \mathcal{F}) \in H^0(\mathrm{Fix}(c), K_{\mathrm{Fix}(c)/S})$ for the image of u under this map.

If $\beta \subset \mathrm{Fix}(c)$ is a closed-open subspace with *proper* structure map $g : \beta \rightarrow S$, then $H^0(\beta, K_{\beta/S}) = H^0(\beta, Rg^!\Lambda)$ is canonically a direct summand of $H^0(\mathrm{Fix}(c), K_{\mathrm{Fix}(c)/S})$, and we can further consider the image of $\mathrm{tr}_c(u, \mathcal{F})$ under the map

$$\begin{aligned} H^0(\mathrm{Fix}(c), K_{\mathrm{Fix}(c)/S}) &\rightarrow H^0(\beta, Rg^!\Lambda) \cong H^0(S, Rg_*Rg^!\Lambda) \\ &\cong H^0(S, Rg!Rg^!\Lambda) \rightarrow H^0(S, \Lambda). \end{aligned}$$

By definition, this is the local term $\mathrm{loc}_\beta(u, \mathcal{F})$. In most situations we care about, S is connected, so $H^0(S, \Lambda) = \Lambda$ and we simply regard $\mathrm{loc}_\beta(u, \mathcal{F})$ as an element of Λ . Note that local terms are additive, in the sense that if $\beta = \beta_1 \amalg \beta_2$, then $\mathrm{loc}_\beta(u, \mathcal{F}) = \mathrm{loc}_{\beta_1}(u, \mathcal{F}) + \mathrm{loc}_{\beta_2}(u, \mathcal{F})$. If S is a geometric point, f and c are proper, and $\pi_0(\mathrm{Fix}(c))$ is a discrete (and therefore finite) set, the usual Lefschetz trace formula holds, and says that

$$\mathrm{tr}(u | R\Gamma_c(X, \mathcal{F})) = \sum_{\beta \in \pi_0(\mathrm{Fix}(c))} \mathrm{loc}_\beta(u, \mathcal{F}).$$

We need to understand how local terms interact with base change on S . More precisely, assume we are given a morphism $a : T \rightarrow S$. Then all objects and morphisms above naturally base change to objects over T . Note that $\text{Fix}(c)_T = \text{Fix}(c_T)$. We write $a_X : X_T \rightarrow X$, $a_C : C_T \rightarrow C$, etc. for the base changes of a . We naturally get a cohomological correspondence u_T on $\mathcal{F}_T = a_X^* \mathcal{F}$ over c_T by taking the image of u under the map

$$\begin{aligned} \text{Hom}(c_1^* \mathcal{F}, Rc_2^! \mathcal{F}) &\rightarrow \text{Hom}(a_C^* c_1^* \mathcal{F}, a_C^* Rc_2^! \mathcal{F}) \cong \text{Hom}(c_{1,T}^* a_X^* \mathcal{F}, a_C^* Rc_2^! \mathcal{F}) \\ &\rightarrow \text{Hom}(c_{1,T}^* a_X^* \mathcal{F}, Rc_{2,T}^! a_X^* \mathcal{F}). \end{aligned}$$

The final arrow here is induced by the canonical map $a_C^* Rc_2^! \mathcal{F} \rightarrow Rc_{2,T}^! a_X^* \mathcal{F}$. This map is a special case of the natural transformation $\beta_{f,g} : \tilde{f}^* Rg^! \rightarrow R\tilde{g}^! f^*$ which exists for any cartesian diagram

$$\begin{array}{ccc} X' & \xrightarrow{\tilde{f}} & Y' \\ \tilde{g} \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

with g representable in nice diamonds. The transformation in question is adjoint to the map $R\tilde{g}^! f^* Rg^! \cong f^* Rg_! Rg^! \rightarrow f^*$ (it is also adjoint to the map $Rg^! \rightarrow Rg^! Rf_* f^* \cong R\tilde{f}_* R\tilde{g}^! f^*$).

In this setup, the next proposition says that formation of local terms commutes with base change along $T \rightarrow S$.

Proposition 5.3.1. *For any given $\beta \subset \text{Fix}(c)$ as above, the natural map*

$$H^0(S, \Lambda) \rightarrow H^0(T, \Lambda)$$

sends $\text{loc}_\beta(u, \mathcal{F})$ to $\text{loc}_{\beta_T}(u_T, \mathcal{F}_T)$. In particular, if S and T are connected, then $\text{loc}_\beta(u, \mathcal{F}) = \text{loc}_{\beta_T}(u_T, \mathcal{F}_T)$ as elements of Λ .

Proof. By a straightforward argument, this reduces to showing that there is a natural map

$$H^0(\text{Fix}(c), K_{\text{Fix}(c)/S}) \rightarrow H^0(\text{Fix}(c)_T, K_{\text{Fix}(c)_T/T})$$

compatible with the map $H^0(S, \Lambda) \rightarrow H^0(T, \Lambda)$, and sending $\text{tr}_c(u, \mathcal{F})$ to $\text{tr}_{c_T}(u_T, \mathcal{F}_T)$. To obtain the map itself, apply $H^0(\text{Fix}(c), -)$ to the composition

$$\begin{aligned} K_{\text{Fix}(c)/S} &= R(f \circ c')^! \Lambda \rightarrow Ra_{\text{Fix}(c)*} a_{\text{Fix}(c)}^* R(f \circ c')^! \Lambda \\ &\xrightarrow{\beta_{a, f \circ c'}} Ra_{\text{Fix}(c)*} R(f_T \circ c'_T)^! a^* \Lambda = Ra_{\text{Fix}(c)*} K_{\text{Fix}(c)_T/T}. \end{aligned}$$

The claim about the relation between tr_c and tr_{c_T} now follows from the fact that the base change functor $\text{CoCorr}_S \rightarrow \text{CoCorr}_T$ is symmetric monoidal, and therefore preserves dualizable objects and traces of endomorphisms thereof. \square

We will also need to compare local terms associated with perfect schemes and with v-sheaves. More precisely, fix a perfect field k/\mathbf{F}_p , and let PSch_k be the category of perfect schemes over k . There is a natural functor $X \mapsto X^\diamond$ from PSch_k to small v-sheaves over $\text{Spd } k$, characterized by $(\text{Spec } R)^\diamond(A, A^+) = \text{Hom}_k(R, A)$. Said differently, X^\diamond sends $\text{Spec } R$ to $\text{Spa}(R, R^+)^\diamond$ where R^+ is the integral closure of k in R . This functor commutes with finite limits. Moreover, if $f : X \rightarrow Y$ is separated and perfectly of finite type, then f^\diamond is representable in locally spatial diamonds and compactifiable with finite dim.trg. By [Sch17, §27], for any X there is a fully faithful symmetric monoidal functor $c_X^* : D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X^\diamond, \Lambda)$ compatible with f^* and $Rf_!$ in the evident senses. Moreover, one has canonical natural transformations

$$c_X^* R\mathcal{H}\text{om}(-, -) \rightarrow R\mathcal{H}\text{om}(c_X^* -, c_X^* -)$$

and $c_X^* Rf^! \rightarrow Rf^{\diamond!} c_Y^*$ for f separated and perfectly of finite type.

Now, let $\text{PSch}_k^{\text{ft}}$ be the full subcategory of schemes separated and perfectly of finite type over k . Fix $X \in \text{PSch}_k^{\text{ft}}$ with structure map $f : X \rightarrow \text{Spec } k$, and let $c : C \rightarrow X \times_k X$ be a correspondence in $\text{PSch}_k^{\text{ft}}$ such that c_1 and $f \circ c'$ are perfectly proper. Let $\mathcal{F} \in D_{\text{ét}}(X, \Lambda)$ be an f -ULA object equipped with a cohomological correspondence u lying over c , so we get local terms $\text{loc}_\beta(u, \mathcal{F}) \in H^0(\text{Spec } k, \Lambda) = \Lambda$ by the schematic version of the recipe recalled above.

On the other hand, applying $(-)^{\diamond}$ and using commutation with finite limits, we get a correspondence $c^\diamond : C^\diamond \rightarrow X^\diamond \times_{\text{Spd } k} X^\diamond$ of v-sheaves over $S = \text{Spd } k$ with $\text{Fix}(c)^\diamond = \text{Fix}(c^\diamond)$, satisfying all of our assumptions from above. Moreover, u naturally induces a cohomological correspondence u^\diamond on $c_X^* \mathcal{F}$ lying over c^\diamond , by taking the image of u under the natural map

$$\begin{aligned} \text{Hom}(c_1^* \mathcal{F}, Rc_2^! \mathcal{F}) &\rightarrow \text{Hom}(c_C^* c_1^* \mathcal{F}, c_C^* Rc_2^! \mathcal{F}) \cong \text{Hom}(c_1^{\diamond*} c_X^* \mathcal{F}, c_C^* Rc_2^! \mathcal{F}) \\ &\rightarrow \text{Hom}(c_1^{\diamond*} c_X^* \mathcal{F}, Rc_2^{\diamond!} c_X^* \mathcal{F}). \end{aligned}$$

Proposition 5.3.2. *Maintain the previous setup and notation. Then $c_X^* \mathcal{F}$ is f^\diamond -ULA, and for any open-closed $\beta \subset \text{Fix}(c)$, we have an equality*

$$\text{loc}_\beta(u, \mathcal{F}) = \text{loc}_{\beta^\diamond}(u^\diamond, c_X^* \mathcal{F})$$

of local terms.

Proof. This is formally identical to the proof of Proposition 5.3.1, using the fact that $(-)^{\diamond}$ induces a symmetric monoidal functor on the appropriate categories of cohomological correspondences. \square

5.4 Independence of g

In this section we prove Proposition 5.2.1. In this section only, we set $S = \text{Spd } C$.

Fix V as in the proposition. Decomposing V into isotypic summands for the action of $Z(\widehat{G})^\circ$, we can assume that \mathcal{S}_V is supported on a single connected component of Gr_G . We can then pick some μ such that \mathcal{S}_V is supported on the Schubert cell $\text{Gr}_{G, \leq \mu}$. Choose some large m such that the L^+G action on $\text{Gr}_{G, \leq \mu}$ factors over the truncated loop group L_m^+G . The sheaf \mathcal{S}_V is naturally the pullback of a sheaf again denoted \mathcal{S}_V on the quotient stack $X = [\text{Gr}_{G, \leq \mu} / L_m^+G]$, so we can consider the characteristic class $\text{cc}_{X/S}(\mathcal{S}_V)$.

To analyze this class, we need to understand the inertia stack of X . For this, we need some notation. Let $L_m^+ G_{\text{sr}}$ be the preimage of the strongly regular semisimple locus $G_{\text{sr}} \subset G$ under the theta map $L_m^+ G \rightarrow G$. Pick any maximal torus $T \subset G$ with Weyl group W , and set $L_m^+ T_{\text{sr}} = L_m^+ T \cap L_m^+ G_{\text{sr}}$.

Proposition 5.4.1. *1. The open substack*

$$\text{In}_S([S/L_m^+ G]_{\text{sr}}) = [L_m^+ G_{\text{sr}} // L_m^+ G] \subset \text{In}_S([S/L_m^+ G])$$

is canonically identified with $[L_m^+ T_{\text{sr}} / (W \times L_m^+ T)]$ via the natural map.

2. The open substack

$$\text{In}_S(X)_{\text{sr}} = \text{In}_S(X) \times_{\text{In}_S([S/L_m^+ G])} [L_m^+ G_{\text{sr}} // L_m^+ G] \subset \text{In}_S(X)$$

is canonically identified with $X_(T)_{\leq \mu} \times^W [L_m^+ T_{\text{sr}} // L_m^+ T]$, such that the natural map $\text{In}_S(X)_{\text{sr}} \rightarrow \text{In}_S([S/L_m^+ G]_{\text{sr}})$ coincides via the identification in part 1. with the evident projection onto $[L_m^+ T_{\text{sr}} / (W \times L_m^+ T)]$.*

Proof. The idea behind (1) is that any $g \in L_m^+ G_{\text{sr}}$ is conjugate to an element of $L_m^+ T_{\text{sr}}$, which is well-defined up to the action of the normalizer of this group, which is $W \times L_m^+ T$.

For (2), we observe that an object of $\text{In}_S(X)_{\text{sr}}$ is a pair (x, g) , where $g \in L_m^+ G_{\text{sr}}$ fixes $x \in \text{Gr}_{G, \leq \mu}$; the automorphisms of this object are $L_m^+ G$. The g can be conjugated to lie in $L_m^+ T_{\text{sr}}$, and then by Proposition 5.1.2, the x can be identified with an element of $X_*(T)_{\leq \mu}$, which is well-defined up to an element of W . \square

Corollary 5.4.2. *There is a natural isomorphism*

$$H^0(\text{In}_S(X)_{\text{sr}}, K_{\text{In}_S(X)_{\text{sr}}/S}) \cong C(X_*(T)_{\leq \mu}, \Lambda)^W$$

which sends $\text{cc}_{X/S}(\mathcal{S}_V)$ to the function sending $\lambda \in X_(T)_{\leq \mu}$ to $\text{loc}_{x_\lambda}(g, \mathcal{S}_V)$, where $x_\lambda \in \text{Gr}_{G, \leq \mu}$ is the T -fixed point corresponding to λ , and $g \in L_m^+ T_{\text{sr}}$. In particular, $\text{loc}_{x_\lambda}(g, \mathcal{S}_V)$ does not depend on the choice of $g \in L_m^+ T_{\text{sr}}$.*

Proof. Combine Theorem 4.6.1 with the description of $\text{In}_S(X)_{\text{sr}}$ from Proposition 5.4.1. \square

5.5 Degeneration to characteristic p

In this section we prove Propositions 5.2.2 and 5.2.3.

Proof of Proposition 5.2.2. The isomorphism $\text{Gr}_{\mathcal{T}} \cong X_*(\mathcal{T})_S$ is [SW20, Proposition 21.3.1]. There is an evident map $f : \text{Gr}_{\mathcal{T}} \rightarrow \text{Gr}_{\mathcal{G}}^g$, and it remains to see that f is an isomorphism. For this, we first note that f is a closed immersion. This follows from the observation the source and target of f are both closed subfunctors of $\text{Gr}_{\mathcal{G}}$. For the source, this follows from [SW20, Proposition 20.3.7], while for the target this follows from the fact that $\text{Gr}_{\mathcal{G}} \rightarrow S$ is separated.

Since f is a closed immersion, it is both qcqs and specializing. By [Sch17, Lemma 12.5] it is enough to check that f is a bijection on rank one geometric points. This can be checked separately on the generic and special fibers. Both cases are handled by Proposition 5.1.2. \square

Proof of Proposition 5.2.3. By two applications of (the connected case of) Proposition 5.3.1, applied to the maps $\eta \rightarrow S$ and $s \rightarrow S$, we get equalities

$$\mathrm{loc}_{\beta_\eta}(g, j^* \mathcal{S}_V) = \mathrm{loc}_\beta(g, \mathcal{S}_V) = \mathrm{loc}_{\beta_s}(g, i^* \mathcal{S}_V),$$

and the result follows. \square

5.6 Local terms on the Witt vector affine Grassmannian

Proof of Proposition 5.2.4. Fix g and V as in the statement, and let $\mathcal{T} \subset \mathcal{G}$ be the connected centralizer of g . For every $\nu \in X_*(\mathcal{T})$, let $S_\nu \subset \mathrm{Gr}_\mathcal{G}^W$ be the associated semi-infinite orbit, with closure $\overline{S}_\nu = \cup_{\nu' \leq \nu} S_{\nu'}$.

Let $X \subset \mathrm{Gr}_\mathcal{G}^W$ be a finite union of closed Schubert cells containing the support of \mathcal{S}_V , so X is a perfectly projective k -scheme by the results in [BS17]. Write $X_\nu = X \cap S_\nu$, $X_{\leq \nu} = X \cap \overline{S}_\nu$, and $\partial X_{\leq \nu} = X_{\leq \nu} \setminus X_\nu$. Note that all of these spaces are stable under g , and in fact under \mathcal{T} . Note also that each X_ν contains a *unique* g -fixed point x_ν .

Proposition 5.6.1. *The compactly supported Euler characteristic of \mathcal{S}_V on X_ν is*

$$\chi_c(X_\nu, \mathcal{S}_V) = (-1)^{\langle 2\rho, \nu \rangle} \mathrm{rank} V[\nu].$$

Proof. This is a consequence of the integral-coefficients version of the Satake equivalence for the Witt vector affine Grassmannian given in [Yu19]. There it is shown (Proposition 4.2) that $H_c^d(X_\nu, \mathcal{S}_V)$ is zero unless $d = \langle 2\rho, \nu \rangle$, and in that degree it corresponds exactly to the ν -weight functor in the Satake category. \square

Any $t \in \mathcal{T}$ must act trivially on $H_c^d(X_\nu, \mathcal{S}_V)$, so $\chi_c(U, \mathcal{S}_V)$ coincides with the trace of g on $R\Gamma_c(X_\nu, \mathcal{S}_V)$. The same is true for $X_{\leq \nu}$ and $\partial X_{\leq \nu}$. We compute:

$$\begin{aligned} (-1)^{\langle 2\rho, \nu \rangle} \mathrm{rank} V[\nu] &= \chi_c(X_\nu, \mathcal{S}_V) \\ &= \chi_c(X_{\leq \nu}, \mathcal{S}_V) - \chi_c(\partial X_{\leq \nu}, \mathcal{S}_V) \\ &= \mathrm{tr}(g | R\Gamma_c(X_{\leq \nu}, \mathcal{S}_V)) - \mathrm{tr}(g | R\Gamma_c(\partial X_{\leq \nu}, \mathcal{S}_V)) \\ &= \sum_{\nu' \leq \nu} \mathrm{loc}_{x_{\nu'}}(g, \mathcal{S}_V|_{X_{\leq \nu}}) - \sum_{\nu' \leq \nu, \nu' \neq \nu} \mathrm{loc}_{x_{\nu'}}(g, \mathcal{S}_V|_{\partial X_{\leq \nu}}) \\ &= \sum_{\nu' \leq \nu} \mathrm{loc}_{x_{\nu'}}(g, \mathcal{S}_V) - \sum_{\nu' \leq \nu, \nu' \neq \nu} \mathrm{loc}_{x_{\nu'}}(g, \mathcal{S}_V) \\ &= \mathrm{loc}_{x_\nu}(g, \mathcal{S}_V). \end{aligned}$$

The penultimate equality is the key technical fact, and follows from Proposition 5.6.2 below together with the assumptions on g . \square

Proposition 5.6.2. *Let k/\mathbf{F}_p be a perfect field, and let X be a perfectly finite type k -scheme with an automorphism $g : X \rightarrow X$ of finite prime-to- p order. Let $A \in D_c^b(X, \mathbf{Z}_\ell)$ be an object equipped with a morphism $u : g^* A \rightarrow A$. Then for every isolated g -fixed point x , the true local term $\mathrm{loc}_x(g, A)$ equals the naive local term $\mathrm{tr}(g|_{A_x})$.*

In particular, if $Z \subset X$ is a g -stable closed subscheme, then $\mathrm{loc}_x(g, A) = \mathrm{loc}_x(g, A|_Z)$.

Proof. With the word “perfectly” deleted, this is a recent result of Varshavsky [Var20] (combine Theorem 4.10(b) and Corollary 5.4(b)). We will reduce to Varshavsky’s result by deperfecting.

Precisely, since $\mathrm{loc}_x(g, A)$ is insensitive to replacing X, A by $U, A|_U$ for $U \subset X$ any g -invariant open neighborhood of x , we can assume that X is affine, so $X = \mathrm{Spec} R$ with R perfectly of finite type. Let $R_0 \subset R$ be a finite type k -algebra with $R_0^{\mathrm{perf}} = R$, and let $R_1 \subset R$ be the k -algebra generated by $g^i R_0$ for all $1 \leq i \leq \mathrm{ord}(g)$. Then $R_1 \subset R$ is a finite-type k -algebra stable under g , with $R_1^{\mathrm{perf}} = A$, so $X_1 = \mathrm{Spec} R_1$ is a deperfection of X equipped with an automorphism g_1 deperfecting g ; since $X \rightarrow X_1$ is a homeomorphism, there is a unique g_1 -fixed point x_1 under x . Next, $g^* A \rightarrow A$ deperfects uniquely to a complex A_1 on X_1 equipped with a map $g_1^* A_1 \rightarrow A_1$, using the equivalence of categories $D(X_{\mathrm{et}}, \Lambda) \cong D(X_{1, \mathrm{et}}, \Lambda)$. Finally, we compute that

$$\mathrm{loc}_x(g, A) = \mathrm{loc}_{x_1}(g_1, A_1) = \mathrm{tr}(g_1|_{A_{1, x_1}}) = \mathrm{tr}(g|_{A_x})$$

where the first equality is formal nonsense (the six functors on k -varieties and on perfectly finite type k -schemes are compatible under perfection), the second equality is Varshavsky’s theorem, and the third equality is trivial. \square

6 Application to the Hecke stacks

In this final chapter we prove Theorem 1.0.2 by applying the technology of Lefschetz-Verdier to the Hecke stacks on Bun_G .

6.1 Bun_G , the local and global Hecke stacks, and their relation to shtuka spaces

Let F/\mathbf{Q}_p be a finite extension, and let G/F be a connected reductive group. Let k be an algebraically closed perfectoid field containing the residue field of F .

For an algebraically closed perfectoid field C/k , there is a bijection [Fara]

$$b \mapsto \mathcal{E}^b$$

between Kottwitz' set $B(G)$ and isomorphism classes of G -bundles on the Fargues-Fontaine curve X_C . Therefore the moduli stack of G -bundles is some geometric version of the set $B(G)$.

Definition 6.1.1 ([FS21, Definition III.0.1 and Theorem III.0.2]). Let Bun_G be the v-stack which assigns to a perfectoid space S/k the groupoid of G -bundles on X_S . Given a class $b \in B(G)$, let $i_b: \mathrm{Bun}_G^b \rightarrow \mathrm{Bun}_G$ be the locally closed substack classifying G -bundles which are isomorphic to \mathcal{E}^b at every geometric point.

Then Bun_G is a cohomologically smooth Artin v-stack over $\mathrm{Spd} k$ [FS21, Theorem I.4.1(vii)]. Central to its study are the substacks Bun_G^b . For each $b \in B(G)$ we have an isomorphism $\mathrm{Bun}_G^b \cong [\mathrm{Spd} k / \tilde{G}_b]$, where

$$\tilde{G}_b = \underline{\mathrm{Aut}} \mathcal{E}_b$$

is a group diamond over $\mathrm{Spd} k$. This fits in an exact sequence of group diamonds over $\mathrm{Spd} k$:

$$0 \rightarrow \tilde{G}_b^\circ \rightarrow \tilde{G}_b \rightarrow G_b(F)_{\mathrm{Spd} k} \rightarrow 0$$

Here, the neutral component $\tilde{G}_b^\circ \subset \tilde{G}_b$ is a cohomologically smooth group diamond over $\mathrm{Spd} k$, and G_b is the automorphism group of the isocrystal b . The group G_b is an inner form of a Levi subgroup of the quasisplit inner form of G . If b is basic, then i_b is an open immersion, and $\tilde{G}_b = G_b(F)_{\mathrm{Spd} k}$.

We next recall the Hecke correspondence on Bun_G and its relation to the local shtuka spaces $\mathrm{Sht}_{G,b,\mu}$. Since our main result on the cohomology of local Shimura varieties does not concern the action of a Weil group, all objects in this discussion will live over the base $S = \mathrm{Spd} C$, where C is an algebraically closed perfectoid field containing F , whose residue field contains k . In particular we have $\mathrm{Bun}_{G,C} = \mathrm{Bun}_G \times_{\mathrm{Spd} k} S$. If T is a perfectoid space over C , the Fargues-Fontaine curve X_T comes equipped with a degree 1 Cartier divisor D_T , corresponding to the untilt T of T^\flat .

We introduce now a diagram of v-stacks over $\mathrm{Spd} C$ containing both local and global Hecke correspondences:

$$\begin{array}{ccccc} \mathrm{Bun}_{G,C} & \xleftarrow{h_2} & \mathrm{Hecke}_{G,C} & \xrightarrow{h_1} & \mathrm{Bun}_{G,C} \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Bun}_{G,C}^{\mathrm{loc}} & \xleftarrow{h_2^{\mathrm{loc}}} & \mathrm{Hecke}_{G,C}^{\mathrm{loc}} & \xrightarrow{h_1^{\mathrm{loc}}} & \mathrm{Bun}_{G,C}^{\mathrm{loc}} \end{array} \quad (6.1.1)$$

in which all squares are cartesian, the morphisms labeled with i are open immersions, and the morphisms h_1 and h_2 are proper.

The top row of (6.1.2) can be described via the diagram:

$$\begin{array}{ccccc} [\mathrm{Gr}_{G,\leq\mu,C}^{1,\mathrm{adm}} / \underline{G(F)}] & \longrightarrow & [\mathrm{Gr}_{G,\leq\mu}^1 / \underline{G(F)}] & \longrightarrow & [*/\underline{G(F)}] \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ \mathrm{Hecke}_{\leq\mu,C}^{b,1} & \xrightarrow{i''_b} & \mathrm{Hecke}_{\leq\mu,C}^{*,1} & \xrightarrow{h_2^{*,1}} & \mathrm{Bun}_{G,C}^1. \end{array}$$

Explanation: $\mathrm{Gr}_{G,\leq\mu,C}^1$ assigns to $T = \mathrm{Spa}(R, R^+)$ the set of pairs (\mathcal{E}, f) , where \mathcal{E} is a G -bundle on X_S , and $f: \mathcal{E}^1|_{X_T \setminus D_T} \cong \mathcal{E}|_{X_T \setminus D_T}$ is an isomorphism, which is bounded by μ along D_T . The bundle \mathcal{E}^1 can be canonically trivialized over $\mathrm{Spa} B_{\mathrm{dR}}^+(R)$, and in so doing we obtain an isomorphism $\mathrm{Gr}_{G,\leq\mu,C}^1 \cong \mathrm{Gr}_{G,\leq\mu,C}$. Within $\mathrm{Gr}_{G,\leq\mu}^1$ we have the open locus $\mathrm{Gr}_{G,\leq\mu}^{1,\mathrm{adm}}$, consisting of those pairs (\mathcal{E}, γ) , where \mathcal{E} is everywhere isomorphic to \mathcal{E}^b .

Similarly the leftmost column of (6.1.2) can be described via the diagram:

$$\begin{array}{ccc} [\mathrm{Gr}_{G,\leq\mu,C}^{b,\mathrm{adm}} / \underline{G_b(F)}] & \xrightarrow{\cong} & \mathrm{Hecke}_{G,\leq\mu,C}^{b,1} \\ \downarrow & & \downarrow i'_1 \\ [\mathrm{Gr}_{G,\leq\mu,C}^b / \underline{J_b(F)}] & \xrightarrow{\cong} & \mathrm{Hecke}_{G,\leq\mu,C}^{b,*} \\ \downarrow & & \downarrow h_1^{b,*} \\ [\mathrm{Spd} C / \underline{G_b(F)}] & \xrightarrow{\cong} & \mathrm{Bun}_{G,C}^b. \end{array}$$

Explanation: $\mathrm{Gr}_{G,\leq\mu,C}^b$ assigns to $T = \mathrm{Spa}(R, R^+)$ the set of pairs (\mathcal{E}, f) , where \mathcal{E} is a G -bundle, and $f: \mathcal{E}|_{X_T \setminus D_T} \cong \mathcal{E}^b|_{X_T \setminus D_T}$ is an isomorphism, which is bounded by μ along D_T . We have an isomorphism $\mathrm{Gr}_{G,\leq\mu,C}^b \cong \mathrm{Gr}_{G,\leq-\mu,C}$. Within $\mathrm{Gr}_{G,\leq\mu,C}^b$, we have the open *admissible locus* $\mathrm{Gr}_{G,\leq\mu,C}^{b,\mathrm{adm}}$ consisting of pairs (\mathcal{E}, f) , where \mathcal{E} is everywhere isomorphic to \mathcal{E}^1 .

The moduli space of local shtukas $\mathrm{Sht}_{G,b,\mu,C}$ appears as the fiber product:

$$\begin{array}{ccc} \mathrm{Sht}_{G,b,\mu,C} & \longrightarrow & \mathrm{Spd} C \\ \downarrow & & \downarrow \\ \mathrm{Hecke}_{G,\leq\mu,C}^{b,1} & \xrightarrow{h_1^{b,1} \times h_2^{b,1}} & \mathrm{Bun}_{G,C}^b \times \mathrm{Bun}_{G,C}^1 \end{array} \quad (6.1.3)$$

where the right vertical morphism corresponds to $\mathcal{E}^b \times \mathcal{E}^1$. This is evident from the definition of $\mathrm{Sht}_{G,b,\mu}$: its S -points are morphisms $f: \mathcal{E}^1_{X_S \setminus D_S} \cong \mathcal{E}^b_{X_S \setminus D_S}$ which are bounded by μ on D_S .

We also have the period morphisms:

$$\begin{array}{ccc} & \mathrm{Sht}_{G,b,\mu,C} & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ \mathrm{Gr}_{G,\leq\mu,C}^b & & \mathrm{Gr}_{G,\leq\mu,C}^1 \end{array} \quad (6.1.4)$$

The morphism π_1 is a $G_b(F)_S$ -equivariant $G(F)_S$ -torsor over the admissible locus $\mathrm{Gr}_{G, \leq \mu, C}^{b, \mathrm{adm}}$. Similarly, π_2 is a $G(F)_S$ -equivariant $G_b(F)_S$ -torsor over the admissible locus $\mathrm{Gr}_{G, \leq \mu, C}^{1, \mathrm{adm}}$.

6.2 The inertia stack of the Hecke stack; admissibility of elliptic fixed points

We continue to put $S = \mathrm{Spd} C$. Here we investigate the inertia stack $\mathrm{In}_S(\mathrm{Hecke}_{G, \leq \mu, C})$, or at least the part of it lying over the strongly regular locus in $\mathrm{In}_S([S/G(F)_S]) \cong [G(F)_S // G(F)_S]$.

It will help to introduce some notation. Suppose \mathcal{E} is a G -bundle on X_C equipped with a trivialization at $\infty = D_C$. Let $T \subset G$ be a maximal torus. We have seen in Proposition 5.1.2 that there is a bijection $\lambda \mapsto L_\lambda$ between $X_*(T)$ and the set of T -fixed points of Gr_G . Given $\lambda \in X_*(T)$, we let $\mathcal{E}[\lambda]$ be the modification of \mathcal{E} corresponding to L_λ .

Lemma 6.2.1 ([CS17, Lemma 3.5.5], see also [CFS17, §2.2], but note that we use the opposite convention concerning Schubert cells). *Let \mathcal{E} be a G -bundle on X_C equipped with a trivialization at ∞ , let $T \subset G$ be a maximal torus, let $\lambda \in X_*(T)$ be a cocharacter, and let $\hat{\lambda} \in X^*(\hat{T})$ be the corresponding character. In the group $X^*(Z(\hat{G})^\Gamma)$ we have*

$$\kappa(\mathcal{E}[\lambda]) = \kappa(\mathcal{E}) + \hat{\lambda}|_{Z(\hat{G})^\Gamma}.$$

Proposition 6.2.2. *Suppose a pair $(g, g') \in G(F)_{\mathrm{sr}} \times G_b(F)_{\mathrm{sr}}$ fixes a point $x \in \mathrm{Sht}_{G, b, \mu}(C)$. Let $T = \mathrm{Cent}(g, G)$ and $T' = \mathrm{Cent}(g', G_b)$. Then $\pi_1(x) \in \mathrm{Gr}_{G, \leq -\mu}^{g'}$ and $\pi_2(x) \in \mathrm{Gr}_{G, \leq \mu}^g$ correspond to cocharacters $\lambda' \in X_*(T')_{\leq -\mu}$ and $\lambda \in X_*(T)_{\leq \mu}$, respectively.*

There exists $y \in G(\check{F})$ such that $\mathrm{ad} y$ is an F -rational isomorphism $T \rightarrow T'$, which carries g to g' and λ onto $-\lambda'$. The invariant $\mathrm{inv}[b](g, g') \in B(T) \cong X_(T)_\Gamma$ agrees with the image of λ under $X_*(T) \rightarrow X_*(T)_\Gamma$. Therefore (g, g', λ) lies in $\mathrm{Rel}_{b, \mu}$.*

Proof. The point x corresponds to an isomorphism $\gamma: \mathcal{E}^1[\lambda] \rightarrow \mathcal{E}^b$, and also to an isomorphism $\gamma': \mathcal{E}^1 \rightarrow \mathcal{E}^b[\lambda']$. Each of these interlaces the action of g with g' , and furthermore $\gamma = \gamma'$ away from ∞ . Trivializing \mathcal{E}^1 and \mathcal{E}^b away from ∞ , we see that g and g' become conjugate over the ring $B_e = H^0(X_C \setminus \{\infty\}, \mathcal{O}_{X_C})$, which implies they are conjugate over \bar{F} and (by Lemma 3.2.1) they are even conjugate over \check{F} . Let $y \in G(\check{F})$ be an element such that $(\mathrm{ad} y)(g) = g'$. Then $\mathrm{ad} y$ is a \check{F} -rational isomorphism $T \rightarrow T'$ which carries g onto g' . In fact, since there is only one such isomorphism, we can conclude that $\mathrm{ad} y$ is an F -rational isomorphism $T \rightarrow T'$. Let $\lambda_0 = (\mathrm{ad} y^{-1})(\lambda') \in X_*(T)$.

Let $b_0 = y^{-1} b y^\sigma$. Then (cf. Definition 3.2.2) we have $b_0 \in T(F)$. The element y induces isomorphisms $y: \mathcal{E}^{b_0} \rightarrow \mathcal{E}^b$ and $y: \mathcal{E}^{b_0}[\lambda_0] \rightarrow \mathcal{E}^b[\lambda']$. Then the isomorphism $y^{-1} \gamma': \mathcal{E}^1 \rightarrow \mathcal{E}^{b_0}[\lambda_0]$ descends to an isomorphism of T -bundles; comparing this with the isomorphism $\gamma y^{-1}: \mathcal{E}^1[\lambda] \rightarrow \mathcal{E}^{b_0}$ shows that $\lambda_0 = -\lambda$. In light of the isomorphism of T -bundles $\mathcal{E}^1[\lambda] \cong \mathcal{E}^{b_0}$, Lemma 6.2.1 implies that the identity $\kappa(\mathcal{E}^{b_0}) = \lambda$ holds in $B(T)$. But also $\kappa(\mathcal{E}^{b_0})$ is the class of $[b_0]$ in $B(T)$, which is $\mathrm{inv}[b](g, g')$ by definition. \square

Proposition 6.2.2 shows that if (g, g') fixes a point of $\mathrm{Sht}_{G, b, \mu}$ and g' is strongly regular, then so is g , and then they are related. However the converse may fail: if a pair of related strongly regular elements (g, g') is given, it is not necessarily true that (g, g') fixes a point of $\mathrm{Sht}_{G, b, \mu}$. Indeed, a

necessary condition for this is that the action of g' on $\mathrm{Gr}_{G,\leq\mu}^b$ has a fixed point in the admissible locus, and this is not automatic.

This result is always true, however, if g (or equivalently, g') lies in the elliptic locus.

Theorem 6.2.3. *Let $g \in G(F)_{\mathrm{ell}}$. Then fixed points of g in $\mathrm{Gr}_{G,\leq\mu}^1$ lie in the admissible locus $\mathrm{Gr}_{G,\leq\mu}^{1,\mathrm{adm}}$. Similarly, if $g' \in G_b(F)_{\mathrm{ell}}$, then fixed points of g' in $\mathrm{Gr}_{G,\leq\mu}^b$ lie in the admissible locus $\mathrm{Gr}_{G,\leq\mu}^{b,\mathrm{adm}}$.*

Proof. We prove the first statement; the second is similar. Let $g \in G(F)_{\mathrm{ell}}$, and let $T = \mathrm{Cent}(g, G)$ be the elliptic maximal torus containing g . Suppose we are given a g -fixed point $x \in \mathrm{Gr}_{G,\leq\mu}(C)$. Then x corresponds to a cocharacter $\lambda \in X_*(T)$, which in turn corresponds to a modification $\mathcal{E}^1[\lambda]$ of the trivial G -bundle \mathcal{E}^1 . We wish to show that $\mathcal{E}^1[\lambda] \cong \mathcal{E}^b$. First we will show that it is semistable.

Let $b' \in G(\check{F})$ be an element whose class in $B(G)$ corresponds to the isomorphism class of $\mathcal{E}^1[\lambda]$. We wish to show that b' is basic. We have the algebraic group $G_{b'}/F$, which is a priori an inner form of a Levi subgroup M^* of G^* , where G^* is the quasi-split inner form of G . Showing that b' is basic is equivalent to showing that $M^* = G^*$.

We have an isomorphism $\gamma: \mathcal{E}^1[\lambda] \cong \mathcal{E}^{b'}$. The action of $g \in T(F)$ on \mathcal{E}^1 extends to an action on $\mathcal{E}^1[\lambda]$, which can be transported via γ to obtain an automorphism $g' \in \tilde{G}_{b'}(C) = \mathrm{Aut} \mathcal{E}_{b'}$. Let \bar{g}' be the image of g' under the projection $\tilde{G}_{b'}(C) \rightarrow G_{b'}(F)$.

The G -bundles \mathcal{E}^1 and $\mathcal{E}^{b'}$ may be trivialized over $\mathrm{Spec} B_{\mathrm{dR}}^+(C)$. In doing so, we obtain embeddings of $G(F) = \mathrm{Aut} \mathcal{E}^1$ and $\tilde{G}_{b'}(C) = \mathrm{Aut} \mathcal{E}^{b'}$ into $G(B_{\mathrm{dR}}^+(C))$; we denote both of these by $h \mapsto h_\infty$. We also have the isomorphism γ_∞ between \mathcal{E}^1 and $\mathcal{E}^{b'}$ over $\mathrm{Spec} B_{\mathrm{dR}}(C)$; we may identify γ_∞ with an element of $G(B_{\mathrm{dR}}(C))$, and then $g'_\infty = \gamma_\infty g_\infty \gamma_\infty^{-1}$ holds in $G(B_{\mathrm{dR}}(C))$.

The element \bar{g}'_∞ is conjugate to g'_∞ , so \bar{g}'_∞ is conjugate to g_∞ in $G(B_{\mathrm{dR}})$. Since g and \bar{g}' are both regular semisimple \bar{F} -points of G , being conjugate in $G(B_{\mathrm{dR}})$ is the same as being conjugate in $G(\bar{F})$. Their centralizers, being F -rational tori, are thus isomorphic over F . Thus $G_{b'}$ contains a maximal torus that is elliptic for G . Elliptic maximal tori transfer across inner forms [Kot86, §10], which means that the Levi subgroup $M^* \subset G^*$ of which $G_{b'}$ is an inner form contains a maximal torus that is elliptic for G^* . Therefore $M^* = G^*$.

We have shown that $\mathcal{E}^1[\lambda] \cong \mathcal{E}^{b'}$ is semistable, implying that $\mathrm{Aut} \mathcal{E}^{b'} = G_{b'}(F)$ and that $g' \in G_{b'}(F)$. Lemma 6.2.1 shows that $\kappa([b'])$ equals the image of λ in $\pi_1(G)_\Gamma$; this is the same as the image of μ , which in turn is the same as $\kappa([b])$ because $b \in B(G, \mu)$. Since b' is basic, we have $[b'] = [b]$ by [Kot85, Proposition 5.6]. \square

Recall the locally profinite set $\mathrm{Rel}_{b,\leq\mu}$ from Definition 3.2.4. This is the set of conjugacy classes of triples (g, g', λ) , where $g \in G(F)$ and $g' \in G_b(F)$ are related strongly regular elements, and λ is a cocharacter of $T = \mathrm{Cent}(g, G)$, bounded by μ , such that $\kappa(\mathrm{inv}[b](g, g'))$ agrees with the image of λ in $X_*(T)_\Gamma$. Let $\mathrm{Rel}_{b,\leq\mu,\mathrm{ell}}$ be the subset where g (equivalently, g') is elliptic.

Theorem 6.2.3 has the following corollary. For a v-stack X , we write $|X|$ for the underlying topological space.

Corollary 6.2.4. *Let $\text{In}_S(\text{Hecke}_{G,\leq\mu,S}^{b,*})_{\text{ell}}$ be the preimage under $\text{In}_S(h_1)$ of $\text{In}_S(\text{Bun}_{G,S}^b)_{\text{ell}}$. Similarly let $\text{In}_S(\text{Hecke}_{G,\leq\mu,S}^{*,1})_{\text{ell}}$ be the preimage under $\text{In}_S(h_2)$ of $\text{In}_S(\text{Bun}_{G,S}^1)_{\text{ell}}$. Then*

$$\text{In}_S(\text{Hecke}_{G,\leq\mu,S}^{b,*})_{\text{ell}} = \text{In}_S(\text{Hecke}_{G,\leq\mu,S}^{*,1})_{\text{ell}} = \text{In}_S(\text{Hecke}_{G,\leq\mu,S}^{b,1})_{\text{ell}}.$$

There is a homeomorphism $\left| \text{In}_S(\text{Hecke}_{G,b,\leq\mu,S}^{b,1})_{\text{ell}} \right| \cong \text{Rel}_{b,\mu,\text{ell}}$.

Proof. The first claim is just the statement that fixed points of elliptic elements on Gr_G^b and Gr_G^1 are admissible. For the second claim: since $\text{Hecke}_{G,\leq\mu,S}^{*,1} \cong [\text{Gr}_{G,\leq\mu,S} / G(F)_S]$, we can think of $\left| \text{In}_S(\text{Hecke}_{G,b,\leq\mu,S}^{b,1})_{\text{ell}} \right|$ as the set of conjugacy classes of pairs (g, λ) , where $g \in G(F)_{\text{ell}}$ and $\lambda \in X_*(T)_{\leq\mu}$, where $T = \text{Cent}(g, G)$. We have an isomorphism $\mathcal{E}^1[\lambda] \cong \mathcal{E}^b$. The element $g \in G(F) \cong \text{Aut } \mathcal{E}^1$ determines an element $g' \in G_b(F) \cong \text{Aut } \mathcal{E}^b$, up to conjugacy. By Proposition 6.2.2, the triple (g, g', λ) determines an element of $\text{Rel}_{b,\mu,\text{ell}}$. Conversely, given such a triple (g, g', λ) , the pair (g, λ) determines an element $g'' \in G_b(F)$ as we have just argued, but then g' and g'' are conjugate by Remark 3.2.5. \square

6.3 Transfer of distributions from G_b to G

We continue to let b be a basic element of $B(G)$. Let Λ be a ring in which p is invertible. Recall the Hecke operator

$$T_{b,\mu}^{G_b \rightarrow G} : C(G_b(F)_{\text{sr}}, \Lambda) \rightarrow C(G(F)_{\text{sr}}, \Lambda)$$

from 3.2.7. As promised, we can now promote this to a transfer of distributions, at least after restriction to elliptic loci (and dependent on choices of Λ -valued Haar measures on $G(F)$ and $G_b(F)$).

Recall the period morphisms:

$$\text{Gr}_{G,\leq\mu,C}^b \xleftarrow{\pi_1} \text{Sht}_{G,b,\mu,C} \xrightarrow{\pi_2} \text{Gr}_{G,\leq\mu,C}^1,$$

in which π_1 is a $G(F)_S$ -torsor over its image, and π_2 is a $G_b(F)_S$ -torsor over its image. Consider the action map on $\text{Sht}_{G,b,\mu,C}$:

$$\alpha_{\text{Sht}} : G(F)_S \times G_b(F)_S \times \text{Sht}_{G,b,\mu,C} \rightarrow \text{Sht}_{G,b,\mu,C}$$

and also those on the period domains:

$$\begin{aligned} \alpha_1 : G(F)_S \times \text{Gr}_{G,\leq\mu,C}^1 &\rightarrow \text{Gr}_{G,\leq\mu,C}^1 \\ \alpha_b : G_b(F)_S \times \text{Gr}_{G,\leq\mu,C}^b &\rightarrow \text{Gr}_{G,\leq\mu,C}^b \end{aligned}$$

For $? \in \{\text{Sht}, 1, b\}$ we can define the elliptic fixed-point locus $\text{Fix}(\alpha?)_{\text{ell}}$ of the corresponding action map, consisting of pairs (g, x) with g elliptic and $g.x = x$; let us think of each $\text{Fix}(\alpha?)$ as a locally profinite set. For instance, $\text{Fix}(\alpha_1)$ is the set of pairs (g, λ) , where $g \in G(F)_{\text{ell}}$, and $\lambda \in X_*(T_g)$ ($T_g = \text{Cent}(g, G)$) is bounded by μ . These fit into a diagram

$$\begin{array}{ccc} \text{Fix}(\alpha_b)_{\text{ell}} & \xleftarrow{p_1} \text{Fix}(\alpha_{\text{Sht}})_{\text{ell}} \xrightarrow{p_2} & \text{Fix}(\alpha_1)_{\text{ell}} \\ q_1 \downarrow & & \downarrow q_2 \\ G_b(F)_{\text{ell}} & & G(F)_{\text{ell}} \end{array} \quad (6.3.1)$$

of locally profinite sets, in which p_1 is a $G_b(F)$ -equivariant $G(F)$ -torsor, p_2 is a $G(F)$ -equivariant $G_b(F)$ -torsor, and q_1 and q_2 are finite étale. (The maps p_i are surjective by Theorem 6.2.3.)

To define our transfer of distributions we need a lemma.

Lemma 6.3.1. *Let H be a locally pro- p group, and let Λ be a commutative ring in which p is invertible. Choose a Λ -valued Haar measure on H . Let $h: \tilde{T} \rightarrow T$ be an H -torsor in locally profinite sets. The integration-along-fibers map $C_c(\tilde{T}, \Lambda) \rightarrow C_c(T, \Lambda)$ induces an isomorphism of $C(T, \Lambda)$ -modules*

$$h_*: C_c(\tilde{T}, \Lambda)_H \rightarrow C_c(T, \Lambda)$$

and, dually, an isomorphism of $C(T, \Lambda)$ -modules

$$h_*: \text{Dist}(\tilde{T}, \Lambda)^H \rightarrow \text{Dist}(T, \Lambda).$$

Proof. The $C(T, \Lambda)$ -modules $C_c(T, \Lambda)$ and $C_c(\tilde{T}, \Lambda)$ are smooth in the sense of Definition B.2.1. Therefore by Lemma B.2.5 the statement is local on T , so we may assume that the torsor $\tilde{T} = T \times H$ is split. Then $C_c(\tilde{T}, \Lambda)_H = C_c(T, \Lambda) \otimes_{\Lambda} C_c(H, \Lambda)_H$. The integration map $C_c(H, \Lambda)_H \rightarrow \Lambda$ is an isomorphism, so that $C_c(\tilde{T}, \Lambda)_H \cong C_c(T, \Lambda)$. \square

Definition 6.3.2. Choose Λ -valued Haar measures dg and dg' on $G(F)$ and on $G_b(F)$. Then, with notation as in (6.3.1), we define a Λ -linear map

$$\tilde{T}_{b,\mu}^{G \rightarrow G_b}: C_c(G(F)_{\text{ell}}, \Lambda)_{G(F)} \rightarrow C_c(G_b(F)_{\text{ell}}, \Lambda)_{G_b(F)} \quad (6.3.2)$$

as the composition

$$\begin{aligned} C_c(G(F)_{\text{ell}}, \Lambda)_{G(F)} &\xrightarrow{q_2^*} C_c(\text{Fix}(\alpha_1)_{\text{ell}}, \Lambda)_{G(F)} \\ &\xrightarrow{(p_2)_*^{-1}} C_c(\text{Fix}(\alpha_{\text{Sht}})_{\text{ell}}, \Lambda)_{G(F) \times G_b(F)} \\ &\xrightarrow{(p_1)_*} C_c(\text{Fix}(\alpha_b)_{\text{ell}}, \Lambda)_{G_b(F)} \\ &\xrightarrow{\cdot K_\mu} C_c(\text{Fix}(\alpha_b)_{\text{ell}}, \Lambda)_{G_b(F)} \\ &\xrightarrow{q_1^*} C_c(G_b(F)_{\text{ell}}, \Lambda)_{G_b(F)}, \end{aligned}$$

where q_2^* means pullback, q_1^* means pushforward (*i.e.*, sum over fibers), the isomorphisms $(p_i)_*$ are induced by our choices of Haar measures as in Lemma 6.3.1, and finally $K_\mu \in C(\text{Fix}(\alpha_b), \Lambda)^{G_b(F)}$ is the function $(g', \lambda') \mapsto \text{rank } V_\mu^\vee[\lambda']$.

Again, we emphasize that this definition depends on the specific choice of Haar measures dg and dg' ; rescaling these choices will rescale $\tilde{T}_{b,\mu}^{G \rightarrow G_b}$.

Proposition 6.3.3. *Assume that $\Lambda = \overline{\mathbf{Q}}_\ell$ and that the Haar measures dg and dg' are compatible in the sense of §3.3. Let $\phi \in C_c(G(F)_{\text{ell}}, \Lambda)$, and let $\phi' \in C_c(G_b(F)_{\text{ell}}, \Lambda)$ be any lift of $\tilde{T}_{b,\mu}^{G \rightarrow G_b} \phi$. Then the orbital integrals of ϕ and ϕ' are related by*

$$\phi'_{G_b} = T_{b,\mu}^{G \rightarrow G_b} \phi_G.$$

Proof. For $g' \in G_b(F)_{\text{ell}}$, with centralizer T' , we have:

$$\begin{aligned} & \phi'_{G_b}(g') \\ &= \int_{h' \in G_b(F)/T'(F)} \phi'(h'g'(h')^{-1}) dh' \\ &= \sum_{\lambda' \in X_*(T')_{\leq -\mu}} \text{rank } V_\mu^\vee[\lambda'] \int_{h' \in \frac{G_b(F)}{T'(F)}} [(p_1)_*(p_2)_*^{-1} q_2^* \phi](h'.(g', \lambda')) dh dh'. \end{aligned}$$

Let T be a transfer of the elliptic torus T' to G . Since $\text{Fix}(\alpha_{\text{Sht}})_{\text{ell}} \rightarrow \text{Fix}(\alpha_b)_{\text{ell}}$ is a $G(F)$ -torsor, we may choose for each λ' a lift $y_{\lambda'} = (g_{\lambda'}, g', x_{\lambda'})$ of (g', λ') to $\text{Fix}(\alpha_{\text{Sht}})$ with $g_{\lambda'} \in T(F)$. Then $\phi'_{G_b}(g')$ equals

$$\sum_{\lambda' \in X_*(T')_{\leq -\mu}} \text{rank } V_\mu^\vee[\lambda'] \int_{h' \in G_b(F)/T'(F)} \int_{h \in G(F)} [(p_2)_*^{-1} q_2^* \phi]((h, h').(y_{\lambda'})) dh dh'$$

We rewrite the inner integral as a nested integral; so that our expression for $\phi'_{G_b}(g')$ equals:

$$\begin{aligned} & \sum_{\lambda'} \text{rank } V_\mu^\vee[\lambda'] \int_{h' \in G_b(F)/T'(F)} \int_{h \in G(F)/T(F)} \int_{t \in T(F)} [(p_2)_*^{-1} q_2^* \phi]((ht^{-1}, h').y_{\lambda'}) dt dh dh' \\ &= \sum_{\lambda'} \text{rank } V_\mu^\vee[\lambda'] \int_{h' \in G_b(F)/T'(F)} \int_{h \in G(F)/T(F)} \int_{t' \in T'(F)} [(p_2)_*^{-1} q_2^* \phi]((h, h't').y_{\lambda'}) dt' dh dh' \end{aligned}$$

Here we have used Proposition 6.2.2: there is an isomorphism $\iota: t \mapsto t'$ between $T(F)$ and $T'(F)$ satisfying $(t, t').y_{\lambda'} = y_{\lambda'}$. This induces a bijection $\lambda \mapsto \lambda' = -\iota_* \lambda$ between $X_*(T)_{\leq \mu}$ and $X_*(T')_{\leq -\mu}$. Given $\lambda \in X_*(T)_{\leq \mu}$, we let $g_\lambda = g_{\lambda'}$. Then by Proposition 6.2.2, the preimage of g' in $\text{Rel}_{b, \mu}$ is exactly $\{(g_\lambda, g, \lambda)\}_{\lambda \in X_*(T)_{\leq \mu}}$.

Noting that $\text{rank } V_\mu[\lambda] = \text{rank } V_\mu^\vee[\lambda']$, we exchange the order of the first two integrals above to obtain

$$\begin{aligned} \phi'_{G_b}(g') &= \sum_{\lambda \in X_*(T)_{\leq \mu}} \text{rank } V_\mu[\lambda] \int_{h \in G(F)/T(F)} \int_{h' \in G_b(F)} [(p_2)_*^{-1} q_2^* \phi]((h, h') \cdot y_{\lambda'}) dh' dh \\ &= \sum_{\lambda \in X_*(T)_{\leq \mu}} \text{rank } V_\mu[\lambda] \int_{h \in G(F)/T(F)} \phi(hg_\lambda h^{-1}) dh \\ &= \sum_{\lambda \in X_*(T)_{\leq \mu}} \text{rank } V_\mu[\lambda] \phi_G(g_\lambda) \\ &= [T_{b, \mu}^{G \rightarrow G_b} \phi_G](g'). \end{aligned}$$

□

Definition 6.3.4. Let

$$\mathcal{T}_{b, \mu}^{G_b \rightarrow G}: \text{Dist}(G_b(F)_{\text{ell}}, \Lambda)^{G_b(F)} \rightarrow \text{Dist}(G(F)_{\text{ell}}, \Lambda)^{G(F)}$$

be the Λ -linear dual of $\tilde{T}_{b, \mu}^{G \rightarrow G_b}$. Again, note that this map depends on the Haar measures dg and dg' .

Proposition 6.3.5. *Assume that $\Lambda = \overline{\mathbf{Q}_\ell}$ and that the Haar measures dg and dg' are compatible in the sense of §3.3. Then the transfer of distributions $\mathcal{T}_{b,\mu}^{G_b \rightarrow G}$ extends the transfer of functions $T_{b,\mu}^{G_b \rightarrow G}$ from Definition 3.2.7.*

Proof. Let $f \in C(G_b(F)_{\text{ell}}, \Lambda)^{G_b(F)}$ be a conjugation-invariant function. Let $\phi \in C_c(G(F)_{\text{ell}}, \Lambda)$, and let $\phi' \in C_c(G_b(F)_{\text{ell}}, \Lambda)$ be a lift of $\tilde{T}_{b,\mu}^{G \rightarrow G_b} \phi$. Using the Weyl integration formula (3.3.1), Lemma 3.3.1, and Proposition 6.3.3 we compute

$$\begin{aligned} \int_{g \in G(F)_{\text{ell}}} \phi(g) \mathcal{T}_{b,\mu}^{G_b \rightarrow G}(f dg') &= \int_{g' \in G_b(F)_{\text{ell}}} f(g') \phi'(g') dg' \\ &= \langle f, \phi'_{G_b} \rangle_{G_b} \\ &= \langle f, T_{b,\mu}^{G \rightarrow G_b} \phi_G \rangle_{G_b} \\ &= \langle T_{b,\mu}^{G_b \rightarrow G} f, \phi_G \rangle_G \\ &= \int_{g \in G(F)_{\text{ell}}} \phi(g) (T_{b,\mu}^{G_b \rightarrow G} f) dg, \end{aligned}$$

so that

$$\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(f dg') = T_{b,\mu}^{G_b \rightarrow G}(f) dg$$

as desired. \square

6.4 Hecke operators on Bun_G and the cohomology of shtuka spaces

We are finally ready to reap our rewards. For the remainder of this chapter, we fix a prime $\ell \neq p$, and write Λ for a \mathbf{Z}_ℓ -algebra. Let \widehat{G} be the Langlands dual group over \mathbf{Z}_ℓ .

We begin by quickly reviewing the results of [FS21] on the categories $D_{\text{lis}}(\text{Bun}_G, \Lambda)$ and $D_{\text{lis}}(\text{Bun}_G^b, \Lambda)$, and the action of Hecke operators on $D_{\text{lis}}(\text{Bun}_G, \Lambda)$.

The first key fact is that for any $b \in B(G)$, there is a natural equivalence of categories

$$D(G_b(F), \Lambda) \cong D_{\text{lis}}(\text{Bun}_G^b, \Lambda) \tag{6.4.1}$$

[FS21, Theorem I.5.1]. For a complex ρ of smooth representations of $G_b(F)$, we will slightly abusively also write ρ for the corresponding object of $D_{\text{lis}}(\text{Bun}_G^b, \Lambda)$.

Next, recall that there is a notion of ULA objects in $D_{\text{lis}}(\text{Bun}_G, \Lambda)$. These admit the following concrete characterization.

Theorem 6.4.1 ([FS21, Theorem I.5.1(v)]). *The following are equivalent for an object $A \in D_{\text{lis}}(\text{Bun}_G, \Lambda)$.*

1. A is ULA over $\text{Spd } k$.
2. For all $b \in B(G)$, the restriction $i_b^* A$, considered as an object of $D(G_b(F), \Lambda)$ via (6.4.1), is admissible in the sense that $(i_b^* A)^K$ is a perfect complex for all pro- p open subgroups $K \subset G_b(F)$.

Moreover, ULA objects are preserved under Verdier duality $\mathbf{D} = \mathbf{D}_{\mathrm{Bun}_G/\mathrm{Spd}k}$, and satisfy Verdier biduality.

Corollary 6.4.2. *Let $b \in B(G)$, and let ρ be an admissible complex in $D(G_b(F), \Lambda)$. The objects $(i_b)_*\rho$ and $(i_b)_!\rho$ of $D_{\mathrm{lis}}(\mathrm{Bun}_G, \Lambda)$ are ULA over $\mathrm{Spd}k$.*

Proof. The object $(i_b)_!\rho$ is clearly ULA by to the criterion in Theorem 6.4.1. Using Verdier duality (P4.) we have $\mathbf{D}((i_b)_!\rho^\vee) \cong (i_b)_*\rho$, so that $(i_b)_*\rho$ is also ULA. \square

Next, recall that any object V of $\mathrm{Rep}_{\widehat{G}}(\Lambda)$ gives rise to a Hecke operator T_V , which is an endofunctor of $D_{\mathrm{lis}}(\mathrm{Bun}_{G,C}, \Lambda)$. When Λ is a torsion ring, there is a natural equivalence $D_{\mathrm{lis}}(\mathrm{Bun}_{G,C}, \Lambda) \cong D_{\mathrm{ét}}(\mathrm{Bun}_{G,C}, \Lambda)$, and the operator T_V is defined concretely as the operation

$$\begin{aligned} T_V: D_{\mathrm{ét}}(\mathrm{Bun}_{G,C}, \Lambda) &\rightarrow D_{\mathrm{ét}}(\mathrm{Bun}_{G,C}, \Lambda) \\ \mathcal{F} &\mapsto h_{2!}(h_1^*\mathcal{F} \otimes \mathcal{S}_V). \end{aligned}$$

Here $\mathcal{S}_V \in D_{\mathrm{ét}}(\mathrm{Hecke}_{G,C}, \Lambda)$ is pulled back from the object $\mathcal{S}_V \in D_{\mathrm{ét}}(\mathrm{Hecke}_{G,C}^{\mathrm{loc}}, \Lambda)$ corresponding to V under the Satake equivalence (Theorem 5.1.1).

Theorem 6.4.3 ([FS21, Theorem IX.0.1]). *The Hecke operators preserve the subcategories of ULA and compact objects in $D_{\mathrm{lis}}(\mathrm{Bun}_{G,C}, \Lambda)$. For any V , T_V has left and right adjoint given by T_{V^\vee} , where V^\vee is the dual representation of \widehat{G} . The actions of Hecke operators are compatible with extension of scalars along any ring map $\Lambda \rightarrow \Lambda'$.*

Next, we explain the relation between the Hecke operators T_V and the cohomology of local shtuka spaces. Let μ be a dominant cocharacter of G , and let $V_\mu \in \mathrm{Rep}(\widehat{G})$ be the associated Weyl module. For any \mathbf{Z}_ℓ -algebra Λ , we write $V_{\mu,\Lambda} \in \mathrm{Rep}(\widehat{G}_\Lambda)$ for the base change of V_μ . Let $\mathcal{S}_\mu = \mathcal{S}_{V_\mu}$ be the corresponding object in the Satake category with \mathbf{Z}_ℓ -coefficients; similarly, if Λ is a torsion ring we write $\mathcal{S}_{\mu,\Lambda} = \mathcal{S}_{V_{\mu,\Lambda}}$ for the corresponding object with Λ -coefficients. We will slightly abuse notation by using the same notations for the pullbacks of \mathcal{S}_μ and $\mathcal{S}_{\mu,\Lambda}$ to various other v-stacks, including $\mathrm{Gr}_{G,\leq\mu,C}$ and $\mathrm{Sht}_{G,b,\mu,C}$ (along the period morphism π_1 from (6.1.4)).

Lemma 6.4.4. *Let $K \subset G(F)$ be an open compact subgroup, and let $I_K = \mathrm{cInd}_K^{G(F)} \mathbf{Z}_\ell$, where cInd is compactly supported induction. Then there is a natural isomorphism*

$$R\Gamma_c(\mathrm{Sht}_{G,b,\mu,K,C}, \mathcal{S}_\mu) \cong i_b^* T_{V_\mu}(i_1)_! I_K$$

in $D(G_b(F), \mathbf{Z}_\ell)$.

Proof. The global sections of $i_b^* T_{V_\mu}(i_1)_! I_K$ over $\mathrm{Spd}C \rightarrow \mathrm{Bun}_G^b$ are

$$R\Gamma(\mathrm{Spd}C, i_b^* h_{2!}(h_1^*(i_1)_! I_K \otimes \mathcal{S}_\mu)) \cong R\Gamma_c(\mathrm{Gr}^{b,1}, I_K|_{\mathrm{Gr}^{b,1}} \otimes \mathcal{S}_\mu).$$

Now use $I_K \cong (j_K)_! \Lambda$ along with proper base change to get the result. \square

Recall that when ρ is any smooth $G_b(F)$ -representation with $\overline{\mathbf{Q}_\ell}$ -coefficients, we defined an object

$$R\Gamma(G, b, \mu)[\rho] \cong \varinjlim_K R\mathrm{Hom}_{G_b(F)}(R\Gamma_c(\mathrm{Sht}_{(G,b,\mu),C}/K, \mathcal{S}_\mu) \otimes \overline{\mathbf{Q}_\ell}, \rho),$$

in $D(G(F), \overline{\mathbf{Q}_\ell})$, cf. Definition 2.4.3. The association $\rho \mapsto R\Gamma(G, b, \mu)[\rho]$ clearly extends to a functor $D(G_b(F), \overline{\mathbf{Q}_\ell}) \rightarrow D(G(F), \overline{\mathbf{Q}_\ell})$. Our next goal is to give an alternative approach to this construction, which is valid for more general coefficient rings and which makes the finiteness properties of this construction transparent.

Proposition 6.4.5. *Let ρ be any object of $D(G_b(F), \overline{\mathbf{Q}_\ell})$. Then there is a natural isomorphism*

$$R\Gamma(G, b, \mu)[\rho] \cong i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} (i_b)_* \rho$$

in $D(G(F), \overline{\mathbf{Q}_\ell})$.

If ρ is admissible, then so is $R\Gamma(G, b, \mu)[\rho]$. If ρ is of finite length, then so is $R\Gamma(G, b, \mu)[\rho]$.

Proof. Let $K \subset G(F)$ be a compact open subgroup. Using Lemma 6.4.4, various adjunctions, and the compatibility of Hecke operators with extension of scalars, we have

$$\begin{aligned} R\mathrm{Hom}_{G_b(F)}(R\Gamma_c(\mathrm{Sht}_{G,b,\mu,C}, \mathcal{S}_V) \otimes \overline{\mathbf{Q}_\ell}, \rho) &\cong R\mathrm{Hom}_{G_b(F)}(i_b^* T_{V_\mu} (i_1)! \mathrm{cInd}_K^{G(F)} \mathbf{Z}_\ell \otimes \overline{\mathbf{Q}_\ell}, \rho) \\ &\cong R\mathrm{Hom}_{G_b(F)}(i_b^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}} (i_1)! \mathrm{cInd}_K^{G(F)} \overline{\mathbf{Q}_\ell}, \rho) \\ &\cong (i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} (i_b)_* \rho)^K. \end{aligned}$$

Taking the colimit over K gives the first claim. The claim about preservation of admissibility now follows from Theorem 6.4.1 combined with Theorem 6.4.3. For the final claim, fix some ρ of finite length. Note that $i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} (i_b)_* \rho$ is the smooth dual of $i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}} (i_b)! \rho^\vee$, so it's enough to show that $i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}} (i_b)! \rho^\vee$ is of finite length. But finite length is equivalent to being both compact and admissible, so we conclude by observing that the operation $i_1^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}} (i_b)! (-)$ preserves compact objects. \square

Definition 6.4.6. For any \mathbf{Z}_ℓ -algebra Λ , we write

$$R\Gamma(G, b, \mu)[-] : D(G_b(F), \Lambda) \rightarrow D(G(F), \Lambda)$$

for the functor $i_1^* T_{V_{\mu, \Lambda}^\vee} (i_b)_* (-)$.

By the previous discussion, this functor is compatible with extension of scalars along any map $\Lambda \rightarrow \Lambda'$, and preserves admissible objects. Moreover, if Λ is Artinian and $\rho \in D(G_b(F), \Lambda)$ is admissible of finite length, then $R\Gamma(G, b, \mu)[\rho]$ is also of finite length, by the same argument as in the proof of Proposition 6.4.5.

We now come to the technical heart of this paper. Choose \mathbf{Z}_ℓ -valued Haar measures on $G(F)$ and $G_b(F)$, inducing Λ -valued Haar measures on the same groups compatibly with varying Λ . Then for any Λ , any admissible representation π of $G(F)$ with coefficients in Λ has a corresponding Λ -valued trace distribution $\mathrm{tr. dist}(\pi)$, and similarly for $G_b(F)$. Recall also that we defined a transfer of Λ -valued distributions $\mathcal{T}_{b,\mu}^{G_b \rightarrow G}$, Definition 6.3.4, which also depends on these choices of Haar measures.

Proposition 6.4.7. *Let Λ be any torsion \mathbf{Z}_ℓ -algebra, and let ρ be any admissible representation of $G_b(F)$ with coefficients in Λ . Then have an equality*

$$\mathrm{tr. dist} R\Gamma(G, b, \mu)[\rho]_{\mathrm{ell}} = \mathcal{T}_{b, \mu}^{G_b \rightarrow G} \mathrm{tr. dist}(\rho)_{\mathrm{ell}}$$

in $\mathrm{Dist}(G(F)_{\mathrm{ell}}, \Lambda)^{G(F)}$.

As a sanity check, note that all dependencies on the choices of Haar measures are compatible: rescaling dg and dg' by C and C' , we see that $\mathrm{tr. dist} R\Gamma(G, b, \mu)[\rho]$ scales by C , $\mathrm{tr. dist}(\rho)$ scales by C' , and $\mathcal{T}_{b, \mu}^{G_b \rightarrow G}$ scales by C/C' .

Proof. In the following proof, we set $S = \mathrm{Spd} C$ and $V = V_{\mu, \Lambda}$ for brevity.

We have an isomorphism

$$H^0(\mathrm{In}_S(\mathrm{Bun}_{G, C}^1), K_{\mathrm{In}_S(\mathrm{Bun}_{G, C}^1)/S}) \cong \mathrm{Dist}(G(F), \Lambda)^{G(F)},$$

and similarly for $G_b(F)$. With respect those isomorphisms, the left side of the desired equality is the characteristic class

$$\mathrm{cc}_{\mathrm{Bun}_{G, C}^1/S}(i_1^* T_{V^\vee}(i_b)_* \rho)$$

restricted to $\mathrm{Dist}(G(F)_{\mathrm{ell}}, \Lambda)^{G(F)}$.

For the remainder of the proof we introduce the abbreviations $B = \mathrm{Bun}_{G, C}$ and $H = \mathrm{Hecke}_{G, \leq \mu, C}$ and $\mathrm{In} = \mathrm{In}_S$. Let us also use a subscript “ell” to mean restriction to the appropriate elliptic locus. Taking inertia stacks in (6.1.2), we obtain a commutative diagram

$$\begin{array}{ccccc} & & \mathrm{In}(H^{*,1})_{\mathrm{ell}} & \xrightarrow{\mathrm{In}(h_2^{*,1})_{\mathrm{ell}}} & \mathrm{In}(B)_{\mathrm{ell}} & \\ & & \downarrow j_1' & & \downarrow j_1 & \\ & \mathrm{id} \swarrow & \mathrm{In}(H^{*,1}) & \xrightarrow{\mathrm{In}(h_2^{*,1})} & \mathrm{In}(B^1) & \\ & & \downarrow \mathrm{In}(i_1') & & \downarrow \mathrm{In}(i_1) & \\ \mathrm{In}(H^{b,*})_{\mathrm{ell}} & \xrightarrow{j_b'} & \mathrm{In}(H^{b,*}) & \xrightarrow{\mathrm{In}(i_b')} & \mathrm{In}(H) & \xrightarrow{\mathrm{In}(h_2)} & \mathrm{In}(B) \\ \downarrow \mathrm{In}(h_1^{b,*})_{\mathrm{ell}} & & \downarrow \mathrm{In}(h_1^{b,*}) & & \downarrow \mathrm{In}(h_1) & & \downarrow \mathrm{In}(i_b) \\ \mathrm{In}(B^b)_{\mathrm{ell}} & \xrightarrow{j_b} & \mathrm{In}(B^b) & \xrightarrow{\mathrm{In}(i_b)} & \mathrm{In}(B) & & \end{array} \quad (6.4.2)$$

in which all squares are cartesian, and the morphism labeled id is the equality from Corollary 6.2.4. The characteristic class in question is

$$\begin{aligned} j_1^* \mathrm{cc}_{B^1/S}(i_1^* T_{V^\vee}(i_b)_* \rho) &\stackrel{\text{Lem. 4.3.6}}{=} j_1^* \mathrm{In}(i_1)^* \mathrm{cc}_{B/S}((h_2)_!(h_1^*(i_b)_* \rho \otimes \mathcal{S}_{V^\vee})) \\ &\stackrel{\text{Cor. 4.3.8}}{=} j_1^* \mathrm{In}(i_1)^* \mathrm{In}(h_2)_* \mathrm{cc}_{H/S}(h_1^*(i_b)_* \rho \otimes \mathcal{S}_{V^\vee}) \\ &= (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_*(j_1)_{\mathrm{ell}}^* \mathrm{In}(i_1')^* \mathrm{cc}_{H/S}(h_1^*(i_b)_* \rho \otimes \mathcal{S}_{V^\vee}) \\ &= (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_*(j_b)_{\mathrm{ell}}^* \mathrm{In}(i_b')^* \mathrm{cc}_{H/S}(h_1^*(i_b)_* \rho \otimes \mathcal{S}_{V^\vee}) \\ &\stackrel{\text{Lem. 4.3.6}}{=} (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_*(j_b)_{\mathrm{ell}}^* \mathrm{cc}_{H^{b,*}/S}((i_b')^* h_1^*(i_b)_* \rho \otimes \mathcal{S}_{V^\vee}) \\ &= (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_*(j_b)_{\mathrm{ell}}^* \mathrm{cc}_{H^{b,*}/S}((h_1^{b,*})^* \rho \otimes \mathcal{S}_{V^\vee}). \end{aligned}$$

Noting that $H^{b,*} \cong [\mathrm{Gr}_{G,\leq-\mu,C}^b / G_b(F)_S]$, we have a cartesian diagram of S -Artin v-stacks:

$$\begin{array}{ccc} H^{b,*} & \longrightarrow & H_m^{\mathrm{loc}} \\ \downarrow & & \downarrow \\ B^b & \longrightarrow & B_m^{\mathrm{loc}} \end{array} \quad (6.4.3)$$

Here $B^b \cong [S/G_b(F)_S]$, $H_m^{\mathrm{loc}} \cong [\mathrm{Gr}_{G,\leq-\mu,C} / L_m^+ G]$, and $B_m^{\mathrm{loc}} \cong [S/L_m^+ G]$; the m here is chosen large enough so that the action of $L^+ G$ on $\mathrm{Gr}_{G,\leq-\mu,C}$ factors through the quotient $L_m^+ G$. Through this, we can identify $(h_1^{b,*})^* \rho \otimes \mathcal{S}_{V^\vee}$ with $\rho \boxtimes_{B_m^{\mathrm{loc}}} \mathcal{S}_{V^\vee}$. It is at this point we apply Theorem 4.5.3, valid because the base $B_m^{\mathrm{loc}} = [S/L_m^+ G]$ satisfies the hypotheses of Lemma 4.5.2. We get

$$\begin{aligned} j_1^* \mathrm{cc}_{B^b/S} (i_1^* T_{V^\vee} (i_b)_* \rho) &= (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_* (j_b)_{\mathrm{ell}}^* \left(\mathrm{cc}_{B^b/S} \rho \boxtimes_{\mathrm{In}(B_m^{\mathrm{loc}})} \mathrm{cc}_{H_m^{\mathrm{loc}}/S} (\mathcal{S}_{V^\vee}) \right). \\ &= (\mathrm{In}(h_2^{*,1})_{\mathrm{ell}})_* \left(\mathrm{cc}_{B^b/S} \rho_{\mathrm{ell}} \boxtimes_{\mathrm{In}(B_m^{\mathrm{loc}})_{\mathrm{sr}}} \mathrm{cc}_{H_m^{\mathrm{loc}}/S} (\mathcal{S}_{V^\vee})_{\mathrm{sr}} \right) \end{aligned}$$

Considering the diagram

$$\begin{array}{ccc} H^0(\mathrm{In}(B^b)_{\mathrm{ell}}, K_{\mathrm{In}(B^b)/S}) & \xrightarrow{\sim} & \mathrm{Dist}(G_b(F)_{\mathrm{ell}}, \Lambda)^{G_b(F)} \\ \downarrow -\boxtimes_{\mathrm{In}(B_m^{\mathrm{loc}})_{\mathrm{sr}}} \mathrm{cc}_{H_m^{\mathrm{loc}}/S} (\mathcal{S}_{V^\vee})_{\mathrm{sr}} & & \downarrow q_1^* (-) \otimes (-)^{\langle 2\rho_{G,\mu} \rangle} \mathrm{rank} V^\vee [-] \\ H^0(\mathrm{In}(H^{b,*})_{\mathrm{ell}}, K_{H^{b,*}/S}) & \xrightarrow{\sim} & \mathrm{Dist}(\mathrm{Fix}(\alpha_b)_{\mathrm{ell}}, \Lambda)^{G_b(F)} \\ = \downarrow & & \downarrow (p_1)_*^{-1} \\ H^0(\mathrm{In}(H^{b,1})_{\mathrm{ell}}, K_{\mathrm{In}(H^{b,1})/S}) & \xrightarrow{\sim} & \mathrm{Dist}(\mathrm{Fix}(\alpha_{\mathrm{Sht}})_{\mathrm{ell}}, \Lambda)^{G(F) \times G_b(F)} \\ = \downarrow & & \downarrow (p_2)_* \\ H^0(\mathrm{In}(H^{*,1})_{\mathrm{ell}}, K_{\mathrm{In}(H^{*,1})/S}) & \xrightarrow{\sim} & \mathrm{Dist}(\mathrm{Fix}(\alpha_1)_{\mathrm{ell}}, \Lambda)^{G(F)} \\ \downarrow \mathrm{In}(h_2^{*,1})_* & & \downarrow (q_2)_* \\ H^0(\mathrm{In}(B^1)_{\mathrm{ell}}, K_{\mathrm{In}(B^1)/S}) & \xrightarrow{\sim} & \mathrm{Dist}(G(F)_{\mathrm{ell}}, \Lambda)^{G(F)} \end{array}$$

our characteristic class is the image of $\mathrm{cc}_{B^b/S}(\rho)_{\mathrm{ell}}$ under the composite vertical map on the left. The diagram is commutative; the hardest thing to check is the commutativity of the top square, which follows from Proposition 6.4.8 below. The composition along the right column is $\mathcal{T}_{b,\mu}^{G_b \rightarrow G}$, giving us the desired equality of distributions. \square

It remains to justify one step in this computation. Maintain the notation and assumptions of the previous theorem. The v-stack $H^{b,*}$ can be expressed as a fiber product as in (6.4.3); we have the ULA object $\rho \boxtimes_{B_m^{\mathrm{loc}}} \mathcal{S}_{V^\vee}$, whose characteristic class can be calculated using Theorem 4.5.3.

Let

$$\alpha_b: G_b(F)_S \times \mathrm{Gr}_{G,\leq-\mu} \rightarrow \mathrm{Gr}_{G,\leq-\mu}$$

be the action map, so that we have an isomorphism

$$\mathrm{In}_S(H^{b,*}) \cong [\mathrm{Fix}(\alpha_b)/G_b(F)_S].$$

Let $\mathrm{Fix}(\alpha_b)_{\mathrm{sr}}$ be the open subset lying over $G_b(F)_{\mathrm{sr}}$ (and use the same convention for other objects); then $\mathrm{Fix}(\alpha_b)_{\mathrm{sr}}$ is a locally profinite set, which is finite over $G_b(F)_{\mathrm{sr}}$ with fibers $X_*(T)_{\leq -\mu}$.

Proposition 6.4.8. *The characteristic class*

$$\mathrm{cc}_{H/S}(\rho \boxtimes \mathcal{S}_{V^\vee})_{\mathrm{sr}} \in H^0(\mathrm{In}_S(H)_{\mathrm{sr}}, K_{\mathrm{In}_S(T)/S}) \cong \mathrm{Dist}(\mathrm{Fix}(\alpha_b)_{\mathrm{sr}}, \Lambda)^{G_b(F)}$$

equals the image of $\mathrm{tr. dist}(\rho) \otimes (-1)^{\langle 2\rho_G, - \rangle} \mathrm{rank} V^\vee[-]$ under the evident map

$$\mathrm{Dist}(G_b(F)_{\mathrm{sr}}, \Lambda)^{G_b(F)} \otimes C(X_*(T)_{\leq -\mu}, \Lambda)^W \rightarrow \mathrm{Dist}(\mathrm{Fix}(\alpha_b)_{\mathrm{sr}}, \Lambda)^{G_b(F)}. \quad (6.4.4)$$

Proof. Take inertia stacks in (6.4.3), and restrict to the strongly regular locus in $\mathrm{In}_S[S/L_m^+G]$ to obtain a cartesian diagram

$$\begin{array}{ccc} [\mathrm{Fix}(\alpha_b)_{\mathrm{sr}}/G_b(F)_S] & \longrightarrow & [G_{b,\mathrm{sr}}(F)_S // G_b(F)_S] \\ \downarrow & & \downarrow \\ X_*(T)_{\leq \mu} \times^W [L_m^+T_{\mathrm{sr}} // T_{\mathrm{sr}}] & \longrightarrow & [L_m^+T_{\mathrm{sr}}/(W \rtimes L_m^+T)] \end{array}$$

The Künneth map (4.5.2) in this situation reduces to (6.4.4) on the level of global sections. The result now follows from Theorems 4.5.3 and 5.1.4. \square

This formally implies the following theorem.

Theorem 6.4.9. *Let ρ be any finite length admissible $G_b(F)$ -representation with $\overline{\mathbf{Q}}_\ell$ -coefficients. Assume that ρ admits a $\overline{\mathbf{Z}}_\ell$ -lattice in the sense of Definition C.2.1. Then for all $\phi \in C_c(G(F)_{\mathrm{ell}}, \overline{\mathbf{Q}}_\ell)$, the equality*

$$\mathrm{tr}(\phi | \mathrm{Mant}_{b,\mu}(\rho)) = \left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho) \right] (\phi)$$

holds.

Recall that by definition, $\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho)(\phi)$ depends only on $(\mathrm{tr. dist} \rho)_{\mathrm{ell}}$.

Proof. Fix ρ and ϕ as in the theorem, and fix a $\overline{\mathbf{Z}}_\ell$ -lattice $\rho^\circ \subset \rho$. After rescaling, we may also assume ϕ is valued in $\overline{\mathbf{Z}}_\ell$. It is clear from the definitions that $\mathrm{tr}(\phi | \mathrm{Mant}_{b,\mu}(\rho)) = \mathrm{tr}(\phi | R\Gamma(G, b, \mu)[\rho^\circ])$ and

$$\left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho) \right] (\phi) = \left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho^\circ) \right] (\phi).$$

For all $n \geq 1$, set $\rho_n^\circ = \rho^\circ \otimes \mathbf{Z}/\ell^n$, and write $\phi_n \in C_c(G(F)_{\mathrm{ell}}, \overline{\mathbf{Z}}_\ell/\ell^n)$ for the obvious reductions of ϕ . Applying Proposition 6.4.7 with $\Lambda = \overline{\mathbf{Z}}_\ell/\ell^n$, we get equalities

$$\mathrm{tr}(\phi_n | R\Gamma(G, b, \mu)[\rho_n^\circ]) = \left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho_n^\circ) \right] (\phi_n)$$

for all $n \geq 1$. The result now follows by taking the inverse limit over n . \square

6.5 Proof of Theorem 1.0.2

We are finally ready to prove the main theorem of the paper, which we restate for the convenience of the reader.

Theorem 6.5.1. *Assume the refined local Langlands correspondence [Kal16a, Conjecture G]. Let $\phi: W_F \times \mathrm{SL}_2 \rightarrow {}^L G$ be a discrete Langlands parameter with coefficients in $\overline{\mathbf{Q}}_\ell$, and let $\rho \in \Pi_\phi(G_b)$ be a member of its L -packet. After ignoring the action of W_E , we have an equality*

$$\mathrm{Mant}_{b,\mu}(\rho) = \sum_{\pi \in \Pi_\phi(G)} [\dim \mathrm{Hom}_{S_\phi}(\delta_{\pi,\rho}, r_\mu)] \pi + \mathrm{err}$$

in $\mathrm{Groth}(G(F))$, where $\mathrm{err} \in \mathrm{Groth}(G(F))$ is a virtual representation whose character vanishes on the locus of elliptic elements of $G(F)$.

If the packet $\Pi_\phi(G)$ consists entirely of supercuspidal representations and the semisimple L -parameter φ_ρ associated with ρ as in [FS21, §I.9.6] is supercuspidal, then in fact $\mathrm{err} = 0$.

The main ingredient in the proof is the following extension of Theorem 6.4.9 to its natural level of generality.

Theorem 6.5.2. *Let ρ be any finite length admissible $G_b(F)$ -representation with $\overline{\mathbf{Q}}_\ell$ -coefficients. Then for all $\phi \in C_c(G(F)_{\mathrm{ell}}, \overline{\mathbf{Q}}_\ell)$, the equality*

$$\mathrm{tr}(\phi | \mathrm{Mant}_{b,\mu}(\rho)) = \left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\mathrm{tr. dist} \rho) \right] (\phi)$$

holds.

In particular, the virtual character of $\mathrm{Mant}_{b,\mu}(\rho)$ restricted to $G(F)_{\mathrm{ell}}$ is equal to $T_{b,\mu}^{G_b \rightarrow G}(\Theta_\rho)$.

We will formally deduce this from Theorem 6.4.9 by a continuity argument.

For the proof of this theorem, it will be convenient to use the language of Grothendieck groups. In particular, by the finiteness results mentioned above, $\mathrm{Mant}_{b,\mu}(-)$ can be regarded as a group homomorphism $\mathrm{Mant}_{b,\mu}(-) : \mathrm{Groth}(G_b(F)) \rightarrow \mathrm{Groth}(G(F))$. Recall that (after choosing a Haar measure) any element $\phi \in C_c(G(F), \overline{\mathbf{Q}}_\ell)$ defines a linear form $\mathrm{tr}(\phi | -) : \mathrm{Groth}(G(F)) \rightarrow \overline{\mathbf{Q}}_\ell$. By definition, a linear form $f : \mathrm{Groth}(G(F)) \rightarrow \overline{\mathbf{Q}}_\ell$ is a *trace form* if it can be written as $\mathrm{tr}(\phi | -)$ for some $\phi \in C_c(G(F), \overline{\mathbf{Q}}_\ell)$. The key ingredient in the proof of Theorem 6.5.2 is the following result, which roughly says that $\mathrm{Mant}_{b,\mu}(\rho)$ is a continuous function of ρ .

Theorem 6.5.3. *For any $\phi \in C_c(G(F), \overline{\mathbf{Q}}_\ell)$, the linear form*

$$\mathrm{tr}(\phi | \mathrm{Mant}_{b,\mu}(-)) : \mathrm{Groth}(G_b(F)) \rightarrow \overline{\mathbf{Q}}_\ell$$

is a trace form.

With future applications in mind, we'll actually prove the following refined form of this theorem which also accounts for the Weil group action.

Theorem 6.5.4. *For any fixed $\phi \in C_c(G(F), \overline{\mathbf{Q}_\ell})$ and $w \in W_E$, the linear form $\text{Groth}(G_b(F)) \rightarrow \overline{\mathbf{Q}_\ell}$ defined by*

$$\rho \mapsto \text{tr}(\phi \times w | R\Gamma(G, b, \mu)[\rho])$$

is a trace form.

In the classical setting of Rapoport-Zink spaces, this was conjectured by Taylor, cf. [Shi12, Conjecture 8.3]. Taking $w = 1$, we deduce Theorem 6.5.3.

Proof. For any reductive group H/F , the trace Paley-Wiener theorem of Bernstein-Deligne-Kazhdan, [BDK86], characterizes trace forms among all linear forms on $\text{Groth}(H(F))$ by the following two conditions:

1. There is some open compact subgroup $K \subset H(F)$ such that $f(\pi) \neq 0$ only if $\pi^K \neq 0$.
2. For any parabolic $P = MU \subset H$ and any irreducible smooth $M(F)$ -representation σ , $f(i_M^H(\sigma\psi))$ is an algebraic function of ψ , where ψ varies over the unramified characters of $M(F)$. Here $i_M^H(-)$ denotes normalized parabolic induction.

We'll prove the theorem by showing that the linear form $\text{tr}(\phi \times w | R\Gamma(G, b, \mu)[-])$ satisfies the conditions of the trace Paley-Wiener theorem, applied to the group $H = G_b$.

Verification of Condition 1. Fix a pro- p open compact subgroup $K \subset G(F)$ such that ϕ is bi- K -invariant. If $\text{tr}(\phi \times w | R\Gamma(G, b, \mu)[\rho]) \neq 0$, then $(i_{\mathbf{1}}^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{b*} \rho)^K \neq 0$. Therefore, it suffices to see that there is some open compact $K' \subset G_b(F)$ such that $(i_{\mathbf{1}}^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{b*} \rho)^K \neq 0$ only if $\rho^{K'} \neq 0$. For this, write

$$\begin{aligned} (i_{\mathbf{1}}^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{b*} \rho)^K &\cong R\text{Hom}(i_{\mathbf{1}!} \text{cInd}_K^{G(F)} \overline{\mathbf{Q}_\ell}, T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{b*} \rho) \\ &\cong R\text{Hom}(T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{\mathbf{1}!} \text{cInd}_K^{G(F)} \overline{\mathbf{Q}_\ell}, i_{b*} \rho) \\ &\cong R\text{Hom}(i_b^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{\mathbf{1}!} \text{cInd}_K^{G(F)} \overline{\mathbf{Q}_\ell}, \rho). \end{aligned}$$

But now $i_b^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{\mathbf{1}!} \text{cInd}_K^{G(F)} \overline{\mathbf{Q}_\ell}$ is compact, and hence supported on only finitely many Bernstein components for $G_b(F)$. This shows that the irreducible ρ 's with $(i_{\mathbf{1}}^* T_{V_{\mu, \overline{\mathbf{Q}_\ell}}^\vee} i_{b*} \rho)^K \neq 0$ are supported on finitely many Bernstein components for $G_b(F)$. Quite generally, if Θ is any finite union of Bernstein components for $G_b(F)$, we can choose an open compact subgroup $K' \subset G_b(F)$ such that $\rho^{K'} \neq 0$ if ρ is supported on Θ . This gives the result.

Verification of Condition 2. Fix $P = MU \subset G_b$ and σ as in Condition 2. Let $X = \text{Spec} R$ be the smooth affine algebraic variety over $\overline{\mathbf{Q}_\ell}$ parametrizing unramified characters of $M(F)$. Let $\psi : M(F) \rightarrow R^\times$ be the universal character, and form $\Pi = i_M^{G_b}(\sigma\psi)$. This is an admissible smooth $R[G_b(F)]$ -module interpolating the parabolic inductions $i_M^{G_b}(\sigma\psi)$ over varying unramified characters ψ in the evident sense.⁴ Since Π is admissible, the pushforward $i_{b*} \Pi \in D_{\text{lis}}(\text{Bun}_G, R)$ is ULA. Since Hecke operators preserve ULA complexes, we deduce that $i_{\mathbf{1}}^* T_{V_{\mu, R}^\vee} i_{b*} \Pi \in D(G(F), R)^{BW_E}$ is an

⁴To see that Π is admissible in our slightly nonstandard sense, observe first that Π^K is finitely generated as an R -module for all pro- p open compact subgroups $K \subset G_b(F)$, since $P(F) \backslash G_b(F) / K$ is finite. But R is a smooth $\overline{\mathbf{Q}_\ell}$ -algebra, so any finitely generated R -module is automatically a perfect complex, giving the desired result.

admissible complex of smooth $R[G(F)]$ -modules with W_E -action, which interpolates the individual complexes

$$R\Gamma(G, b, \mu)[i_M^{G_b}(\sigma\psi)] = i_{\mathbf{1}}^* T_{V_{\mu, \mathbf{Q}_\ell}^\vee} i_{b*} i_M^{G_b}(\sigma\psi)$$

in the evident sense.

Now fix a pro- p open compact subgroup $K \subset G(F)$ such that ϕ is bi- K -invariant, so $\phi \times w$ defines an endomorphism of the perfect complex

$$(i_{\mathbf{1}}^* T_{V_{\mu, R}^\vee} i_{b*} \Pi)^K \in \text{Perf}(R).$$

Let $f \in R$ be the trace of this endomorphism. Unwinding definitions, we see that for any unramified character $\psi : M(F) \rightarrow \overline{\mathbf{Q}_\ell}^\times$ with associated point $x_\psi \in X(\overline{\mathbf{Q}_\ell})$, there is an equality

$$f(x_\psi) = \text{tr} \left(\phi \times w | R\Gamma(G, b, \mu)[i_M^{G_b}(\sigma\psi)] \right).$$

This shows that $\text{tr}(\phi \times w | R\Gamma(G, b, \mu)[i_M^{G_b}(\sigma\psi)])$ is an algebraic function of ψ , as desired. \square

Let us say a subset $S \subset \text{Irr}_{\overline{\mathbf{Q}_\ell}}(G(F))$ is dense if any trace form on $\text{Groth}(G(F))$ which vanishes on S vanishes identically. For instance, the Langlands classification implies that (for any fixed choice of isomorphism $\mathbf{C} \simeq \overline{\mathbf{Q}_\ell}$) tempered representations are dense, cf. [Kaz86, Theorem 0].

Lemma 6.5.5. *The subset of irreducible representations $\pi \in \text{Irr}_{\overline{\mathbf{Q}_\ell}}(G(F))$ admitting $\overline{\mathbf{Z}_\ell}$ -lattices is dense.*

It seems reasonable to think of this lemma as an ℓ -adic analogue of the density of tempered representations.

Proof. Let f be a trace form, and assume that $f(\tau) = 0$ for every $\tau \in \text{Irr}_{\overline{\mathbf{Q}_\ell}}(G(F))$ admitting a $\overline{\mathbf{Z}_\ell}$ -lattice.

By Proposition C.2.2, it's enough to show that $f(i_M^G(\sigma\psi)) = 0$ for any parabolic $P = MU \subset G$, any unramified character ψ of $M(F)$, and any $\sigma \in \text{Irr}_{\overline{\mathbf{Q}_\ell}}(M(F))$ admitting a $\overline{\mathbf{Z}_\ell}$ -lattice. Fix P and σ , and consider the function g on unramified characters of $M(F)$ sending ψ to $f(i_M^G(\sigma\psi))$. By the easy direction of the trace Paley-Wiener theorem, g is a regular function on the variety of unramified characters of $M(F)$.

Let us say an unramified character ψ is integral if it takes values in $\overline{\mathbf{Z}_\ell}^\times$. If ψ is integral, then $\sigma\psi$ admits a $\overline{\mathbf{Z}_\ell}$ -lattice, and hence also $i_M^G(\sigma\psi)$ admits a $\overline{\mathbf{Z}_\ell}$ -lattice. In particular, if ψ is integral and $i_M^G(\sigma\psi)$ is irreducible, then $g(\psi) = 0$ by combining these observations with our assumption on f . Now integral characters are Zariski-dense in the variety of unramified characters of $M(F)$, and the subset T of integral characters such that $i_M^G(\sigma\psi)$ is irreducible is also Zariski-dense (use [Dat05, Theorem 5.1]). Since $g(\psi) = 0$ for all $\psi \in T$, we deduce that $g \equiv 0$, so in particular

$$0 = g(\psi) = f(i_M^G(\sigma\psi))$$

for all ψ . This gives the result. \square

Proof of Theorem 6.5.2. Fix ϕ as in the statement of the theorem, and consider the linear form

$$f(-) = \text{tr}(\phi | \text{Mant}_{b,\mu}(-)) - \left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\text{tr. dist } -) \right] (\phi)$$

on $\text{Groth}(G_b(F))$. By Theorem 6.4.9, we know that $f(\rho) = 0$ if ρ admits a lattice. We need to show that f vanishes identically.

The key observation is that f is a trace form. Indeed, $\text{tr}(\phi | \text{Mant}_{b,\mu}(-))$ is a trace form by Theorem 6.5.3. Moreover, $\left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\text{tr. dist } -) \right] (\phi)$ is a trace form, since we can rewrite $\left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\text{tr. dist } \rho) \right] (\phi)$ as the trace of $\tilde{T}_{b,\mu}^{G \rightarrow G_b}(\phi) \in C_c(G_b(F)_{\text{ell}}, \overline{\mathbf{Q}}_\ell)$ acting on ρ . Thus f is a difference of trace forms, and hence a trace form. Since $f(\rho) = 0$ for any ρ admitting a lattice, Lemma 6.5.5 now implies the desired result.

For the final claim about virtual characters, choose compatible $\overline{\mathbf{Q}}_\ell$ -valued Haar measures dg and dg' on $G(F)$ and $G_b(F)$. Fix some ρ , and let $\Xi \in C(G(F)_{\text{sr}} // G(F), \overline{\mathbf{Q}}_\ell)$ be the virtual character of $\text{Mant}_{b,\mu}(\rho)$. Pick any $\phi \in C_c(G(F)_{\text{ell}}, \overline{\mathbf{Q}}_\ell)$. Then

$$\text{tr}(\phi | \text{Mant}_{b,\mu}(\rho)) = \int_{G(F)} \Xi(g) \phi(g) dg$$

by definition. On the other hand,

$$\left[\mathcal{T}_{b,\mu}^{G_b \rightarrow G}(\text{tr. dist } \rho) \right] (\phi) = \int_{G(F)} T_{b,\mu}^{G_b \rightarrow G}(\Theta_\rho)(g) \phi(g) dg$$

by compatibility of the Haar measures and Proposition 6.3.5. Combining these observations, we get an equality

$$\int_{G(F)} T_{b,\mu}^{G_b \rightarrow G}(\Theta_\rho)(g) \phi(g) dg = \int_{G(F)} \Xi(g) \phi(g) dg$$

for any $\phi \in C_c(G(F)_{\text{ell}}, \overline{\mathbf{Q}}_\ell)$. The result now follows by varying ϕ . \square

Proof of Theorem 6.5.1. The claimed equality in $\text{Groth}(G(F))$ is an immediate consequence of Theorem 6.5.2 and Theorem 3.2.9.

For the claim regarding the error term, consider the virtual representation

$$\text{err} = \text{Mant}_{b,\mu}(\rho) - \sum_{\pi \in \Pi_\phi(G)} \dim \text{Hom}_{S_\phi}(\delta_{\pi,\rho}, r_\mu) \pi.$$

By the first half of the theorem, we know that err is non-elliptic. By Theorem C.1.1, it thus suffices to show that err is a virtual sum of supercuspidal representations. Since the packet $\Pi_\phi(G)$ is supercuspidal by assumption, we're reduced to showing that $\text{Mant}_{b,\mu}(\rho)$ is a virtual supercuspidal representation. By definition, this is the Grothendieck class of the complex $A = i_{\mathbf{1}}^* T_{V_{\mu, \overline{\mathbf{Q}}_\ell}} i_{b*} \rho \in D(G(F), \overline{\mathbf{Q}}_\ell)$, so we need to see that any irreducible τ occurring in the Jordan-Holder series of $H^*(A)$ is supercuspidal. Since $\varphi_\tau = \varphi_\rho$ by the commutation of Hecke operators with excursion operators, the claim now follows from the assumption on φ_ρ and [FS21, Theorem I.9.6.viii]. \square

6.6 Application to inner forms of GL_n

We give an application to the local Langlands correspondence. Recall that for any G/F , any $b \in B(G)$ and any $\tau \in \mathrm{Irr}(G_b(F))$, the construction in [FS21, Proposition I.9.1] (applied to $A = i_b! \tau$) gives rise to a semisimple L -parameter $\varphi_\tau : W_F \rightarrow {}^L G(\overline{\mathbf{Q}}_\ell)$ associated with τ . This construction is canonical and satisfies a long list of desirable properties [FS21, Theorem I.9.6]. However, it is a highly nontrivial problem to compare this construction with “previously known” realizations of the local Langlands correspondence.

Theorem 6.6.1. *Let G be any inner form of GL_n/F , and let π be an irreducible smooth representation of $G(F)$. Then the L -parameter φ_π associated with π as in [FS21, §I.9] agrees with the usual semisimplified L -parameter attached to π .*

Proof. By [FS21, Theorem I.9.6.viii], we can assume π is supercuspidal. Pick some basic b with $G_b = \mathrm{GL}_n/F$, and let $\rho \in \mathrm{Irr}(G_b(F))$ be the Jacquet-Langlands transfer of π [DKV84], so the (usual) semisimple L -parameters of ρ and π agree. By [FS21, Theorem 1.9.6.viii-ix], we know that φ_ρ agrees with the usual semisimple L -parameter of ρ . To prove the theorem, it thus suffices to show that $\varphi_\pi = \varphi_\rho$.

Pick any μ such that $b \in B(G, \mu)$. By Theorem 6.5.2 and the usual character relation characterizing the Jacquet-Langlands correspondence, we have an equality $\mathrm{Mant}_{b, \mu}(\rho) = \dim V_\mu \cdot \pi + e$ in $R(G)$, where e is a non-elliptic virtual representation. Since π is supercuspidal, this implies that π occurs as a subquotient of some cohomology group of the complex $A = i_{\mathbf{1}}^* T_{V, \overline{\mathbf{Q}}_\ell}^{\vee} i_{b*} \rho \in D(G(F), \overline{\mathbf{Q}}_\ell)$. But Hecke operators commute with excursion operators, so $\varphi_\tau = \varphi_\rho$ for any irreducible τ occurring in the Jordan-Holder series of $H^*(A)$. \square

A Endoscopy

A.1 Endoscopic character relations

We recall here the endoscopic character identities, which are part of the refined local Langlands correspondence, following the formulation of [Kal16b, §5.4], also recalled in [Kal16a, §4.2]. They will be an important ingredient in the proof of our main result.

We will use the following notation.

- F/\mathbf{Q}_p is a finite extension, F^{nr}/F a maximal unramified extension.
- G is a connected reductive group defined over F .
- G^* is a quasi-split connected reductive group defined over F .
- Ψ is a G^* -conjugacy class of inner twists $\psi: G^* \rightarrow G$.
- $\bar{z}_\sigma = \psi^{-1}\sigma(\psi) \in G_{\text{ad}}^*$, so that $\bar{z} \in Z^1(F, G_{\text{ad}}^*)$.
- $z \in Z^1(u \rightarrow W, Z(G^*) \rightarrow G^*)$ is a lift of \bar{z} .
- $b \in G(F^{\text{nr}})$ is a decent basic element.
- G_b is the corresponding inner form of G .
- $\xi: G_{F^{\text{nr}}} \rightarrow G_{b, F^{\text{nr}}}$ is the identity map.
- $z_b \in Z^1(u \rightarrow W, Z(G) \rightarrow G)$ is the image of b under (2.3.1).
- \mathfrak{w} is a Whittaker datum for G^* .
- $\phi: W_F \rightarrow {}^L G$ is a discrete parameter.
- $S_\phi = \text{Cent}(\phi, \widehat{G})$.
- S_ϕ^+ is the group defined in Definition 2.3.1.
- λ_z resp. λ_{z_b} the image of the class of z resp. z_b under the isomorphism $H^1(u \rightarrow W, Z(G^*) \rightarrow G^*) \rightarrow \pi_0(Z(\widehat{G})^+)^*$.

Associated to ϕ are the L -packets $\Pi_\phi(G)$ and $\Pi_\phi(G_b)$ and the bijections

$$\Pi_\phi(G) \rightarrow \text{Irr}(\pi_0(S_\phi^+), \lambda_z), \quad \Pi_\phi(G_b) \rightarrow \text{Irr}(\pi_0(S_\phi^+), \lambda_z + \lambda_{z_b})$$

denoted by $\pi \mapsto \tau_{z, \mathfrak{w}, \pi}$ and $\rho \mapsto \tau_{z, \mathfrak{w}, \rho}$.

We now choose a semi-simple element $s \in S_\phi$ and an element $\dot{s} \in S_\phi^+$ which lifts s . Let $e(G)$ and $e(G_b)$ be the Kottwitz signs of the groups G and G_b , as defined in [Kot83]. Consider the virtual characters

$$e(G) \sum_{\pi \in \Pi_\phi(G)} \text{tr } \tau_{z, \mathfrak{w}, \pi}(\dot{s}) \cdot \Theta_\pi \quad \text{and} \quad e(G_b) \sum_{\rho \in \Pi_\phi(G_b)} \text{tr } \tau_{z, \mathfrak{w}, \rho}(\dot{s}) \cdot \Theta_\rho.$$

The endoscopic character identities are equations which relate these two virtual characters to virtual characters on an endoscopic group H_1 . From the pair (ϕ, \dot{s}) one obtains a refined elliptic endoscopic datum

$$\dot{\mathbf{e}} = (H, \mathcal{H}, \dot{s}, \eta) \tag{A.1.1}$$

in the sense of [Kal16b, §5.3] as follows. Let $\widehat{H} = \text{Cent}(s, \widehat{G})^\circ$. The image of ϕ is contained in $\text{Cent}(s, \widehat{G})$, which in turns acts by conjugation on its connected component \widehat{H} . This gives a homomorphism $W_F \rightarrow \text{Aut}(\widehat{H})$. Letting $\Psi_0(\widehat{H})$ be the based root datum of \widehat{H} [Kot84b, §1.1] and $\Psi_0^\vee(\widehat{H})$ its dual, we obtain the homomorphism

$$W_F \rightarrow \text{Aut}(\widehat{H}) \rightarrow \text{Out}(\widehat{H}) = \text{Aut}(\Psi_0(\widehat{H})) = \text{Aut}(\Psi_0^\vee(\widehat{H})).$$

Since the target is finite, this homomorphism extends to Γ_F and we obtain a based root datum with Galois action, hence a quasi-split connected reductive group H defined over F . Its dual group is by construction equal to \widehat{H} . We let $\mathcal{H} = \widehat{H} \cdot \phi(W_F)$, noting that the right factor normalizes the left so their product \mathcal{H} is a subgroup of ${}^L G$. Finally, we let $\eta : \mathcal{H} \rightarrow {}^L G$ be the natural inclusion. Note that by construction ϕ takes image in \mathcal{H} , i.e. it factors through η .

We can realize the L -group of H as ${}^L H = \widehat{H} \rtimes W_F$, but we caution the reader that W_F does not act on \widehat{H} via the map $W_F \rightarrow \text{Aut}(\widehat{H})$ given by ϕ as above. Rather, we have to modify this action to ensure that it preserves a pinning of \widehat{H} . More precisely, after fixing an arbitrary pinning of \widehat{H} we obtain a splitting $\text{Out}(\widehat{H}) \rightarrow \text{Aut}(\widehat{H})$ of the projection $\text{Aut}(\widehat{H}) \rightarrow \text{Out}(\widehat{H})$ and the action of W_F on \widehat{H} we use to form ${}^L H$ is given by composing the above map $W_F \rightarrow \text{Out}(\widehat{H})$ with this splitting.

Both ${}^L H$ and \mathcal{H} are thus extensions of W_F by \widehat{H} , but they need not be isomorphic. If they are, we fix arbitrarily an isomorphism $\eta_1 : \mathcal{H} \rightarrow {}^L H$ of extensions. Then $\phi^s = \eta_1 \circ \phi$ is a supercuspidal parameter for H .

In the general case we need to introduce a z -pair $\mathfrak{z} = (H_1, \eta_1)$ as in [KS99, §2]. It consists of a z -extension $H_1 \rightarrow H$ (recall this means that H_1 has a simply connected derived subgroup and the kernel of $H_1 \rightarrow H$ is an induced torus) and $\eta_1 : \mathcal{H} \rightarrow {}^L H_1$ is an L -embedding that extends the natural embedding $\widehat{H} \rightarrow \widehat{H}_1$. As is shown in [KS99, §2.2], such a z -pair always exists. Again we set $\phi^s = \eta_1 \circ \phi$ and obtain a supercuspidal parameter for H_1 . In the situation where an isomorphism $\eta_1 : \mathcal{H} \rightarrow {}^L H$ does exist, we will allow ourselves to take $H = H_1$ and so regard $\mathfrak{z} = (H, \eta_1)$ as a z -pair, even though in general H will not have a simply connected derived subgroup.

The virtual character on H_1 that the above virtual characters on G and G_b are to be related to is

$$S\Theta_{\phi^s} := \sum_{\pi^s \in \Pi_{\phi^s}(H_1)} \dim(\tau_{\pi^s}) \Theta_{\pi^s}.$$

Here $\pi^s \mapsto \tau_{\pi^s}$ is a bijection $\Pi_{\phi^s}(H_1) \rightarrow \text{Irr}(\pi_0(\text{Cent}(\phi^s, \widehat{H}_1)/Z(\widehat{H}_1)^\Gamma))$ determined by an arbitrary choice of Whittaker datum for H_1 . The argument in the proof of Lemma 2.3.3 shows the independence of $\dim(\tau_{\pi^s})$ of the choice of a Whittaker datum for H_1 .

The relationship between the virtual characters on G , G_b , and H_1 , is expressed in terms of the Langlands-Shelstad transfer factor $\Delta'_{\text{abs}}[\dot{\mathbf{e}}, \mathfrak{z}, \mathbf{w}, (\psi, z)]$ for the pair of groups (H_1, G) and the corresponding Langlands-Shelstad transfer factor $\Delta'_{\text{abs}}[\dot{\mathbf{e}}, \mathfrak{z}, \mathbf{w}, (\xi \circ \psi, \psi^{-1}(z_b) \cdot z)]$ for the pair of groups (H_1, G_b) , both of which are defined by [Kal16b, (5.10)]. We will abbreviate both of them

to just Δ . It is a simple consequence of the Weyl integration formula that the character relation [Kal16b, (5.11)] can be restated in terms of character functions (rather than character distributions) as

$$e(G) \sum_{\pi \in \Pi_\phi(G)} \operatorname{tr} \tau_{z, \mathfrak{w}, \pi}(\dot{s}) \Theta_\pi(g) = \sum_{h_1 \in H_1(F)/\text{st.}} \Delta(h_1, g) S \Theta_{\phi^s}(h_1) \quad (\text{A.1.2})$$

for any strongly regular semi-simple element $g \in G(F)$. The sum on the right runs over stable conjugacy classes of strongly regular semi-simple elements of $H_1(F)$. We also have the analogous identity for G_b :

$$e(G_b) \sum_{\rho \in \Pi_\phi(G_b)} \operatorname{tr} \tau_{z, \mathfrak{w}, \rho}(\dot{s}) \Theta_\rho(g') = \sum_{h_1 \in H_1(F)/\text{st.}} \Delta(h_1, g') S \Theta_{\phi^s}(h_1). \quad (\text{A.1.3})$$

For the purposes of this paper, we are only interested in the right hand sides of these two equations as a bridge between their left-hand sides. Essential for this bridge is a certain compatibility between the transfer factors appearing on both right-hand sides:

Lemma A.1.1.

$$\Delta(h_1, g') = \Delta(h_1, g) \cdot \langle \operatorname{inv}[b](g, g'), s_{h,g}^\natural \rangle. \quad (\text{A.1.4})$$

We need to explain the second factor. Given maximal tori $T_H \subset H$ and $T \subset G$, there is a notion of an admissible isomorphism $T_H \rightarrow T$, for which we refer the reader to [Kal16a, §1.3]. Two strongly regular semi-simple elements $h \in H(\mathbf{Q}_p)$ and $g \in G(\mathbf{Q}_p)$ are called *related* if there exists an admissible isomorphism $T_h \rightarrow T_g$ between their centralizers mapping h to g . If such an isomorphism exists, it is unique, and in particular defined over F , and shall be called $\varphi_{h,g}$. An element $h_1 \in H_1(F)$ is called related to $g \in G(F)$ if and only if its image $h \in H(F)$ is so. Since g and g' are stably conjugate, an element $h_1 \in H_1(F)$ is related to g if and only if it is related to g' . If that is not the case, both $\Delta(h_1, g')$ and $\Delta(h_1, g)$ are zero and (A.1.4) is trivially true. Thus assume that h_1 is related to both g and g' . Let $s^\natural \in S_\phi$ be the image of \dot{s} under (2.3.2). Note that $s^\natural \in s \cdot Z(\widehat{G})^{\circ, \Gamma}$ and hence the preimage of s^\natural under η belongs to $Z(\widehat{H})^\Gamma$, which in turns embeds naturally into \widehat{T}_h^Γ . Using the admissible isomorphism $\varphi_{h,g}$ we transport s^\natural into \widehat{T}_g^Γ and denote it by $s_{h,g}^\natural$. It is then paired with $\operatorname{inv}[b](g, g')$ via the isomorphism $B(T_g) \cong X^*(\widehat{T}_g^\Gamma)$ of [Kot85, §2.4].

Proof. For every finite subgroup $Z \subset Z(G) \subset T_g$ one obtains from $\varphi_{h,g}$ an isomorphism $T_h/\varphi_{h,g}^{-1}(Z) \rightarrow T_g/Z$. Using the subgroups Z_n from the previous subsection we form the quotients $T_{h,n} = T_h/\varphi_{h,g}^{-1}(Z_n)$ and $T_{g,n} = T_g/Z_n$. From $\varphi_{h,g}$ we obtain an isomorphism

$$\widehat{T}_h \rightarrow \widehat{T}_g$$

between the limits over n of the tori dual to $T_{h,n}$ and $T_{g,n}$. Let $\dot{s}_{h,g} \in [\widehat{T}_g]^+$ be the image of \dot{s} under this isomorphism. Let $\operatorname{inv}[z_b](g, g') \in H^1(u \rightarrow W, Z(G) \rightarrow T_g)$ be the invariant defined in [Kal16b, §5.1]. If we replace $\langle \operatorname{inv}[b](g, g'), s_{h,g}^\natural \rangle$ by $\langle \operatorname{inv}[z_b](g, g'), \dot{s}_{h,g} \rangle$ then the lemma follows immediately from the defining formula [Kal16b, (5.10)] of the transfer factors. The lemma follows from the equality $\langle \operatorname{inv}[b](g, g'), s_{h,g}^\natural \rangle = \langle \operatorname{inv}[z_b](g, g'), \dot{s}_{h,g} \rangle$ proved in [Kal18, §4.2]. \square

A.2 The Kottwitz sign

We will give a formula for the Kottwitz sign $e(G)$ in terms of the dual group \widehat{G} . Fix a quasi-split inner form G^* and an inner twisting $\psi: G^* \rightarrow G$. Let $h \in H^1(\Gamma, G_{\text{ad}}^*)$ be the class of $\sigma \mapsto \psi^{-1}\sigma(\psi)$. Via the Kottwitz homomorphism [Kot86, Theorem 1.2] the class h corresponds to a character $\nu \in X^*(Z(\widehat{G}_{\text{sc}})^\Gamma)$.

Choose an arbitrary Borel pair $(\widehat{T}_{\text{sc}}, \widehat{B}_{\text{sc}})$ of \widehat{G}_{sc} and let $2\rho \in X_*(\widehat{T}_{\text{sc}})$ be the sum of the \widehat{B}_{sc} -positive coroots. The restriction map $X^*(\widehat{T}_{\text{sc}}) \rightarrow X^*(Z(\widehat{G}_{\text{sc}}))$ is surjective and we can lift ν to $\dot{\nu} \in X^*(\widehat{T}_{\text{sc}})$ and form $\langle 2\rho, \dot{\nu} \rangle \in \mathbf{Z}$. A different lift $\dot{\nu}$ would differ by an element of $X^*(\widehat{T}_{\text{ad}})$, and since $\rho \in X_*(\widehat{T}_{\text{ad}})$ we see that the image of $\langle 2\rho, \dot{\nu} \rangle$ in $\mathbf{Z}/2\mathbf{Z}$ is independent of the choice of lift $\dot{\nu}$. We thus write $\langle 2\rho, \nu \rangle \in \mathbf{Z}/2\mathbf{Z}$. Since any two Borel pairs in \widehat{G}_{sc} are conjugate $\langle 2\rho, \nu \rangle$ does not depend on the choice of $(\widehat{T}_{\text{sc}}, \widehat{B}_{\text{sc}})$.

Lemma A.2.1.

$$e(G) = (-1)^{\langle 2\rho, \nu \rangle}.$$

Proof. We fix Γ -invariant Borel pairs $(T_{\text{ad}}, B_{\text{ad}})$ in G_{ad}^* and $(\widehat{T}_{\text{sc}}, \widehat{B}_{\text{sc}})$ in \widehat{G}_{sc} . Then we have the identification $X^*(T_{\text{ad}}) = X_*(\widehat{T}_{\text{sc}})$. Let $(T_{\text{sc}}, B_{\text{sc}})$ be the preimage in G_{sc}^* of $(T_{\text{ad}}, B_{\text{ad}})$.

By definition the Kottwitz sign is the image of h under

$$H^1(\Gamma, G_{\text{ad}}^*) \xrightarrow{\delta} H^2(\Gamma, Z(G_{\text{sc}}^*)) \xrightarrow{\rho} H^2(\Gamma, \{\pm 1\}) \longrightarrow \{\pm 1\},$$

where $\rho \in X^*(T_{\text{sc}})$ is half the sum of the B_{sc} -positive roots and its restriction to $Z(G_{\text{sc}}^*)$ is independent of the choice of $(T_{\text{ad}}, B_{\text{ad}})$. By functoriality of the Tate-Nakayama pairing this is the same as pairing $\delta h \in H^2(\Gamma, Z(G_{\text{sc}}^*))$ with $\rho \in H^0(\Gamma, X^*(Z(G_{\text{sc}}^*)))$. The canonical pairing $X^*(T_{\text{ad}}) \otimes X^*(\widehat{T}_{\text{sc}}) \rightarrow \mathbf{Z}$ induces the perfect pairing $X^*(T_{\text{sc}})/X^*(T_{\text{ad}}) \otimes X^*(\widehat{T}_{\text{sc}})/X^*(\widehat{T}_{\text{ad}}) \rightarrow \mathbf{Q}/\mathbf{Z}$ and hence the isomorphism $X^*(Z(G_{\text{sc}}^*)) \rightarrow \text{Hom}_{\mathbf{Z}}(X^*(Z(\widehat{G}_{\text{sc}})), \mathbf{Q}/\mathbf{Z}) = Z(\widehat{G}_{\text{sc}})$, where the last equality uses the exponential map. Under this isomorphism $\rho \in X^*(Z(G_{\text{sc}}^*))^\Gamma$ maps to the element $(-1)^{2\rho} \in Z(\widehat{G}_{\text{sc}})^\Gamma$ obtained by mapping $(-1) \in \mathbf{C}^\times$ under $2\rho \in X^*(T_{\text{ad}}) = X_*(\widehat{T}_{\text{sc}})$. The lemma now follows from [Kot86, Lemma 1.8]. \square

B Elementary Lemmas

B.1 Homological algebra

Lemma B.1.1. *Let R be a discrete valuation ring with maximal ideal \mathfrak{m} . Let $\kappa = R/\mathfrak{m}$ be the residue field, and let $\Lambda = R/\mathfrak{m}^k$ for some $k > 0$. For a Λ -module M we have the dual module $M^* = \text{Hom}_\Lambda(M, \Lambda)$ and the natural morphisms $M \rightarrow M^{**}$ and $(M^* \otimes M) \rightarrow (M \otimes M^*)^*$.*

*The morphism $M \rightarrow M^{**}$ is an isomorphism if and only if M is finitely generated.*

Proof. For the ‘‘if’’ direction of the first point we note that the structure theorem for R -modules implies that a finitely generated Λ -module is a direct sum of finitely many cyclic Λ -modules, and each cyclic Λ -module is isomorphic to its own double dual.

Conversely assume that $M \rightarrow M^{**}$ is an isomorphism. We induct on k . If $k = 1$, then Λ is a field, and this is well-known. For general k we consider $N = M/\mathfrak{m}M$. The ring Λ is an Artinian

serial ring, and hence it is injective as a module over itself. Thus the dualization functor is exact, and we get a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathfrak{m}M & \longrightarrow & M & \longrightarrow & N \longrightarrow 0 \\
& & \downarrow & & \downarrow \cong & & \downarrow \\
0 & \longrightarrow & (\mathfrak{m}M)^{**} & \longrightarrow & M^{**} & \longrightarrow & N^{**} \longrightarrow 0,
\end{array} \tag{B.1.1}$$

which shows that the right-most vertical map is surjective and the left-most vertical map is injective.

We have an isomorphism of Λ -modules $\mathfrak{m}^{m-1}\Lambda \rightarrow \kappa$, from which we obtain

$$N^* = \mathrm{Hom}_\Lambda(N, \Lambda) = \mathrm{Hom}_\Lambda(N, \mathfrak{m}^{m-1}\Lambda) \cong \mathrm{Hom}_\kappa(N, \kappa).$$

Thus N^{**} is also the double dual of N in the category of κ -vector spaces, and it is easy to check that the right-most vertical map in (B.1.1) is the canonical map in that category. Thus, this map is an isomorphism, and N is finitely generated as a κ -vector space.

By the Snake Lemma, the left-most vertical arrow in (B.1.1) is an isomorphism. We can apply the inductive hypothesis to the $(\Lambda/\mathfrak{m}^{m-1})$ -module $\mathfrak{m}M$ and conclude that it is finitely generated. Thus so is M . □

Lemma B.1.2. *Let Λ be an arbitrary ring, and let $D(\Lambda)$ be the derived category of Λ -modules. For an object M of $D(\Lambda)$, let $\mathbf{D}M = \mathrm{RHom}(M, \Lambda[0])$.*

1. *Assume that Λ is self-injective. The natural morphism $M \rightarrow \mathbf{D}\mathbf{D}M$ is an isomorphism if and only if each $H^i(M)$ is finitely generated.*
2. *The following are equivalent:*
 - (a) *The natural maps $M \rightarrow \mathbf{D}\mathbf{D}M$ and $\mathbf{D}M \otimes M \rightarrow \mathbf{D}(M \otimes \mathbf{D}M)$ are isomorphisms.*
 - (b) *The natural map $M \otimes \mathbf{D}M \rightarrow \mathrm{RHom}(M, M)$ is an isomorphism.*
 - (c) *M is strongly dualizable; that is, for any object N , $N \otimes \mathbf{D}M \rightarrow \mathrm{RHom}(M, N)$ is an isomorphism.*
 - (d) *M is a compact object; that is, the functor $N \mapsto \mathrm{RHom}(M, N)$ commutes with colimits.*
 - (e) *M is a perfect complex; that is, M is isomorphic to a bounded complex of finitely generated projective Λ -modules.*

(Throughout, the \otimes means derived tensor product.)

Proof. For the first statement, the injectivity of Λ implies that $H^i(\mathbf{D}M) \cong H^{-i}(M)^*$, so that $H^i(\mathbf{D}\mathbf{D}M) \cong H^i(M)^{**}$. Therefore $M \rightarrow \mathbf{D}\mathbf{D}M$ is an isomorphism if and only if each $H^i(M) \rightarrow H^i(M)^{**}$ is an isomorphism. By Lemma B.1.1, this is equivalent to each $H^i(M)$ being finitely generated.

We now turn to the second statement. For (a) \implies (b), assume that $M \rightarrow \mathbf{D}\mathbf{D}M$ and $\mathbf{D}M \otimes M \rightarrow \mathbf{D}(M \otimes \mathbf{D}M)$ are isomorphisms. Then $\mathrm{RHom}(M, M) \cong \mathrm{RHom}(M, \mathbf{D}\mathbf{D}M) \cong \mathrm{RHom}(M \otimes \mathbf{D}M, \Lambda) \cong \mathbf{D}(M \otimes \mathbf{D}M) \cong \mathbf{D}M \otimes M$.

For (b) \implies (c), the identity map on M induces a morphism $\varepsilon: \Lambda[0] \rightarrow \mathrm{RHom}(M, M) \xrightarrow{\sim} M \otimes \mathbf{D}M$ (the coevaluation map). The required inverse to $N \otimes \mathbf{D}M \rightarrow \mathrm{RHom}(M, N)$ is

$$\mathrm{RHom}(M, N) \xrightarrow{\mathrm{id} \otimes \varepsilon} \mathrm{RHom}(M, N) \otimes M \otimes \mathbf{D}M \rightarrow N \otimes \mathbf{D}M.$$

For (c) \implies (d), we use the fact that \otimes commutes with colimits.

For (d) \implies (e), we use the fact that compact objects of $D(\Lambda)$ are perfect [Sta21, Tag 07LT].

Finally, for (e) implies (a), we can write M as a bounded complex of finitely generated projective Λ -modules. Then duals and derived tensor products can be computed on the level of chain complexes. We are reduced to showing, for finitely generated projective Λ -modules A and B , that $A \rightarrow A^{**}$ and $A^* \otimes B \rightarrow (A \otimes B^*)^*$ are isomorphisms. After localizing on Λ , we may assume that A and B are free of finite rank (since duals commute over direct sums), where these statements are easy to check. \square

We thank Bhargav Bhatt for helping us with the above proof.

B.2 Sheaves on locally profinite sets

Let S be a locally profinite set and Λ a discrete ring. We have the ring $C(S, \Lambda)$ of locally constant functions on S and the ring $C_c(S, \Lambda)$ of locally constant compactly supported functions on S . For each compact open subset $U \subset S$ let $\mathbf{1}_U$ denote the characteristic function. Then $C(U, \Lambda)$ is a principal ideal of both $C_c(S, \Lambda)$ and $C(S, \Lambda)$ generated by $\mathbf{1}_U$. Multiplication by $\mathbf{1}_U$ is a homomorphism $C(S, \Lambda) \rightarrow C(U, \Lambda)$ of rings with unity. In this way every $C(U, \Lambda)$ -module becomes a $C(S, \Lambda)$ -module.

Definition B.2.1. We call a $C(S, \Lambda)$ -module M

1. *smooth* if it satisfies the following equivalent conditions
 - (a) The multiplication map $M \otimes_{C_c(S, \Lambda)} C(S, \Lambda) \rightarrow M$ is an isomorphism.
 - (b) The natural map $\varinjlim (\mathbf{1}_U \cdot M) \rightarrow M$ is an isomorphism, where the colimit runs over the open compact subsets $U \subset S$ and the transition map $\mathbf{1}_U \cdot M \rightarrow \mathbf{1}_V \cdot M$ for $U \subset V$ is given by the natural inclusion.
2. *complete* if the natural map $M \rightarrow \varprojlim_U (\mathbf{1}_U \cdot M)$ is an isomorphism, where again U runs over the open compact subsets of S and the transition map $\mathbf{1}_U \cdot M \rightarrow \mathbf{1}_V \cdot M$ for $V \subset U$ is multiplication by $\mathbf{1}_V$.

Lemma B.2.2. *Let $V \subset S$ be compact open and let M be any $C(S, \Lambda)$ -module. Then*

1. $\mathbf{1}_V \cdot M$ is a submodule of $\varinjlim_U (\mathbf{1}_U \cdot M)$ and equals $\mathbf{1}_V \cdot \varinjlim (\mathbf{1}_U \cdot M)$.
2. $\mathbf{1}_V \cdot M$ is a submodule of $\varprojlim_U (\mathbf{1}_U \cdot M)$ and equals $\mathbf{1}_V \cdot \varprojlim (\mathbf{1}_U \cdot M)$.

Lemma B.2.3. 1. *The functor $M \mapsto M^s := \varinjlim (\mathbf{1}_U \cdot M)$ is a projector onto the category of smooth modules.*

2. *The functor $M \mapsto M^c := \varprojlim (\mathbf{1}_U \cdot M)$ is a projector onto the category of complete modules.*

3. *The two functors give mutually inverse equivalences of categories between the categories of smooth and complete modules.*

Let \mathcal{B} be the set of open compact subsets of S . Then \mathcal{B} is a basis for the topology of S and is closed under taking finite intersections and finite unions. Restriction gives an equivalence between the category of sheaves on S and the category of sheaves on \mathcal{B} . Define $R(U) = C(U, \Lambda)$. This is a sheaf of rings on S .

Let \mathcal{F} be an R -module sheaf on S . For $U \in \mathcal{B}$ we extend the $R(U)$ -module structure on $\mathcal{F}(U)$ to a $C(S, \Lambda)$ -module structure as remarked above. Then the restriction map $\mathcal{F}(S) \rightarrow \mathcal{F}(U)$ becomes a morphism of $C(S, \Lambda)$ -modules.

Lemma B.2.4. *1. For any $U \in \mathcal{B}$ the restriction map $\mathcal{F}(S) \rightarrow \mathcal{F}(U)$ is surjective and its restriction to $\mathbf{1}_U \cdot \mathcal{F}(S)$ is an isomorphism $\mathbf{1}_U \cdot \mathcal{F}(S) \rightarrow \mathcal{F}(U)$.*

2. *We have $\mathcal{F}(S) = \varprojlim_U \mathcal{F}(U)$, where the transition maps are the restriction maps.*

Let M be an $C(S, \Lambda)$ -module. Let $\mathcal{F}_M(U) = R(U)M = \mathbf{1}_U M$. This is a $C(S, \Lambda)$ -submodule of M . Given $V, U \in \mathcal{B}$ with $V \subset U$ we have the map $\mathcal{F}_M(U) \rightarrow \mathcal{F}_M(V)$ defined by multiplication by $\mathbf{1}_V$. In this way \mathcal{F}_M becomes an R -module sheaf.

Let $f : M \rightarrow N$ be a morphism of $C(S, \Lambda)$ -modules. We define for each U the morphism $f_U : \mathcal{F}_M(U) \rightarrow \mathcal{F}_N(U)$ simply by restricting f to $\mathcal{F}_M(U)$. One checks immediately that $(f_U)_U$ is a morphism of sheaves of R -modules. Therefore we obtain a functor from the category of $C(S, \Lambda)$ -modules to the category of sheaves of R -modules.

Given a sheaf \mathcal{F} on S we can define the smooth module $M_{\mathcal{F}}^s$ and the complete module $M_{\mathcal{F}}^c$ by

$$M_{\mathcal{F}}^s = \varinjlim_U \mathcal{F}(U) \quad M_{\mathcal{F}}^c = \varprojlim_U \mathcal{F}(U)$$

where the limit is taken over the restriction maps, and the colimit is taken over their sections given by Lemma B.2.4, and in both cases U runs over \mathcal{B} . Conversely, given any $C(S, \Lambda)$ -module M we have the sheaf \mathcal{F}_M .

Lemma B.2.5. *These functors give mutually inverse equivalences of categories from the category of smooth (resp. complete) $C(S, \Lambda)$ -modules to the category of R -module sheaves. These equivalences commute with the equivalence between the categories of smooth and complete modules. Furthermore $\mathcal{F}_M(S) = M^c$.*

C Some representation theory

Let G be a reductive group over a finite extension F/\mathbf{Q}_p . For a parabolic subgroup P of G we write i_P^G for the functor of normalized parabolic induction and r_G^P for the normalized Jacquet module functor.

Fix a minimal parabolic $P_0 = M_0 U_0$. A parabolic subgroup P is called standard if it contains P_0 . There is a unique Levi factor M of P that contains M_0 , and conversely M determines P . In that situation we may write i_M^G and r_G^M in place of i_P^G and r_G^P .

C.1 Non-elliptic representations

Our goal in this section is the following result.

Theorem C.1.1. *Let $\pi \in \text{Groth}(G(F))$ be any finite-length virtual $G(F)$ -representation with \mathbf{C} -coefficients, or with $\overline{\mathbf{Q}_\ell}$ -coefficients. Then π is non-elliptic if and only if it can be expressed as a \mathbf{Q} -linear virtual combination of representations induced from proper parabolic subgroups of G .*

When $G(F)$ has compact center, this is (a weaker version of) a classical result of Kazhdan [Kaz86]. The general statement seems to be well-known to experts, but we were unable to find an explicit formulation in the literature.

Proof. It suffices to treat the case of complex coefficients. Parabolic inductions are non-elliptic by van Dijk’s formula [vD72], so the “if” direction is clear. We will deduce the “only if” direction from [Dat00]; in what follows we freely use various notations from loc. cit., in particular writing $\mathcal{R}(G)$ for the Grothendieck group of finite length smooth \mathbf{C} -representations of $G(F)$.

Suppose that $\pi \in \mathcal{R}(G)$ is non-elliptic. Following the notation of [Dat00], pick any $f \in \overline{\mathcal{H}^{d(G)}}(G)$. Then all regular semisimple non-elliptic orbital integrals of f vanish by [Dat00, Theorem 3.2.iii], so $\text{tr}(f|\pi) = 0$ by our assumption on π and the Weyl integration formula. Therefore $\pi \in \overline{\mathcal{H}^{d(G)}}(G)^\perp$, so $\pi \in \mathcal{R}_{\mathbf{C}_{d(G)}}(G)$ by [Dat00, Theorem 3.2.ii]. Now applying [Dat00, Proposition 2.5.i] to the Hopf system $\mathcal{A}(-) = \mathcal{R}(-) \otimes \mathbf{Q}$ with $d = d(G)$, we see that $\pi \in \mathcal{R}(G) \otimes \mathbf{Q}$ is annihilated by the operator $1 - \sum_{d(M) > d(G)} c_d(M) i_M^G r_G^M$ for some rational numbers $c_d(M)$. Therefore

$$\pi = \sum_{d(M) > d(G)} c_d(M) i_M^G r_G^M(\pi),$$

and the right-hand side is a \mathbf{Q} -linear virtual combination of proper parabolic inductions, giving the result. \square

C.2 Integral representations and parabolic inductions

Fix a prime $\ell \neq p$. As usual, let $\text{Groth}(G(F))$ be the Grothendieck group of finite-length smooth $\overline{\mathbf{Q}_\ell}$ -representations of $G(F)$.

Definition C.2.1. Let π be an admissible smooth $\overline{\mathbf{Q}_\ell}$ -representation of $G(F)$. We say π admits a $\overline{\mathbf{Z}_\ell}$ -lattice if there exists an admissible smooth $\overline{\mathbf{Z}_\ell}[G(F)]$ -module L together with an isomorphism $L[1/\ell] \simeq \pi$.

Recall that our convention on the meaning of admissible is slightly non-standard, so in particular any such L has the property that L^K is a finite free $\overline{\mathbf{Z}_\ell}$ -module for all open compact pro- p subgroups $K \subset G(F)$, and whence L is a free $\overline{\mathbf{Z}_\ell}$ -module. The existence of a $\overline{\mathbf{Z}_\ell}$ -lattice in our sense implies, but is strictly stronger than, the existence of a “ $\overline{\mathbf{Z}_\ell}G(F)$ -réseau” in the sense of [Vig96]. Note also that if π is a finite-length admissible representation admitting a $\overline{\mathbf{Z}_\ell}$ -lattice, then any such lattice is finitely generated as a $\overline{\mathbf{Z}_\ell}[G(F)]$ -module by [Vig04].

The goal of this section is to prove the following result.

Proposition C.2.2. *The group $\text{Groth}(G(F))$ is generated by representations of the form $i_M^G(\sigma \otimes \psi)$, where $i_M^G(-)$ is the normalized parabolic induction functor associated with a standard Levi subgroup M , ψ is an unramified character of $M(F)$, and σ is an irreducible admissible $\overline{\mathbf{Q}_\ell}$ -representation of $M(F)$ admitting a $\overline{\mathbf{Z}_\ell}$ -lattice.*

We will deduce this from Dat's theory of ν -tempered representations [Dat05]. In particular, we will apply the theory from [Dat05] with $\mathcal{K} = \overline{\mathbf{Q}_\ell}$ or with $\mathcal{K} = E \subset \overline{\mathbf{Q}_\ell}$ a finite extension of \mathbf{Q}_ℓ , equipped with the usual norms, so ν is a positive multiple of the usual ℓ -adic valuation.

Lemma C.2.3. *Let π be any irreducible smooth $\overline{\mathbf{Q}_\ell}$ -representation of $G(F)$. If π is ν -tempered, then π admits a $\overline{\mathbf{Z}_\ell}$ -lattice.*

Proof. Suppose given π as in the lemma. By [Vig96, II.4.7], we may choose some E and some admissible E -representation π_E together with an isomorphism $\pi_E \otimes_E \overline{\mathbf{Q}_\ell} = \pi$. By definition, π is ν -tempered if and only if π_E is ν -tempered, [Dat05, Lemma 3.3]. Since π_E is ν -tempered, it admits an \mathcal{O}_E -lattice L by [Dat05, Proposition 6.3]. Then $L \otimes_{\mathcal{O}_E} \overline{\mathbf{Z}_\ell}$ is the desired $\overline{\mathbf{Z}_\ell}$ -lattice in π . \square

We will now freely use all the notation and results of [Dat05, §2-3], with $\mathcal{K} = \overline{\mathbf{Q}_\ell}$. A triple (M, σ, ψ) consisting of a standard Levi subgroup $M \subset G$, a ν -tempered irreducible representation σ of $M(F)$, and an unramified character ψ of $M(F)$ with $-\nu(\psi) \in (\mathfrak{a}_P)^{*,+}$, is called a Langlands triple. The corresponding representation $i_P^G(\sigma \otimes \psi)$ has a unique irreducible quotient, which we will denote by $j_P^G(\sigma \otimes \psi)$. Every irreducible smooth representation π of $G(F)$ is isomorphic to $j_P^G(\sigma \otimes \psi)$ for a (essentially) unique Langlands triple, cf. [Dat05, Theorem 3.11]. The uniqueness of the triple (M, σ, ψ) with a given irreducible quotient π allows us to index the representation $i_P^G(\sigma \otimes \psi)$ by π . We shall write $I(\pi)$ for this representation, and refer to it as the *standard representation* associated with π . Note that there is a natural surjection $I(\pi) \rightarrow \pi$.

On the other hand, by [Dat05, Theorem 3.11.ii], $\lambda_\pi := -\nu(\psi) \in \mathfrak{a}_{M_0}^*$ is also a well-defined invariant of π . Note that π is ν -tempered if and only if $\lambda_\pi = 0$, and that M can be read off from λ_π . The following key lemma is the analogue of [BW00, Lemma XI.2.13] in our setting.

Lemma C.2.4. *Let π be any irreducible representation. Write $\pi = j_P^G(\sigma \otimes \psi)$, and let π' be any nonzero irreducible subquotient of $I(\pi) = i_M^G(\sigma \otimes \psi)$. Then $\lambda_{\pi'} \leq \lambda_\pi$ in the usual partial ordering on $\mathfrak{a}_{M_0}^*$, and $\lambda_{\pi'} < \lambda_\pi$ if π' is a subquotient of $\ker(I(\pi) \rightarrow \pi)$.*

Proof. After twisting, we may assume π and π' have integral central characters. Write $\pi' = j_Q^G(\sigma' \otimes \psi')$ for some Langlands triple (L, σ', ψ') . By the proof of [Dat05, Theorem 3.11.i], $\lambda_{\pi'}$ occurs in $-\nu(\mathcal{E}(A_L, r_G^{\overline{\mathbf{Q}_\ell}}(\pi')))$, so the result now follows from the subsequent proposition. \square

Proposition C.2.5. *Let $MU = P$ and $LN = Q$ be standard parabolic subgroups of G . Let σ be a ν -tempered irreducible representation of $M(F)$, and $\psi : M(F) \rightarrow \overline{\mathbf{Q}_\ell}^\times$ an unramified character with $\mu = -\nu(\psi) \in (\mathfrak{a}_P^G)^{*,+}$. Let π' be a subquotient of $i_P^G(\sigma \otimes \psi)$ and let $\mu' \in -\nu(\mathcal{E}(A_L, r_G^{\overline{\mathbf{Q}_\ell}}(\pi')))$. Then*

1. $\mu' \leq \mu$.
2. If π' is a subquotient of $\ker(i_P^G(\sigma \otimes \psi) \rightarrow j_P^G(\sigma \otimes \psi))$, then $\mu' < \mu$.

Proof. The exponents of π' are a subset of the exponents of $r_G^{\overline{Q}}(i_P^G(\sigma \otimes \psi))$. These were analyzed in the proof of [Dat05, Lemma 3.7], where it was shown that if ψ' is such an exponent and $\mu' = -\nu(\psi')$ is such an exponent, then

$$\mu - \mu' = \mu - [\mu]_L^G + [\mu - w^{-1}\mu]_L^G - \overline{+(\mathfrak{a}_Q^G)^*},$$

for some $w \in W_M \setminus W/W_L$. It was moreover shown that $\mu - [\mu]_L^G$ and $[\mu - w^{-1}\mu]_L^G$ belong to $\overline{+(\mathfrak{a}_M^G)^*}$, which shows $\mu - \mu' \geq 0$, hence (1).

For (2) we may replace π' with $\ker(i_P^G(\sigma \otimes \psi) \rightarrow j_P^G(\sigma \otimes \psi))$, since the exponents of the former are again a subset of the exponents of the latter. We assume by way of contradiction that $\mu = \mu'$. We have $\mu \in (\mathfrak{a}_P^G)^{*,+}$ and $\mu' \in (\mathfrak{a}_L^G)^*$, so $\mu' = \mu$ implies that the intersection $(\mathfrak{a}_P^G)^* \cap (\mathfrak{a}_L^G)^*$ is non-empty. Since $(\mathfrak{a}_P^G)^{*,+}$ is an open subset of $(\mathfrak{a}_M^G)^*$, we see that $(\mathfrak{a}_M^G)^* \subset (\mathfrak{a}_L^G)^*$, hence $Q \subset P$. According to the formula $r_G^{\overline{Q}}(\pi') = r_M^{\overline{Q} \cap M}(r_G^{\overline{P}}(\pi'))$, $\mu' \in -\nu(\mathcal{E}(A_M, r_G^{\overline{P}}(\pi')))$.

By construction of the Langlands quotient $j_P^G(\sigma \otimes \psi)$ we have the exact sequence

$$0 \rightarrow \pi' \rightarrow i_P^G(\sigma \otimes \psi) \rightarrow i_{\overline{P}}^G(\sigma \otimes \psi),$$

where the map between the two parabolic inductions is the intertwining operator $J_{\overline{P}, P}$ of [Dat05, Lemma 3.7]. We recall that this intertwining operator was obtained via Frobenius reciprocity from the unique (up to scalar) element of $\text{Hom}_M(r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)), \sigma \otimes \psi)$. This element is the unique retraction of the natural embedding of $\sigma \otimes \psi$ into $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi))$.

We can describe this element in a slightly different way that is more suitable for our purposes. The representation $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi))$ has a filtration indexed by elements of $W_M \setminus W_G/W_M$ (strictly speaking, one has to choose a total order that refines the Bruhat order), and the natural embedding of $\sigma \otimes \psi$ into $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi))$ identifies $\sigma \otimes \psi$ with the beginning part of this filtration, indexed by $w = 1$. It is shown in equation (3.9) of the proof of [Dat05, Lemma 3.7] that for any exponent ψ'' of a subquotient corresponding to $w \neq 1$, $\mu'' = -\nu(\psi'')$ satisfies $\mu'' < \mu$. On the other hand, all exponents of $\sigma \otimes \psi$ have image μ under $-\nu$. Therefore, the retraction $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)) \rightarrow \sigma \otimes \psi$ is simply the projection onto the μ -direct summand of the the exponent decomposition of $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi))$.

Applying $r_G^{\overline{P}}$ to the above displayed exact sequence we obtain the exact sequence

$$0 \rightarrow r_G^{\overline{P}}(\pi') \rightarrow r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)) \rightarrow r_G^{\overline{P}}(i_{\overline{P}}^G(\sigma \otimes \psi)).$$

Therefore $r_G^{\overline{P}}(\pi')$, being the kernel of $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)) \rightarrow r_G^{\overline{P}}(i_{\overline{P}}^G(\sigma \otimes \psi))$, is contained in the kernel of the composition of this map with the evaluation-at-1 map $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)) \rightarrow \sigma \otimes \psi$. But that composition is, by construction of $J_{\overline{P}, P}$ via Frobenius reciprocity, equal to the projection $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)) \rightarrow (r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi)))_{\mu}$. Thus the exponents of the kernel of that projection are those exponents of $r_G^{\overline{P}}(i_P^G(\sigma \otimes \psi))$ whose image μ'' under $-\nu$ is not equal to μ . By what was said in the previous paragraph, these satisfy $\mu'' < \mu$. \square

Proof of Proposition C.2.2. Fix a point θ in the Bernstein variety for $G(F)$, and let $\text{Irr}(G(F))_{\theta} \subset \text{Irr}(G(F))$ be the finite set of irreducible representations with cuspidal support θ . Let

$$\text{Groth}(G(F))_{\theta} \subset \text{Groth}(G(F))$$

be the subgroup generated by $\text{Irr}(G(F))_\theta$, so

$$\text{Groth}(G(F)) = \oplus_\theta \text{Groth}(G(F))_\theta.$$

By Lemma C.2.3, it suffices to prove that $\text{Groth}(G(F))_\theta$ is generated by representations $i_M^G(\sigma \otimes \psi)$ for Langlands triples (M, σ, ψ) , i.e. by standard representations. Note that $\pi \in \text{Irr}(G(F))_\theta$ implies $I(\pi) \in \text{Groth}(G(F))_\theta$, cf. [BDK86, Proposition 2.4]. We will prove the finer result that the standard representations $I(\pi), \pi \in \text{Irr}(G(F))_\theta$ give a basis for $\text{Groth}(G(F))_\theta$.

Set

$$S = \{\lambda_\pi, \pi \in \text{Irr}(G(F))_\theta\} \subset \mathfrak{a}_{M_0}^*.$$

Note that S is a finite set, and inherits a natural partial order from the partial order on $\mathfrak{a}_{M_0}^*$. Pick any $\pi \in \text{Irr}(G(F))_\theta$. If λ_π is minimal in S , then the natural map $I(\pi) \rightarrow \pi$ is an isomorphism by Lemma C.2.4. In general, if λ_π is not minimal in S , then by Lemma C.2.4 and induction on S , we may assume that $\ker(I(\pi) \rightarrow \pi)$ is a \mathbf{Z} -linear combination of standard representations $I(\pi'), \pi' \in \text{Irr}(G(F))_\theta$. Then also $\pi = I(\pi) - \ker(I(\pi) \rightarrow \pi)$ is a \mathbf{Z} -linear combination of standard representations, giving the desired result. \square

References

- [BDK86] J. Bernstein, P. Deligne, and D. Kazhdan, *Trace Paley-Wiener theorem for reductive p -adic groups*, J. Analyse Math. **47** (1986), 180–192. MR 874050
- [BS17] Bhargav Bhatt and Peter Scholze, *Projectivity of the Witt vector affine Grassmannian*, Invent. Math. **209** (2017), no. 2, 329–423. MR 3674218
- [BT72] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local*, Inst. Hautes Études Sci. Publ. Math. (1972), no. 41, 5–251.
- [BT84] ———, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. (1984), no. 60, 197–376.
- [BW00] A. Borel and N. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, second ed., Mathematical Surveys and Monographs, vol. 67, American Mathematical Society, Providence, RI, 2000. MR 1721403
- [CFS17] Miaofen Chen, Laurent Fargues, and Xu Shen, *On the structure of some p -adic period domains*, 2017.
- [CS17] Ana Caraiani and Peter Scholze, *On the generic part of the cohomology of compact unitary Shimura varieties*, Ann. of Math. (2) **186** (2017), no. 3, 649–766. MR 3702677
- [Dat00] J.-F. Dat, *On the K_0 of a p -adic group*, Invent. Math. **140** (2000), no. 1, 171–226. MR 1779801
- [Dat05] ———, *ν -tempered representations of p -adic groups. I. l -adic case*, Duke Math. J. **126** (2005), no. 3, 397–469. MR 2120114
- [DKV84] P. Deligne, D. Kazhdan, and M.-F. Vignéras, *Représentations des algèbres centrales simples p -adiques*, Representations of reductive groups over a local field, Travaux en Cours, Hermann, Paris, 1984, pp. 33–117. MR 771672
- [Fal94] Gerd Faltings, *The trace formula and Drinfeld’s upper halfplane*, Duke Math. J. **76** (1994), no. 2, 467–481.
- [Fara] L. Fargues, *G -torseurs en théorie de Hodge p -adique*, To appear in Compositio Math.
- [Farb] ———, *Geometrization of the local Langlands correspondence: an overview*.
- [FF] L. Fargues and J.-M. Fontaine, *Courbes et fibrés vectoriels en théorie de Hodge p -adique*, To appear in Astérisque.
- [FKS19] Jessica Fintzen, Tasho Kaletha, and Loren Spice, *A twisted Yu construction, Harish-Chandra characters, and endoscopy*, arXiv:1912.03286 (2019).
- [FS21] Laurent Fargues and Peter Scholze, *Geometrization of the local Langlands correspondence*, preprint, 2021.

- [GHW] Daniel Gulotta, David Hansen, and Jared Weinstein, *An enhanced six functor formalism for diamonds*, In preparation.
- [Har15] Michael Harris, *Mathematics without apologies: Portrait of a problematic vocation*, Science Essentials, Princeton University Press, 2015.
- [Ima] Naoki Imai, *Convolution morphisms and Kottwitz conjecture*.
- [Kal13] Tasho Kaletha, *Genericity and contragredience in the local Langlands correspondence*, Algebra Number Theory **7** (2013), no. 10, 2447–2474.
- [Kal14] ———, *Supercuspidal L -packets via isocrystals*, Amer. J. Math. **136** (2014), no. 1, 203–239.
- [Kal16a] ———, *The local Langlands conjectures for non-quasi-split groups*, Families of automorphic forms and the trace formula, Simons Symp., Springer, [Cham], 2016, pp. 217–257. MR 3675168
- [Kal16b] ———, *Rigid inner forms of real and p -adic groups*, Ann. of Math. (2) **184** (2016), no. 2, 559–632. MR 3548533
- [Kal18] ———, *Rigid inner forms vs isocrystals*, J. Eur. Math. Soc. (JEMS) **20** (2018), no. 1, 61–101. MR 3743236
- [Kal19a] ———, *Regular supercuspidal representations*, J. Amer. Math. Soc. **32** (2019), no. 4, 1071–1170. MR 4013740
- [Kal19b] ———, *Supercuspidal L -packets*, arXiv:1912.03274 (2019).
- [Kaz86] David Kazhdan, *Cuspidal geometry of p -adic groups*, J. Analyse Math. **47** (1986), 1–36. MR 874042
- [Kot] Robert E. Kottwitz, *$B(G)$ for all local and global fields*, arXiv:1401.5728.
- [Kot83] ———, *Sign changes in harmonic analysis on reductive groups*, Trans. Amer. Math. Soc. **278** (1983), no. 1, 289–297.
- [Kot84a] ———, *Shimura varieties and twisted orbital integrals*, Math. Ann. **269** (1984), no. 3, 287–300.
- [Kot84b] ———, *Stable trace formula: cuspidal tempered terms*, Duke Math. J. **51** (1984), no. 3, 611–650.
- [Kot85] ———, *Isocrystals with additional structure*, Compositio Math. **56** (1985), no. 2, 201–220.
- [Kot86] ———, *Stable trace formula: elliptic singular terms*, Math. Ann. **275** (1986), no. 3, 365–399.
- [Kot97] ———, *Isocrystals with additional structure. II*, Compositio Math. **109** (1997), no. 3, 255–339.

- [KS99] Robert E. Kottwitz and Diana Shelstad, *Foundations of twisted endoscopy*, Astérisque (1999), no. 255, vi+190.
- [LZ] Yifeng Liu and Weizhe Zheng, *Enhanced six operations and base change theorem for higher Artin stacks*, <https://arxiv.org/abs/1211.5948>.
- [LZ20] Qing Lu and Weizhe Zheng, *Categorical traces and a relative Lefschetz-Verdier formula*, arXiv e-prints (2020), arXiv:2005.08522.
- [Man04] Elena Mantovan, *On certain unitary group Shimura varieties*, no. 291, 2004, Variétés de Shimura, espaces de Rapoport-Zink et correspondances de Langlands locales, pp. 201–331. MR 2074715
- [Mie] Yoichi Mieda, *Lefschetz trace formula and ℓ -adic cohomology of Rapoport-Zink tower for $GS(4)$* , <http://www.ms.u-tokyo.ac.jp/~mieda/pdf/RZ-LTF.pdf>.
- [Mie12] Yoichi Mieda, *Lefschetz trace formula and ℓ -adic cohomology of Lubin-Tate tower*, Math. Res. Lett. **19** (2012), no. 1, 95–107.
- [Mie14a] ———, *Geometric approach to the local Jacquet-Langlands correspondence*, Amer. J. Math. **136** (2014), no. 4, 1067–1091.
- [Mie14b] ———, *Lefschetz trace formula for open adic spaces*, J. Reine Angew. Math. **694** (2014), 85–128.
- [Rap95] Michael Rapoport, *Non-Archimedean period domains*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), Birkhäuser, Basel, 1995, pp. 423–434.
- [RV14] Michael Rapoport and Eva Viehmann, *Towards a theory of local Shimura varieties*, Münster J. Math. **7** (2014), no. 1, 273–326.
- [RZ96] M. Rapoport and Th. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996.
- [Sch17] Peter Scholze, *Étale cohomology of diamonds*, preprint, 2017.
- [SGA77] *Cohomologie l -adique et fonctions L* , Lecture Notes in Mathematics, Vol. 589, Springer-Verlag, Berlin-New York, 1977, Séminaire de Géométrie Algébrique du Bois-Marie 1965–1966 (SGA 5), Edité par Luc Illusie.
- [She14] Xu Shen, *Cell decomposition of some unitary group rapoport-zink spaces*, Math. Ann. **360** (2014), 825–899.
- [Shi11] Sug Woo Shin, *Galois representations arising from some compact Shimura varieties*, Ann. of Math. (2) **173** (2011), no. 3, 1645–1741. MR 2800722
- [Shi12] ———, *On the cohomology of Rapoport-Zink spaces of EL -type*, Amer. J. Math. **134** (2012), no. 2, 407–452. MR 2905002

- [Sta21] The Stacks Project Authors, *Stacks Project*, <http://stacks.math.columbia.edu>, 2021.
- [Ste65] Robert Steinberg, *Regular elements of semisimple algebraic groups*, Inst. Hautes Études Sci. Publ. Math. (1965), no. 25, 49–80.
- [Str05] Matthias Strauch, *On the Jacquet-Langlands correspondence in the cohomology of the Lubin-Tate deformation tower*, Astérisque (2005), no. 298, 391–410, Automorphic forms. I.
- [Str08] ———, *Deformation spaces of one-dimensional formal modules and their cohomology*, Adv. Math. **217** (2008), no. 3, 889–951.
- [SW13] Peter Scholze and Jared Weinstein, *Moduli of p -divisible groups*, Camb. J. Math. **1** (2013), no. 2, 145–237.
- [SW20] ———, *Berkeley lectures on p -adic geometry*, Annals of Math. Studies, no. 207, Princeton University Press, 2020.
- [Tit79] J. Tits, *Reductive groups over local fields*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 29–69.
- [Var07] Yakov Varshavsky, *Lefschetz-Verdier trace formula and a generalization of a theorem of Fujiwara*, Geom. Funct. Anal. **17** (2007), no. 1, 271–319.
- [Var20] Yakov Varshavsky, *Local terms for transversal intersections*, arXiv e-prints (2020), arXiv:2003.06815.
- [vD72] G. van Dijk, *Computation of certain induced characters of p -adic groups*, Math. Ann. **199** (1972), 229–240. MR 338277
- [Vig96] Marie-France Vignéras, *Représentations l -modulaires d’un groupe réductif p -adique avec $l \neq p$* , Progress in Mathematics, vol. 137, Birkhäuser Boston, Inc., Boston, MA, 1996. MR 1395151
- [Vig04] ———, *On highest Whittaker models and integral structures*, Contributions to automorphic forms, geometry, and number theory, Johns Hopkins Univ. Press, Baltimore, MD, 2004, pp. 773–801. MR 2058628
- [Yu19] Jize Yu, *The integral geometric Satake equivalence in mixed characteristic*, arXiv e-prints (2019), arXiv:1903.11132.