RATIONAL MAPS WITH A PREPERIODIC CRITICAL POINT

XAVIER BUFF, ADAM L. EPSTEIN, AND SARAH KOCH

Abstract. We show that the set of conjugacy classes of cubic polynomials with a prefixed critical point, of preperiod $k \geq 1$, is an irreducible algebraic curve. We also establish an analogous result for quadratic rational maps. We then study a closely related question concerning the irreducibility (over $\mathbb{Q}$) of the set of conjugacy classes of unicritical polynomials, of degree $D \geq 2$, with a preperiodic critical point. Our proofs are purely algebraic.

Contents

Introduction 1
1. Cubic polynomials 2
2. Quadratic rational maps 6
3. Unicritical polynomials 9
References 18

Introduction

Let $f : \mathbb{CP}^1 \to \mathbb{CP}^1$ be a rational map. A point $z \in \mathbb{CP}^1$ is

- periodic for f with period $n \geq 1$ if $f^{\circ n}(z) = z$ and n is the least such integer;
- preperiodic for f with preperiod $k \geq 0$ if $f^{\circ k}(z)$ is periodic for f and k is the least such integer.

The moduli space $\mathscr{P}_3$ of affine conjugacy classes of cubic polynomials is isomorphic to $\mathbb{C}^2$. Similarly, the moduli space $\mathscr{M}_2$ of Möbius conjugacy classes of quadratic rational maps is isomorphic to $\mathbb{C}^2$. In both cases, requiring that one critical point is preperiodic to a cycle of period $n \geq 1$ with preperiod $k \geq 0$ (with $k \neq 1$ in the case of quadratic rational maps) defines an algebraic curve. In [M1] and [M2], John Milnor introduced these curves and raised various questions about their geometry. In this article, we prove that the curves consisting of those maps with a prefixed critical point are irreducible.

We first study the case of cubic polynomials. Given $k \geq 0$ and $n \geq 1$, the affine conjugacy classes of cubic polynomials with a critical point preperiodic to a cycle of period n with preperiod k form an algebraic curve $\mathscr{S}_{k,n} \subset \mathscr{P}_3$. The following conjecture goes back to John Milnor [M2, Question 5.3] in the case $k = 0$.

Conjecture. *For $k \geq 0$ and $n \geq 1$, the curve $\mathscr{S}_{k,n}$ is irreducible.*

The research of the first author was supported in part by the ANR grant Lambda ANR-13-BS01-0002.

The research of the third author was supported in part by the NSF.

A proof in the case $k = 0$ has recently been announced by Matthieu Arfeux and Jan Kiwi [AK]; it relies on a result of Mary Rees in [R], that the set of fixed points of an endomorphism on a certain Teichmüller space is connected. We prove the following result.

Theorem 1. *For $k \geq 0$, the curve $\mathscr{S}_{k,1}$ is irreducible.*

Our proof is purely algebraic. It is largely inspired by the proof of Thierry Bousch [Bo] that for $n \geq 1$, the set of $(c, z) \in \mathbb{C}^2$ such that z is periodic of period n for $f_c : w \mapsto w^2 + c$ is irreducible. The proof will be given in §1.

In §2, we explain how the proof presented for cubic polynomials adapts to the case of quadratic rational maps. Given $k \geq 0$ with $k \neq 1$ and $n \geq 1$, the Möbius conjugacy classes of quadratic rational maps with a critical point preperiodic to a cycle of period n, with preperiod k, form an algebraic curve $\mathscr{V}_{k,n} \subset \mathscr{M}_2$.

Conjecture. *For $n \geq 1$, the curve $\mathscr{V}_{0,n}$ is irreducible. For $k \geq 2$ and $n \geq 1$, the curve $\mathscr{V}_{k,n}$ is irreducible.*

In this article, we prove the following result.

Theorem 2. *For $k \geq 2$, the curve $\mathscr{V}_{k,1}$ is irreducible.*

The proofs of Theorems 1 and 2 rely on the following result due to Vefa Goksel [G]. Assume $D \in \{2, 3\}$. Let $b_1 \in \mathbb{C}$ and $b_2 \in \mathbb{C}$ be two algebraic numbers such that 0 is preperiodic to a fixed point of $z \mapsto z^D + b_1$ and $z \mapsto z^D + b_2$, with the same preperiod $k \geq 2$. Then, b_1 and b_2 are Galois conjugate.

More generally, if $D \geq 2$ is an integer, the unicritical polynomials $z \mapsto z^D + b_1$ and $z \mapsto z^D + b_2$ are affine conjugate if and only if $b_1^{D-1} = b_2^{D-1}$. John Milnor [M3] asked whether one can classify the Galois conjugacy classes of parameters b^{D-1} such that the critical point of $z \mapsto z^D + b$ is preperiodic. In §3, we characterize those Galois conjugacy classes when the period is 1 or 2 for any prime power $D = p^e$, and when the period is 3 for $D = 2$ and $D = 8$.

1. Cubic polynomials

Every cubic polynomial is affine conjugate to a polynomial of the form

$$F_{a,b}(z) = z^3 - 3a^2 z + 2a^3 + b, \quad (a, b) \in \mathbb{C}^2.$$

Those polynomials have critical points at $\pm a$ and $b = F_{a,b}(a)$ is a critical value. A conjugacy between two such polynomials either preserves or exchanges the two critical points. Consequently, the moduli space $\mathscr{P}_3$ is obtained by identifying (a, b) with $(-a, -b)$. It follows that in order to prove Theorem 1, it is enough to show that the set $\mathcal{S}_k$ of parameters $(a, b) \in \mathbb{C}^2$ such that a is preperiodic to a fixed point with preperiod $k \geq 0$ is irreducible.

Note that for $k = 0$, the critical point a is fixed if and only if (a, b) belongs to the line $\mathcal{L}_0 := \{b = a\} \subset \mathbb{C}^2$. Thus, $\mathcal{S}_0 = \mathcal{L}_0$ is irreducible.

Note that for $k = 1$, the critical value $b = F_{a,b}(a)$ is fixed if and only if

$$b = F_{a,b}(b) = b^3 - 3a^2 b + 2a^3 = b + (a - b)^2(2a + b).$$

Consequently, $\mathcal{S}_1 = \mathcal{L}_1 \smallsetminus \mathcal{L}_0 = \mathcal{L}_1 \smallsetminus \big\{(0, 0)\big\}$, with $\mathcal{L}_1 := \{b = -2a\} \subset \mathbb{C}^2$. Thus, $\mathcal{S}_1$ is irreducible.

For the remainder of §1, we assume that $k \geq 2$.

1.1. **An equation for $\mathcal{S}_k$.** On the one hand, if a is preperiodic to a fixed point of $F_{a,b}$ with preperiod k, then the points $F_{a,b}^{\circ(k-1)}(a)$ and $F_{a,b}^{\circ k}(a)$ are distinct and have the same image under $F_{a,b}$. For $j \geq 0$, let $P_j \in \mathbb{Z}[a,b]$ be the polynomial defined by

$$P_j(a,b) := F_{a,b}^{\circ j}(a).$$

Then,

$$P_0(a,b) = a, \quad P_1(a,b) = b, \quad \text{and} \quad P_{j+1} = P_j^3 - 3a^2 P_j + 2a^3 + b,$$

so that for $j \geq 1$, the polynomial P_j has degree 3^{j-1}. Note that

$$F_{a,b}(z) - F_{a,b}(w) = (z-w)H(z,w) \quad \text{with} \quad H(z,w) = z^2 + zw + w^2 - 3a^2.$$

Thus, the polynomial

$$Q_k := H(P_{k-1}, P_k) \in \mathbb{Z}[a,b]$$

has degree $2 \cdot 3^{k-1}$ and vanishes on $\mathcal{S}_k$.

On the other hand, $H(z,z) = 0$ if and only if $z^2 = a^2$, i.e. $z = \pm a$. In particular, if $a = F_{a,b}(a)$, i.e. if $a = b$, then $P_{k-1}(a,b) = P_k(a,b) = a$ and $Q_k(a,b) = 0$. Thus, $b - a$ divides Q_k and so,

$$Q_k = (b-a)R_k \quad \text{with} \quad R_k \in \mathbb{Z}[a,b].$$

The polynomial R_k has degree $2 \cdot 3^{k-1} - 1$ and vanishes on $\mathcal{S}_k$. Set

$$\Sigma_k := \big\{(a,b) \in \mathbb{C}^2 \ ; \ R_k(a,b) = 0\big\}.$$

Then, $\mathcal{S}_k \subset \Sigma_k$. Note that there are points in $\Sigma_k \smallsetminus \mathcal{S}_k$:

(i) either $F_{a,b}^{\circ(k-1)}(a) = F_{a,b}^{\circ k}(a) = a$ in which case a is fixed;
(ii) or $F_{a,b}^{\circ(k-1)}(a) = F_{a,b}^{\circ k}(a) = -a$ in which case $-a$ is fixed and a is prefixed to $-a$ with preperiod j for some $j \in [\![2, k-1]\!]$.

Remark. Note that case (i) occurs if and only if $a = b = 0$, i.e., $\Sigma_k \cap \mathcal{S}_1 = \big\{(0,0)\big\}$. Indeed, for $j \geq 1$,

$$\frac{\partial P_{j+1}}{\partial b} = 3(P_j^2 - a^2)\frac{\partial P_j}{\partial b} + 1.$$

Since $P_j(a,a) = a$, it follows by induction that

$$\frac{\partial P_j}{\partial b}(a,a) = 1.$$

Since

$$\frac{\partial Q_k}{\partial b} = (2P_{k-1} + P_k)\frac{\partial P_{k-1}}{\partial b} + (P_{k-1} + 2P_k)\frac{\partial P_k}{\partial b},$$

we deduce that

$$R_k(a,a) = \frac{\partial Q_k}{\partial b}(a,a) = 6a.$$

Thus, on the line $\{a = b\} \subset \mathbb{C}^2$, the polynomial R_k only vanishes at $(0,0)$.

Theorem 1 is a corollary of the following result, the proof of which occupies the remainder of §1.

Proposition 3. *For $k \geq 2$, the polynomial $R_k \in \mathbb{Z}[a,b]$ is irreducible over $\mathbb{C}$.*



FIGURE 1. Three curves drawn in $\mathbb{R}^2$: $\mathcal{S}_0$ is red, $\mathcal{S}_1$ is blue, and $\mathcal{S}_2$ is green.

1.2. **Behavior near the origin.** We now show that in order to prove Proposition 3, it is enough to prove that R_k is irreducible over $\mathbb{Q}$.

Proposition 4. *The polynomial $R_k \in \mathbb{Z}[a,b]$ is irreducible over $\mathbb{C}$ if and only if it is irreducible over $\mathbb{Q}$.*

Proof. To prove the proposition, we use the following criterion.

Lemma 5. *Let $R \in \mathbb{Q}[a,b]$ be a polynomial vanishing at the origin with nonzero linear part. Then, R is irreducible over $\mathbb{C}$ if and only if R is irreducible over $\mathbb{Q}$.*

Proof. Clearly, if R is irreducible over $\mathbb{C}$, then it is irreducible over $\mathbb{Q}$.

Conversely, suppose that R is irreducible over $\mathbb{Q}$. We will show that R is irreducible over $\mathbb{C}$. Suppose that $R = S \cdot T$ where $S \in \mathbb{C}[a,b]$ is irreducible and vanishes at the origin. Such a polynomial S exists because R vanishes at the origin. It then follows that $T \in \mathbb{C}[a,b]$ does not vanish at the origin, since otherwise, the linear part of R at the origin would vanish. Multiplying S by a nonzero constant, we may assume that $T(0,0) = 1$. In that case, the linear parts of R and S at the origin coincide.

Since $R \in \mathbb{Q}[a,b]$, the polynomials S and T have algebraic coefficients. We claim that the coefficients of S are in fact rational. Indeed, assume $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Let S^σ be the image of S under the action of σ. Then, S^σ is an irreducible factor of R^σ and $R^\sigma = R$ since $R \in \mathbb{Q}[a,b]$. Note that S and S^σ are equal up to multiplication by a constant since otherwise, $S \cdot S^\sigma$ would divide R, and the linear part of R at the origin would vanish. In addition, the linear part of S^σ is equal to the linear part of $R^\sigma = R$. Thus, $S^\sigma = S$. Since this holds for all $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, the coefficients of S are rational.

Since $S \in \mathbb{Q}[a,b]$ is a factor of R and since R is irreducible over $\mathbb{Q}$, we have that $S = R$. This completes the proof since S is irreducible over $\mathbb{C}$ by assumption. □

To apply this lemma, we need to study the behavior of R_k at the origin.

Lemma 6. *The homogeneous part of least degree of R_k is $3(a+b)$.*

Proof. An elementary induction on $j \geq 1$ shows that the homogeneous part of least degree of P_j is b. As a consequence, the homogeneous part of least degree of Q_k is $3b^2 - 3a^2$. Factoring out $b - a$ to get R_k yields the required result. □

Thus, R_k vanishes at the origin with nonzero linear part $(a, b) \mapsto 3(a + b)$. This completes the proof of the proposition. □

What really matters in the proof of Lemma 5 is that the curve $\{R = 0\}$ has a single irreducible component containing the origin (indeed, since the derivative of R at the origin is non zero, the curve is smooth at the origin) and that the origin is fixed by the action of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. In fact, we have the following more general result (that we do not use in this article).

Lemma 7. *Let $R \in \mathbb{Q}[a, b]$ be a polynomial. Assume the affine curve $\{R = 0\}$ contains a point $(a_0, b_0) \in \mathbb{Q}^2$ and has a unique locally irreducible (over $\mathbb{C}$) branch at (a_0, b_0). Then R is irreducible over $\mathbb{C}$ if and only if R is irreducible over $\mathbb{Q}$.*

1.3. **The family $z^3 + b$, $b \in \mathbb{C}$.** We now study the intersection of $\mathcal{S}_k$ with the line $\mathcal{L}_2 := \{a = 0\} \subset \mathbb{C}^2$. Note that the map $f_b := F_{0,b}$ is a unicritical polynomial:

$$f_b(z) = z^3 + b.$$

For $j \geq 1$, define $p_j \in \mathbb{Z}[b]$ by

$$p_j(b) := P_j(0, b) \quad \text{so that} \quad p_1 = b \quad \text{and} \quad p_{j+1} = p_j^3 + b.$$

Let $q_k \in \mathbb{Z}[b]$ and $r_k \in \mathbb{Z}[b]$ be defined by

$$q_k(b) := Q_k(0, b) \quad \text{and} \quad r_k(b) := R_k(0, b),$$

so that

$$q_k = p_{k-1}^2 + p_{k-1}p_k + p_k^2 \quad \text{and} \quad q_k = br_k.$$

An easy induction on $j \geq 1$ shows that p_j is a monic polynomial of degree 3^{j-1} with least degree term b. It follows that q_k is a monic polynomial of degree $2 \cdot 3^{k-1}$ with least degree term $3b^2$. Thus, $q_k = br_k = b^2 s_k$ where $s_k \in \mathbb{Z}[b]$ is a monic polynomial of degree $2 \cdot 3^{k-1} - 2$ with $s_k(0) = 3$. The proof of the following result goes back to [G] (see also §3.2).

Proposition 8. *For $k \geq 2$, the polynomial $s_k \in \mathbb{Z}[b]$ is irreducible over $\mathbb{Q}$.*

Proof. Working in $\mathbb{F}_3[b]$, we have that $(x + y)^3 \equiv x^3 + y^3 \pmod 3$. An elementary induction on $j \geq 1$ yields

$$p_j \equiv b^{3^{j-1}} + b^{3^{j-2}} + \cdots + b^3 + b \pmod 3.$$

It follows that

$$p_k - p_{k-1} \equiv b^{3^{k-1}} \pmod 3 \quad \text{and} \quad (p_k - p_{k-1})q_k = (p_k - p_{k-1})^3 \equiv b^{3^k} \pmod 3.$$

Thus,

$$q_k \equiv b^{2 \cdot 3^{k-1}} \pmod 3 \quad \text{and} \quad s_k \equiv b^{2 \cdot 3^{k-1} - 2} \pmod 3.$$

Since $s_k(0) = 3$ is not a multiple of 9, the Eisenstein criterion implies that s_k is irreducible over $\mathbb{Q}$. □

1.4. **Behavior near infinity.** We now study the behavior of R_k when a or b is large.

Lemma 9. *The homogeneous part of greatest degree of R_k is*

$$(b-a)^{4\cdot 3^{k-2}-1}\cdot(2a+b)^{2\cdot 3^{k-2}}.$$

Proof. We first determine the homogeneous part H_k of greatest degree of P_j for $j \geq 2$. Since

$$P_2 = b^3 - 3a^2b + 2a^3 + b = (b-a)^2(2a+b) + b \quad \text{and} \quad P_{j+1} = P_j^3 - 3a^3P_j + 2a^3 + b,$$

we have $H_2 = (b-a)^2(2a+b)$ and an elementary induction on $j \geq 2$ yields that $H_j = (H_2)^{3^{j-2}}$. It follows that the homogeneous part of greatest degree of $Q_k = P_{k-1}^2 + P_{k-1}P_k + P_k^2 - 3a^2$ is $(H_2)^{2\cdot 3^{k-2}} = (b-a)^{4\cdot 3^{k-2}} \cdot (2a+b)^{2\cdot 3^{k-2}}$. Factoring out $b-a$ to get R_k yields the required result. □

Let us embed $\mathbb{C}^2$ in $\mathbb{CP}^2$ in the usual way, sending (a,b) to $[a:b:1]$.

Corollary 10. *The closure of Σ_k in $\mathbb{CP}^2$ intersects the line at infinity at only two points: $[1:1:0]$ with multiplicity $4\cdot 3^{k-2}-1$, and $[1:-2:0]$ with multiplicity $2\cdot 3^{k-2}$.*

1.5. **Irreducibility over $\mathbb{Q}$.** We may now complete the proof of Proposition 3.

Proposition 11. *For $k \geq 2$, the polynomial $R_k \in \mathbb{Z}[a,b]$ is irreducible over $\mathbb{Q}$.*

Proof. Assume by contradiction that $R_k = T_1 \cdot T_2$ with $T_1 \in \mathbb{Z}[a,b]$, $T_2 \in \mathbb{Z}[a,b]$, $\text{degree}(T_1) < \text{degree}(R_k)$, and $\text{degree}(T_2) < \text{degree}(R_k)$.

We first prove that either T_1 or T_2 must have degree 1. Let $t_1 \in \mathbb{Z}[b]$ and $t_2 \in \mathbb{Z}[b]$ be defined by

$$t_1(b) := T_1(0,b) \quad \text{and} \quad t_2(b) := T_2(0,b).$$

Then, $r_k = t_1 \cdot t_2$ with $\text{degree}(t_1) \leq \text{degree}(T_1) < \text{degree}(R_k) = \text{degree}(r_k)$. Similarly, $\text{degree}(t_2) < \text{degree}(r_k)$. Since $r_k = bs_k$ with r_k monic and s_k irreducible over $\mathbb{Q}$, exchanging T_1 and T_2 if necessary, this implies that $t_1 = \pm b$ and $t_2 = \pm s_k$. Then, $\text{degree}(T_2) \geq \text{degree}(s_k) = \text{degree}(R_k) - 1$ and $\text{degree}(T_1) = 1$.

According to Lemma 6, the homogeneous part of least degree of R_k is $3(a+b)$. Thus, T_1 divides $3(a+b)$; in fact, since $t_1 = \pm b$, we have that $T_1 = \pm(a+b)$. So, the closure of Σ_k in $\mathbb{CP}^2$ intersects the line at infinity at the point $[1:-1:0]$. This contradicts Corollary 10. □

2. Quadratic rational maps

To prove Theorem 2, it is convenient to work in a space of dynamically marked quadratic rational maps. A quadratic rational map whose conjugacy class belongs to $\mathscr{V}_{k,1}$ with $k \geq 2$ has a critical point ω whose orbit contains a fixed point α. There is a fixed point $\beta \neq \alpha$ since otherwise, α would be a triple fixed point and its parabolic basin would contain both critical orbits. Note that $\beta \neq \omega$ since ω is not fixed. The conjugacy class may therefore be represented by a rational map f such that

$$\alpha = 0, \quad \beta = \infty \quad \text{and} \quad \omega = 1.$$

The critical value $a = f(1)$ belongs to $\mathbb{C}\smallsetminus\{0\}$ and $f^{-1}(0) = \{0, b\}$ with $b \in \mathbb{C}\smallsetminus\{1\}$. So, the rational map is

$$G_{a,b}(z) := \frac{az(b-z)}{1+(b-2)z} \quad \text{with} \quad (a,b) \in \Lambda := \big(\mathbb{C}\smallsetminus\{0\}\big) \times \big(\mathbb{C}\smallsetminus\{1\}\big).$$

In addition, (a, b) belongs to the curve

$$\mathcal{V}_k := \big\{(a,b) \in \Lambda \ ; \ G_{a,b}^{\circ(k-2)}(a) = b\big\}.$$

Conversely, if (a, b) belongs to the curve $\mathcal{V}_k$, then the conjugacy class of $G_{a,b}$ belongs to $\mathscr{V}_{k,1}$. So, in order to prove Theorem 2, it is enough to prove that the curve $\mathcal{V}_k$ is irreducible.

Remark. A generic conjugacy class in $\mathscr{V}_{k,1}$ has two representatives in $\mathcal{V}_k$ corresponding to the choice of the marked fixed point β. It follows that the quotient map $\mathcal{V}_k \to \mathscr{V}_{k,1}$ has degree 2.

2.1. **An equation for $\mathcal{V}_k$.** Here, we define a polynomial $R_k \in \mathbb{Z}[a, b]$ vanishing on $\mathcal{V}_k$. This polynomial should not be confused with the polynomial R_k defined in §1. However, since they play parallel roles, we keep the same notation. Let us first observe that for $j \geq 2$,

$$G_{a,b}^{\circ(j-2)}(a) = \frac{P_j(a,b)}{Q_j(a,b)}$$

where $P_j \in \mathbb{Z}[a, b]$ and $Q_j \in \mathbb{Z}[a, b]$ are defined recursively by

$$P_2 = a, \quad Q_2 = 1, \quad P_{j+1} = aP_j \cdot (bQ_j - P_j) \quad \text{and} \quad Q_{j+1} = Q_j^2 + (b-2)P_jQ_j.$$

So, $\mathcal{V}_k$ is the set of parameters $(a, b) \in \Lambda$ such that

$$R_k(a,b) = 0 \quad \text{with} \quad R_k := P_k - bQ_k \in \mathbb{Z}[a,b].$$

This shows that $\mathcal{V}_k$ is an algebraic subset of Λ and that Theorem 2 follows from the following result.

Proposition 12. *For $k \geq 2$, the polynomial $R_k \in \mathbb{Z}[a, b]$ is irreducible over $\mathbb{C}$.*

Note that $R_2 = a - b$ is irreducible over $\mathbb{C}$. For the remainder of §2, devoted to the proof of Proposition 12, we assume that $k \geq 3$.

2.2. **Behavior near the origin.** As in §1.2, we first prove that it is enough to show that R_k is irreducible over $\mathbb{Q}$. And here also, we deduce this from Lemma 5, studying the behavior of R_k near the origin. There is however a fundamental difference between the two approaches, even if this does not appear in the proof. In the case of cubic polynomials, the origin corresponds to the cubic polynomial $z \mapsto z^3$ which belongs to the family we are studying, whereas here, the origin does not belong to our parameter space Λ.

Lemma 13. *For $k \geq 3$, the homogeneous part of least degree of R_k is $-b$.*

Proof. An elementary induction shows that for $j \geq 2$, the homogeneous part of least degree of P_j is $a^{j-1}b^{j-2}$ and the homogeneous part of least degree of Q_j is 1. The result follows immediately. □

As a consequence $R_k \in \mathbb{Z}[a, b]$ vanishes at the origin with nonzero linear part. According to Lemma 5, the polynomial R_k is irreducible over $\mathbb{C}$ if and only if it is irreducible over $\mathbb{Q}$.



FIGURE 2. Three curves drawn in $\mathbb{R}^2$: $\mathcal{V}_2$ is red, $\mathcal{V}_3$ is blue, and $\mathcal{V}_4$ is green.

2.3. **The family $az(2-z)$, $a \in \mathbb{C}$.** We now study the intersection of $\mathcal{V}_k$ with the line $\mathcal{L} := \{b = 2\} \subset \mathbb{C}^2$. Note that the map $g_a := G_{a,2}$ is a quadratic polynomial:

$$g_a(z) = az(2-z).$$

For $j \geq 2$, define $p_j \in \mathbb{Z}[a]$ and $q_j \in \mathbb{Z}[a]$ by

$$p_j(a) := P_j(a, 2), \quad q_j(a) := Q_j(a, 2)$$

so that

$$p_2 = a, \quad p_{j+1} = -ap_j^2 + 2ap_j, \quad q_1 = 1 \quad \text{and} \quad q_{j+1} = q_j^2.$$

In particular, for $j \geq 3$, the polynomial $-p_j$ is monic with degree $2^{j-1} - 1$, and its constant coefficient is 0; and $q_j = 1$. Let $r_k \in \mathbb{Z}[a]$ be defined by

$$r_k(a) := R_k(a, 2) \quad \text{so that} \quad r_k = p_k - 2.$$

Then, r_k has degree $2^{k-1} - 1$ and its constant coefficient is -2.

Lemma 14. *The degree of R_k is $2^{k-1} - 1$.*

Proof. An elementary induction shows that the degree of P_k is at most $2^{k-1} - 1$ and the degree of Q_k is at most $2^{k-1} - 2$. Consequently, the degree of R_k is at most $2^{k-1} - 1$.

Since the polynomial p_k has degree $2^{k-1} - 1$, the polynomial $r_k = p_k - 2$ also has degree $2^{k-1} - 1$. Thus,

$$2^{k-1} - 1 = \text{degree}(r_k) \leq \text{degree}(R_k) \leq 2^{k-1} - 1$$

and the result follows. □

The proof of the following result goes back to [G] (see also §3.2).

Proposition 15. *For all $k \geq 2$, the polynomial $r_k \in \mathbb{Z}[a]$ is irreducible over $\mathbb{Q}$.*

Proof. Working in $\mathbb{F}_2[a]$, we have that for $j \geq 2$,

$$p_{j+1} \equiv a p_j^2 \pmod 2 \quad \text{so that} \quad r_k \equiv p_k \equiv a^{2^{k-1}} \pmod 2.$$

The constant coefficient of r_k is -2. It follows from the Eisenstein criterion that r_k is irreducible over $\mathbb{Q}$. □

2.4. **Irreducibility over $\mathbb{Q}$.** We may now complete the proof of Proposition 12.

Proposition 16. *The polynomial $R_k \in \mathbb{Z}[a,b]$ is irreducible over $\mathbb{Q}$.*

Proof. Assume by contradiction that $R_k = T_1 \cdot T_2$ with $T_1 \in \mathbb{Z}[a,b]$, $T_2 \in \mathbb{Z}[a,b]$, $\text{degree}(T_1) < \text{degree}(R_k)$ and $\text{degree}(T_2) < \text{degree}(R_k)$. Consider the polynomials $t_1 \in \mathbb{Z}[a]$ and $t_2 \in \mathbb{Z}[a]$ defined by

$$t_1(a) := T_1(a,2) \quad \text{and} \quad t_2(a) := T_2(a,2).$$

Then, $r_k = t_1 \cdot t_2$ with $\text{degree}(t_1) \leq \text{degree}(T_1) < \text{degree}(R_k) = \text{degree}(r_k)$. Similarly, $\text{degree}(t_2) < \text{degree}(r_k)$. This is not possible since r_k is irreducible over $\mathbb{Q}$. □

3. Unicritical polynomials

The previous discussion motivates a more systematic study of irreducibility over $\mathbb{Q}$ within families of unicritical polynomials. This section is devoted to such a study. It can be read independently of the rest of the article. Consider the polynomials $f_a : \mathbb{C} \to \mathbb{C}$ defined by

$$f_a(z) = az^D + 1, \quad a \in \mathbb{C}.$$

The polynomial f_a is unicritical: it has a unique critical point at $z = 0$. We are interested in parameters a such that the critical point is preperiodic for f_a. Note that the preperiod k cannot be equal to 1.

For $n \geq 1$, let $P_n \in \mathbb{Z}[a]$ be the polynomial

$$P_n(a) := f_a^{\circ n}(0).$$

Andrew Gleason observed that the discriminant of P_n is 1 (mod D), and thus P_n has simple roots. It follows that

$$P_n = \prod_{m|n} R_m \quad \text{with} \quad R_n := \prod_{m|n} P_m^{\mu(n/m)} \in \mathbb{Z}[a],$$

where μ is the Möbius function defined by $\mu(i) = (-1)^j$ if i is the product of j distinct primes with $j \geq 0$ and $\mu(i) = 0$ otherwise. For example,

$$R_1 = P_1 = 1, \quad R_2 = P_2 = a + 1 \quad \text{and} \quad R_3 = P_3 = a(a+1)^D + 1.$$

It is conjectured that when $D = 2$, the polynomials R_n are irreducible over $\mathbb{Q}$ for all $k \geq 2$. The following result shows that this is not true when $D \equiv 1 \pmod 6$.

Proposition 17 ([Bu])**.** *The polynomial R_3 is irreducible over $\mathbb{Q}$ if and only if D is not congruent to 1 modulo 6. When $D \equiv 1 \pmod 6$, the polynomial R_3 has exactly two irreducible factors over $\mathbb{Q}$, one of which is $a^2 + a + 1$.*

Assume now that 0 is preperiodic for f_a with preperiod $k \geq 2$ and period $n \geq 1$. Then,

$$f_a^{\circ(k+n-1)}(0) = \omega f_a^{\circ(k-1)}(0) \quad \text{with} \quad \omega^D = 1 \quad \text{and} \quad \omega \neq 1. \tag{1}$$

In fact, Equation (1) is satisfied if and only if either 0 is periodic for f_a with period dividing $\gcd(n, k-1)$, or 0 is preperiodic for f_a with preperiod k and period dividing n.

For $k \geq 2$, $n \geq 1$ and $d \geq 2$ dividing D, we therefore consider the monic polynomial $R_{k,n,d}$ whose roots are the parameters $a \in \mathbb{C}$ such that

- 0 is preperiodic for f_a with preperiod k and period n, and
- Equation (1) is satisfied for some primitive d-th root of unity ω.

We claim that $R_{k,n,d} \in \mathbb{Z}[a]$. Indeed, let $\Phi_d \in \mathbb{Z}[X, Y]$ be the (homogenized) d-th cyclotomic polynomial: if Ω_d is the set of primitive d-th roots of unity, then

$$\Phi_d := \prod_{\omega \in \Omega_d} (X - \omega Y).$$

Let $P_{k,n,d} \in \mathbb{Z}[a]$ be the polynomial defined by

$$P_{k,n,d} := \Phi_d(P_{k+n-1}, P_{k-1}) = \prod_{\omega \in \Omega_d} (P_{k+n-1} - \omega P_{k-1}).$$

The polynomial $P_{k+n-1} - \omega P_{k-1}$ has simple roots (see [Bu] for example). In addition, the common roots of P_{k+n-1} and P_{k-1} are the roots of $P_{\gcd(n,k-1)}$. It follows that the multiple roots of $P_{k,n,d}$ are the roots of $P_{\gcd(n,k-1)}$ with multiplicities $\varphi(d) = \deg(\Phi_d)$, where φ is the Euler totient function. As a consequence,

$$P_{k,n,d} = P_{\gcd(n,k-1)}^{\varphi(d)} \cdot \prod_{m|n} R_{k,m,d} \tag{2}$$

and according to the Möbius Inversion Formula,

$$R_{k,n,d} = \prod_{m|n} \left(\frac{P_{k,m,d}}{P_{\gcd(m,k-1)}^{\varphi(d)}} \right)^{\mu(n/m)} \in \mathbb{Z}[a].$$

We also consider the polynomials $P_{k,n} \in \mathbb{Z}[a]$ and $R_{k,n} \in \mathbb{Z}[a]$ defined by

$$P_{k,n} := \prod_{\substack{d|D \\ d \neq 1}} P_{k,n,d} = \frac{P_{k+n-1}^D - P_{k-1}^D}{P_{k+n-1} - P_{k-1}} = \sum_{i+j=D-1} P_{k+n-1}^i \cdot P_{k-1}^j.$$

and

$$R_{k,n} := \prod_{\substack{d|D \\ d \neq 1}} R_{k,n,d} \in \mathbb{Z}[a] \quad \text{so that} \quad P_{k,n} = P_{\gcd(n,k-1)}^{D-1} \cdot \prod_{m|n} R_{k,m}. \tag{3}$$

We study the following conjecture of John Milnor [M3] (compare with [HT]).

Conjecture. *For all $k \geq 2$, $n \geq 1$, and $d \geq 2$ that divide $D \geq 2$, the polynomial $R_{k,n,d}$ is irreducible over $\mathbb{Q}$.*

There are few cases where the expression of $R_{k,n,d}$ is sufficiently simple so that existing results in the literature directly apply (see §3.4).

Theorem 18 ([G])**.** *If D is a prime number, then $R_{k,1}(c^{D-1}) \in \mathbb{Z}[c]$ is irreducible for all $k \geq 2$. If $D = 2$, then $R_{k,2}$ is irreducible for all $k \geq 2$.*

We prove the following theorem. In the remainder of the article, p is a prime number.

Theorem 19. *Assume $D = p^e$ is a prime power. Then $R_{k,1,d}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$, and for all $d \geq 2$ that divide D. More generally, if $n \geq 2$ and the polynomial $R_n \pmod p$ is irreducible over $\mathbb{F}_p$, then $R_{k,n,d}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$, and for all $d \geq 2$ that divide D.*

Corollary 20. *Assume $D = p^e$ is a prime power. Then $R_{k,2,d}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$, and for all $d \geq 2$ that divide D.*

Proof. The reduction of $R_2 = a + 1$ modulo p is irreducible over $\mathbb{F}_p$. □

Corollary 21. *If $D = 2$ then $R_{k,3}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$.*

Proof. If $D = 2$, then $R_3 = a(a+1)^2 + 1 \equiv 1 + a + a^3 \pmod 2$ and $R_3 \pmod 2$ is irreducible over $\mathbb{F}_2$. □

Corollary 22. *If $D = 8$, then $R_{k,3,2}$, $R_{k,3,4}$ and $R_{k,3,8}$ are irreducible over $\mathbb{Q}$ for all $k \geq 2$.*

Proof. If $D = 8$, then $R_3 = a(a+1)^8 + 1 \equiv 1 + a + a^9 \pmod 2$ and $R_3 \pmod 2$ is irreducible over $\mathbb{F}_2$. □

Remark. The only values of $D = p^e$ and $n \geq 2$ for which the polynomial $R_n \pmod p$ is irreducible over $\mathbb{F}_p$ are the one listed previously: $n = 2$ for any prime power degree D, and $n = 3$ for both $D = 2$ and $D = 8$ (see §3.5).

Our proof of Theorem 19 relies on the following two results (see §3.3).

Lemma 23. *Assume $d \geq 2$ divides $D \geq 2$. Assume $k \geq 2$, $n \geq 1$ and $m \geq 1$. Then,*

$$\operatorname{resultant}(R_{k,m,d}, R_n) = \begin{cases} \pm p^{\deg(R_n)} & \text{if } n = m \text{ and } d = p^e \text{ is a prime power} \\ \pm 1 & \text{otherwise.} \end{cases}$$

Lemma 24. *Assume $D = p^e$ is a prime power and $d \geq 2$ is a divisor of D. Then for all $k \geq 2$, the polynomials $R_{k,1,d} \pmod p$ are powers of $a \in \mathbb{F}_p[a]$; and for all $k \geq 2$ and all $n \geq 2$, the polynomials $R_{k,n,d} \pmod p$ are powers of $R_n \pmod p$.*

Remark. Lemma 23 shows a connection between the polynomials $R_{k,n,d}$ and the polynomials R_n, valid for all degrees $D \geq 2$. Lemma 24 shows a stronger connection between these polynomials, but only valid for prime power degrees $D = p^e$. We think that it is worth investigating what this relation becomes when D is no longer a prime power.

3.1. **The critical orbit.** We first study some properties of the polynomials $P_k \in \mathbb{Z}[a]$. Recall that by definition, for all $k \geq 1$,

$$P_k(a) := f_a^{\circ k}(0).$$

For $k \geq 0$, set

$$N_k := \frac{D^k - 1}{D - 1} \quad \text{so that} \quad 1 + DN_k = \frac{D - 1 + D^{k+1} - D}{D - 1} = N_{k+1}.$$

Lemma 25. *For all $k \geq 1$, the polynomial P_k has constant coefficient 1 and is monic of degree N_{k-1}.*

Proof. First, note that $P_1 = 1$ and for all $k \geq 1$, $P_{k+1} = aP_k^D + 1$. It follows that the constant coefficient of P_{k+1} is 1. Second, let us prove by induction on $k \geq 1$ that P_k is monic of degree N_{k-1}. The property holds for $k = 1$: indeed, $P_1 = 1$ and $N_0 = 0$. Now, if the result holds for some integer $k \geq 1$, then $P_{k+1} = aP_k^D + 1$ is monic of degree $1 + DN_{k-1} = N_k$. □

Lemma 26. *Assume $D = p^e$ is a prime power. For all $k \geq 1$,*

$$P_{k+1} - P_k \equiv a^{N_k} \pmod{p}.$$

Proof. We prove the result by induction on $k \geq 1$. For $k = 1$,

$$P_2 - P_1 = a + 1 - 1 = a = a^{N_1}.$$

Now, assume the property holds for some $k \geq 1$. Since $D = p^e$,

$$\begin{aligned} P_{k+2} - P_{k+1} &= (aP_{k+1}^D + 1) - (aP_k^D + 1) \\ &= a \cdot (P_{k+1}^D - P_k^D) \equiv a \cdot (P_{k+1} - P_k)^D \pmod{p}. \end{aligned}$$

Thus,

$$P_{k+2} - P_{k+1} \equiv a^{1+DN_k} \pmod{p} \equiv a^{N_{k+1}} \pmod{p}.$$ □

We conclude this section by the following observation due to Poonen.

Lemma 27 (Poonen)**.** *For $m \neq n$, we have that* resultant$(R_m, R_n) = \pm 1$.

Proof. Assume $n > m$. It is not hard to see by induction on $k \geq 1$, that

$$P_{m+k} \equiv P_k \pmod{P_m^D}.$$

Indeed, $P_{m+1} = aP_m^D + 1 = P_1 + aP_m^D$ and if $P_{m+k} \equiv P_k \pmod{P_m^D}$, then

$$P_{m+k+1} = aP_{m+k}^D + 1 \equiv aP_k^D + 1 \pmod{P_m^D} \equiv P_k \pmod{P_m^D}.$$

This implies that, $P_{mn} \equiv P_m \pmod{P_m^D}$. Since $m < n$, $P_m R_n$ divides P_{mn}. So, there are polynomials $A \in \mathbb{Z}[a]$ and $B \in \mathbb{Z}[a]$ such that

$$AP_m R_n = P_{mn} = P_m + BP_m^D.$$

Dividing by P_m yields $AR_n - BP_m^{D-1} = 1$. It follows that R_m and R_n are relatively prime in $\mathbb{Z}[a]$ and resultant$(R_m, R_n) = \pm 1$. □

3.2. **When the critical point is preperiodic to a fixed point.** As a warm up, we first prove the following proposition that is due to Vefa Goksel. Our proof differs significantly from the one given in [G].

Proposition 28. *If D is prime, then $R_{k,1}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$.*

Proof. Our proof relies on the following two lemmas.

Lemma 29. *For $k \geq 2$ and $n \geq 1$, the polynomial $P_{k,n}$ has constant coefficient D and is monic of degree $(D-1)N_{k+n-2}$.*

Proof. By Lemma 25, if $i + j = D - 1$, the polynomial $P_{k+n-1}^i \cdot P_{k-1}^j$ has constant coefficient 1 and is monic of degree

$$i \cdot N_{k+n-2} + j \cdot N_{k-2} \leq (D-1)N_{k+n-2}$$

with equality if and only if $i = D - 1$ and $j = 0$. There are D pairs $(i, j) \in \mathbb{N}^2$ such that $i + j = D - 1$. Only one pair contributes to the leading term. Thus the polynomial is monic. Every pair contributes to the constant coefficient, which therefore is equal to D. □

Lemma 30. *If D is prime, then for all $k \geq 1$,*

$$R_{k,1} = P_{k,1} \equiv a^{(D-1)N_{k-1}} \pmod{D}.$$

Proof. Assume D is prime. On the one hand, according to Lemma 26:

$$P_k^D - P_{k-1}^D \equiv (P_k - P_{k-1})^D \pmod{D} \equiv a^{DN_{k-1}} \pmod{D}. \tag{4}$$

On the other hand, by definition of $P_{k,1}$:

$$P_k^D - P_{k-1}^D = (P_k - P_{k-1}) \cdot P_{k,1} \equiv a^{N_{k-1}} P_{k,1} \pmod{D}.$$

As a consequence,

$$a^{N_{k-1}} P_{k,1} \equiv a^{DN_{k-1}} \pmod{D} \quad \text{so that} \quad P_{k,1} \equiv a^{(D-1)N_{k-1}} \pmod{D}. \qquad \square$$

The proposition now follows from the Eisenstein criterion: $R_{k,1}$ is monic, D divides all the coefficients except the one of the leading term, and D^2 does not divide the constant coefficient. $\square$

3.3. **The general case.** This section is devoted to the proof of Theorem 19. We first prove Lemmas 23 and 24.

Proof of Lemma 23. Assume $d \geq 2$ divides $D \geq 2$, $k \geq 2$, $n \geq 1$ and $m \geq 1$. We need to show that

$$\operatorname{resultant}(R_{k,m,d}, R_n) = \begin{cases} \pm p^{\deg(R_n)} & \text{if } n = m \text{ and } d = p^e \text{ is a prime power} \\ \pm 1 & \text{otherwise.} \end{cases}$$

The proof splits in several cases.

Case 1: n does not divide m. Assume α is a root of R_n. Then, $P_{j_1}(\alpha) = P_{j_2}(\alpha)$ if and only if $j_1 \equiv j_2 \pmod{n}$. Since n does not divide m, for all $k \geq 2$,

$$P_{k+m-1}(\alpha) - P_{k-1}(\alpha) \neq 0 \quad \text{and} \quad \alpha P_{k,m}(\alpha) = \frac{P_{k+m}(\alpha) - P_k(\alpha)}{P_{k+m-1}(\alpha) - P_{k-1}(\alpha)},$$

so that

$$\alpha^n \prod_{j=0}^{n-1} P_{k+j,m}(\alpha) = 1.$$

The polynomial R_n is monic with constant coefficient 1. So, α is an algebraic unit. Thus,

$$\prod_{j=0}^{n-1} \operatorname{resultant}(P_{k+j,m}, R_n) = \prod_{j=0}^{n-1} \prod_{\alpha \in R_n^{-1}(0)} P_{k+j,m}(\alpha) = \prod_{j=0}^{n-1} \prod_{\alpha \in R_n^{-1}(0)} \frac{1}{\alpha^n} = \pm 1.$$

Since $R_{k,m,d}$ divides $P_{k,m}$, it follows that

$$\operatorname{resultant}(R_{k,m}, R_n) = \pm 1.$$

Case 2: n divides m. Set

$$\nu := \Phi_d(1,1) = \begin{cases} p & \text{if } d = p^e \text{ is a prime power} \\ 1 & \text{otherwise.} \end{cases}$$

It is enough to prove that

$$\prod_{\ell \mid m} \operatorname{resultant}(R_{k,\ell,d}, R_n) = \pm \nu^{\deg(R_n)}. \tag{5}$$

Indeed, assume Equation (5) holds. We have seen that $\text{resultant}(R_{k,\ell,d}, R_n) = \pm 1$ when n does not divide ℓ. So, for $m = n$,

$$\begin{aligned}\pm\nu^{\deg(R_n)} &= \text{resultant}(R_{k,n,d}, R_n) \cdot \prod_{\substack{\ell \mid n \\ \ell \neq n}} \text{resultant}(R_{k,\ell,d}, R_n) \\ &= \pm\text{resultant}(R_{k,n,d}, R_n).\end{aligned}$$

Now, if n divides $m \neq n$, the polynomial $R_{k,n,d} \cdot R_{k,m,d}$ divides $P_{k,m,d}$; and

$$\text{resultant}(R_{k,n,d} \cdot R_{k,m,d}, R_n) = \pm\nu^{\deg(R_n)} \cdot \text{resultant}(R_{k,m,d}, R_n)$$

divides

$$\text{resultant}(P_{k,m,d}, R_n) = \pm\nu^{\deg(R_n)}.$$

This forces

$$\text{resultant}(R_{k,m,d}, R_n) = \pm 1.$$

So, it is enough to prove that Equation (5) holds.

Case 2.a: n does not divide $k-1$. Assume α is a root of R_n. Since n divides m, we have that $P_{k+m-1}(\alpha) = P_{m-1}(\alpha)$ and

$$P_{k,m,d}(\alpha) = \Phi_d\big(P_{k+m-1}(\alpha), P_{k-1}(\alpha)\big) = P_{k-1}^{\varphi(d)}(\alpha) \cdot \Phi_d(1,1) = \nu P_{k-1}^{\varphi(d)}(\alpha).$$

It follows that

$$\begin{aligned}\text{resultant}(P_{k,m,d}, R_n) &= \prod_{\alpha \in R_n^{-1}(0)} P_{k,m,d}(\alpha) \\ &= \nu^{\deg(R_n)} \cdot \prod_{\alpha \in R_n^{-1}(0)} P_{k-1}^{\varphi(d)}(\alpha) = \nu^{\deg(R_n)} \cdot \text{resultant}(P_{k-1}^{\varphi(d)}, R_n).\end{aligned}$$

Since n does not divide $k-1$, Lemma 27 yields $\text{resultant}(R_\ell, R_n) = \pm 1$ for any divisor ℓ of $k-1$. Thus,

$$\begin{aligned}\text{resultant}(P_{k,m,d}, R_n) &= \nu^{\deg(R_n)} \cdot \text{resultant}(P_{k-1}^{\varphi(d)}, R_n) \\ &= \nu^{\deg(R_n)} \cdot \prod_{\ell \mid k-1} \big(\text{resultant}(R_\ell, R_n)\big)^{\varphi(d)} = \pm\nu^{\deg(R_n)}.\end{aligned}$$

Equation (5) now follows from Equation (2).

Case 2.b: n divides $k-1$. As in the proof of Lemma 27, if n divides ℓ, then

$$P_\ell = P_n \pmod{P_n^D} = P_n \cdot (1 + H_\ell)$$

with $H_\ell \in \mathbb{Z}[a]$ divisible by P_n. It follows that

$$P_{k,m,d} = \Phi_d\big(P_{k+m-1}, P_{k-1}\big) = P_n^{\varphi(d)} \cdot (\nu + H_{k,m,d})$$

with $H_{k,m,d} \in \mathbb{Z}[a]$ divisible by P_n. Since n divides $\gcd(m, k-1)$, Equation (2) yields

$$\left(\prod_{\substack{\ell \mid \gcd(m,k-1) \\ \ell \text{ does not divide } n}} R_\ell^{\varphi(d)}\right) \cdot \left(\prod_{\ell \mid m} R_{k,\ell,d}\right) = \nu + H_{k,m,d} P_n^{D-1}$$

and since resultant$(R_\ell, R_n) = \pm 1$ for $\ell \neq n$, we deduce that

$$\prod_{\ell|m} \text{resultant}(R_{k,\ell,d}, R_n) = \text{resultant}(\nu + H_{k,m,d} P_n^{D-1}, R_n)$$
$$= \text{resultant}(\nu, R_n) = \pm \nu^{\deg(R_n)}.$$

This is Equation (5).

The proof of Lemma 23 is completed □

Proof of Lemma 24. Assume $D = p^e$ is a prime power and $d \geq 2$ is a divisor of D. We need to show that for all $k \geq 2$, the polynomials $R_{k,1,d} \pmod p$ are powers of $a \in \mathbb{F}_p[a]$; and for all $k \geq 2$ and $n \geq 2$, the polynomials $R_{k,n,d} \pmod p$ are powers of $R_n \pmod p$. Since $R_{k,n,d}$ divides $R_{k,n}$ for all $n \geq 1$, it is enough to prove that for all $k \geq 2$, the polynomials $R_{k,1} \pmod p$ are powers of $a \in \mathbb{F}_p[a]$; and for all $k \geq 2$ and $n \geq 2$, the polynomials $R_{k,n} \pmod p$ are powers of $R_n \pmod p$.

For $k \geq 2$, set $M_{k,1} := (D-1)N_{k-1}$ and for $n \geq 2$, set

$$M_{k,n} := \begin{cases} (D-1)(D^{k-1}-1) & \text{if } n \text{ divides } k-1 \\ (D-1)D^{k-1} & \text{if } n \text{ does not divide } k-1. \end{cases}$$

We prove that for $k \geq 2$ and $n \geq 2$,

$$R_{k,1} \equiv a^{M_{k,1}} \pmod p \quad \text{and} \quad R_{k,n} \equiv R_n^{M_{k,n}} \pmod p. \tag{6}$$

Note that $N_{i+j} - N_i = D^i N_j$ for all integers $i \geq 0$ and $j \geq 0$. So, according to Lemma 26, if $k \geq 2$ and $n \geq 1$,

$$\begin{aligned} P_{k+n-1} - P_{k-1} &\equiv a^{N_{k-1}} + a^{N_k} + \cdots + a^{N_{k+n-2}} \pmod p \\ &\equiv a^{N_{k-1}} \cdot \left(a^{D^{k-1}N_0} + a^{D^{k-1}N_1} + \cdots + a^{D^{k-1}N_{n-1}} \right) \pmod p \\ &\equiv a^{N_{k-1}} \cdot \left(a^{N_0} + a^{N_1} + \cdots + a^{N_{n-1}} \right)^{D^{k-1}} \pmod p \\ &\equiv a^{N_{k-1}} P_n^{D^{k-1}} \pmod p. \end{aligned}$$

As a consequence,

$$P_{k+n-1}^D - P_{k-1}^D \equiv a^{DN_{k-1}} P_n^{D^k} \pmod p$$

and

$$P_{k,n} \equiv a^{(D-1)N_{k-1}} P_n^{D^k - D^{k-1}} \pmod p \equiv a^{M_{k,1}} P_n^{(D-1)D^{k-1}} \pmod p.$$

In particular, for $n = 1$, this yields

$$R_{k,1} = P_{k,1} \equiv a^{M_{k,1}} \pmod p.$$

According to Equation (3),

$$\left(\prod_{m|\gcd(n,k-1)} R_m^{D-1} \right) \cdot \left(\prod_{m|n} R_{k,m} \right) = P_{k,n} \equiv a^{M_{k,1}} \cdot \prod_{m|n} R_m^{(D-1)D^{k-1}} \pmod p$$

and since $R_1 = 1$ and $R_{k,1} \equiv a^{M_{k,1}} \pmod p$,

$$\prod_{\substack{m|n \\ m \neq 1}} R_{k,m} = \prod_{\substack{m|n \\ m \neq 1}} R_m^{M_{k,m}} \pmod p.$$

Equation (6) now follows from the Möbius inversion formula, completing the proof of Lemma 24. □

To complete the proof of Theorem 19, we use the following generalization of the Eisenstein criterion.

Lemma 31. *Assume $A \in \mathbb{Z}[a]$ and $B \in \mathbb{Z}[a]$ are monic polynomials and p is a prime number such that*

- *$A = B^N \pmod p$ for some integer $N \geq 1$;*
- *the polynomial $B \pmod p$ is irreducible over $\mathbb{F}_p$;*
- *$p^{2\deg(B)}$ does not divide $\operatorname{resultant}(A, B)$.*

Then, A is irreducible over $\mathbb{Q}$.

Proof. Assume by contradiction that A is reducible over $\mathbb{Q}$, so that $A = A_1A_2$ with $A_1 \in \mathbb{Z}[a]$ and $A_2 \in \mathbb{Z}[a]$ non constant. Let $\overline{A}_1$, $\overline{A}_2$ and $\overline{B}$ be the reductions of the polynomials modulo p. Then, $\overline{A}_1\overline{A}_2 = \overline{B}^N$ and since $\overline{B}$ is irreducible over $\mathbb{F}_p$, we have that $\overline{A}_1 = \overline{B}^{N_1}$ and $\overline{A}_2 = \overline{B}^{N_2}$ for some positive integers $N_1 \geq 1$ and $N_2 \geq 1$. In other words, $A_1 = B^{N_1} + pC_1$ and $A_2 = B^{N_2} + pC_2$ for some polynomials $C_1 \in \mathbb{Z}[a]$ and $C_2 \in \mathbb{Z}[a]$. In that case,

$$\begin{aligned}\operatorname{resultant}(A, B) = \operatorname{resultant}(A_1A_2, B) &= \operatorname{resultant}(A_1, B) \cdot \operatorname{resultant}(A_2, B)\\ &= \operatorname{resultant}(pC_1, B) \cdot \operatorname{resultant}(pC_2, B)\\ &= p^{2\deg(B)}\operatorname{resultant}(C_1C_2, B).\end{aligned}$$

This contradicts the assumption that $p^{2\deg(B)}$ does not divide $\operatorname{resultant}(A, B)$. □

We now complete the proof of Theorem 19. Assume $D = p^e$ is a prime power and $d \geq 2$ is a divisor of D. Then d is a power of p.

According to Lemma 24, the polynomial $R_{k,1,d} \pmod p$ is a power of $a \in \mathbb{F}_p[a]$, which is irreducible over $\mathbb{F}_p$; and according to Lemma 23, $p^{2\deg(R_n)}$ does not divide $\operatorname{resultant}(R_{k,1,d}, R_1) = \pm p^{\deg(R_n)}$. It follows from Lemma 31 that $R_{1,k,d}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$.

Similary, according to Lemma 24, if $n \geq 2$, the polynomial $R_{k,n,d} \pmod p$ is a power of $R_n \pmod p$; and according to Lemma 23, $p^{2\deg(R_n)}$ does not divide $\operatorname{resultant}(R_{k,n,d}, R_n) = \pm p^{\deg(R_n)}$. It follows from Lemma 31 that when $R_n \pmod p$ is irreducible over $\mathbb{F}_p$, the polynomial $R_{k,n,d}$ is irreducible over $\mathbb{Q}$ for all $k \geq 2$.

This completes the proof of Theorem 19.

3.4. **Particular cases.** For small values of k and n, the expression of $R_{k,n,d}$ is quite simple and we may obtain irreducibility as follows.

Proposition 32. *For all $D \geq 2$ and all d that divide D, the polynomial $R_{2,1,d}$ is irreducible over $\mathbb{Q}$.*

Proof. We have that

$$R_{2,1,d} = \Phi_d(a + 1, 1).$$

Since cyclotomic polynomials are irreducible over $\mathbb{Q}$, so is $R_{2,1,d}$. □

Proposition 33. *For all $D \geq 2$ even, the polynomial $R_{3,1,2}$ is irreducible over $\mathbb{Q}$.*

Proof. Setting $b := a + 1$, we have that

$$R_{3,1,2} = \Phi_2(P_3, P_2) = P_3 + P_2 = a(a + 1)^D + 1 + (a + 1) = b^{2d+1} - b^{2d} + b + 1.$$

By [FJ, Theorem 2], this quadrinomial is irreducible for all $d \geq 1$. □

Proposition 34. *For all $D \geq 2$ even, the polynomial $R_{2,2,2}$ is irreducible over $\mathbb{Q}$.*

Proof. Assume $D = 2d$ is even. Then setting $b = a + 1$ as previously,

$$\begin{aligned} R_{2,2,2} = \frac{\Phi_2(P_3, P_1)}{\Phi_2(P_2, P_1)} &= \frac{P_3 + P_1}{P_2 + P_1} \\ &= \frac{a(a+1)^D + 2}{a+2} \\ &= \frac{b^{2d+1} - b^{2d} + 2}{b+1} = b^{2d} - 2b^{2d-1} + 2b^{2d-2} \ - \cdots - 2b + 2. \end{aligned}$$

According to the Eisenstein criterion, this polynomial is irreducible over $\mathbb{Q}$. □

3.5. **Irreducibility over $\mathbb{F}_p$.** Here, $D = p^e$ is a prime power, and we work over the field $\mathbb{F}_p$ or its algebraic closure $\overline{\mathbb{F}}_p$. Abusing notation, we keep the notation P_n and R_n for their reductions modulo p. In other words, $P_n \in \mathbb{F}_p[a]$ and $R_n \in \mathbb{F}_p[a]$ are defined by

$$P_n := \sum_{k=0}^{n-1} a^{N_k} \quad \text{with} \quad N_k := \frac{D^k - 1}{D - 1} \quad \text{and} \quad R_n := \prod_{m|n} P_m^{\mu(n/m)}.$$

We study the irreducibility of R_n over $\mathbb{F}_p$. Note that

$$R_1 = 1 \quad \text{and} \quad R_2 = a + 1.$$

So, we restrict our study to the case $n \geq 3$.

Proposition 35. *Assume $D = p^e$ is a prime power and $n \geq 3$. Then, the polynomial $R_n \in \mathbb{F}_p[a]$ is irreducible over $\mathbb{F}_p$ if and only if either $n = 3$ and $D = 2$, or $n = 3$ and $D = 8$.*

Proof. Let $f : \overline{\mathbb{F}}_p \to \overline{\mathbb{F}}_p$ be the Frobenius automorphism $x \mapsto x^p$.

Lemma 36. *If $\alpha \in \overline{\mathbb{F}}_p$ is a root of R_n, then α is a periodic point of f of period dividing $n \cdot e$.*

Proof. Assume α is a root of R_n. Then, $P_n(\alpha) = 0$, so that

$$\begin{aligned} 1 = 1 + \alpha P_n^D(\alpha) &= 1 + \alpha P_n(\alpha^D) \\ &= 1 + \sum_{k=0}^{n-1} \alpha^{1+DN_k} \\ &= 1 + \sum_{k=0}^{n-1} \alpha^{N_{k+1}} = P_n(\alpha) + \alpha^{N_n} = \alpha^{N_n}. \end{aligned}$$

It follows that

$$f^{\circ(n\cdot e)}(\alpha) = \alpha^{D^n} = \alpha^{1+(D-1)N_n} = \alpha \cdot \left(\alpha^{N_n}\right)^{D-1} = \alpha. \qquad \square$$

As a consequence, if R_n is irreducible over $\mathbb{F}_p$, then the degree of R_n divides $n \cdot e$. The degree of R_n is

$$\deg(R_n) = \sum_{m|n} \mu\left(\frac{n}{m}\right) \deg(P_m) = \sum_{m|n} \mu\left(\frac{n}{m}\right) N_{m-1} \geq D^{n-2}.$$

So, if R_n is irreducible over $\mathbb{F}_p$, then $p^{(n-2)e} \leq n \cdot e$.

Set $\kappa := (n-2)\log(p) > 0$. The function $(0,+\infty) \ni x \mapsto \exp(\kappa x)/x \in (0,+\infty)$ reaches a minimum at $x = 1/\kappa$ with value $\kappa \cdot \exp(1)$. It follows that for $n \geq 3$,

$$\frac{p^{(n-2)e}}{n \cdot e} \geq \left(1 - \frac{2}{n}\right) \log(p) \exp(1).$$

If $n \geq 3$ and $p \geq 5$, or if $n \geq 4$ and $p = 3$, or if $n \geq 5$ and $p = 2$, this is greater than 1. So, it is enough to study the following cases.

Case $n = 3$ and $p = 2$. In that case, for $e \geq 1$,

$$\deg(R_n) = 1 + D = 2^e + 1 \quad \text{and} \quad n \cdot e = 3e.$$

The function $(0,+\infty) \ni x \mapsto (2^x + 1)/(3x) \in (0,+\infty)$ is increasing on $[2,+\infty)$ and takes the values 1 at $x = 1$, $5/6$ at $x = 2$ and 1 at $x = 3$. It follows that $\deg(R_n)$ divides $n \cdot e$ if and only if $e = 1$ or $e = 3$, i.e. $D = 2$ or $D = 8$; in those two cases, R_3 is irreducible.

Case $n = 3$ and $p = 3$. In that case, for $e \geq 1$,

$$\deg(R_n) = 1 + D = 3^e + 1 > 3e = n \cdot e = 3e.$$

So, R_n cannot be irreducible in that case.

Case $n = 4$ and $p = 2$. In that case, for $e \geq 1$,

$$\deg(R_n) = 1 + D + D^2 = 1 + 3^e + 3^{2e} > 4e = n \cdot e.$$

So, R_n cannot be irreducible in that case. □

References

[AK] M. Arfeux & J. Kiwi *Irreducibility of the set of cubic polynomials with one periodic critical point*, Preprint, `https://arxiv.org/abs/1611.09281`

[Bo] T. Bousch *Sur quelques problèmes de dynamique holomorphe*, Ph.D. thesis, Université de Paris- Sud, Orsay, 1992.

[Bu] X. Buff *On Postcritically Finite Unicritical Polynomials*, Preprint, `https://www.math.univ-toulouse.fr/~buff/Preprints/Gleason/Gleason.pdf`.

[E] A. L. Epstein *Integrality and rigidity for postcritically finite polynomials*, Bull. London Math. Soc. 44 (2012), 39–46.

[FJ] C. Finch & L. Jones *On the irreducibility of $\{-1,0,1\}$-quadrinomials*, Integers 6 (2006).

[G] V. Goksel *On the orbit of a post-critically finite polynomial of the form $x^d + c$*, Preprint, `https://arxiv.org/abs/1806.01208`.

[HT] B. Hutz & A. Towsley *Misiurewicz points for polynomial maps and transversality*, New York J. Math. 21 (2015) 297–319.

[M1] J. Milnor *Geometry and dynamics of quadratic rational maps*, Experiment. Math. Volume 2, Issue 1 1993, 37–83.

[M2] J. Milnor *Cubic polynomials with periodic critical orbit, Part I*, In "Complex Dynamics Families and Friends", ed. D. Schleicher, A. K. Peters 2009, pp 333–411.

[M3] J. Milnor *Arithmetic of unicritical polynomial maps*, Frontiers in Complex Dynamics: In Celebration of John Milnor's 80th Birthday 2012, 15–23.

[R] M. Rees *View of Parameter Space: Topographer and Resident*, Astérisque 288, 2003.

E-mail address: `xavier.buff@math.univ-toulouse.fr`

Institut de Mathématiques de Toulouse, UMR5219, Université de Toulouse, CNRS, UPS, F-31062 Toulouse Cedex 9, France

E-mail address: `adame@maths.warwick.ac.uk`

Mathematics institute, University of Warwick, Coventry CV4 7AL, United Kingdom

E-mail address: `kochsc@umich.edu`

Department of Mathematics, 530 Church Street, East Hall, University of Michigan, Ann Arbor MI 48109, United States