## Warm-up questions

(These warm-up questions are optional, and won't be graded.)

1. Consider  $\mathbb{R}$  with the Euclidean metric. Which of the following maps  $f:\mathbb{R}\to\mathbb{R}$  are homeomorphisms (Assignment Question 2)?

- (a) f(x) = ax + b (b)  $f(x) = x^2$
- (c)  $f(x) = x^3$
- (d)  $f(x) = \sin(x)$

2. Let  $f: X \to Y$  be an **invertible** function of sets, and let  $S \subseteq Y$ . The notation  $f^{-1}(S)$  could denote either the preimage of S under f, or the image of S under the inverse function  $f^{-1}$ . Show that these two sets are equal, so there is no ambiguity in using the notation  $f^{-1}(S)$ .

3. Let  $f: X \to Y$  be an **invertible** function of sets.

- (a) Show that, for subsets  $B \subseteq Y$ , there is equality  $f(f^{-1}(B)) = B$ .
- (b) Show that, for subsets  $A \subseteq X$ , there is equality  $f^{-1}(f(A)) = A$ .

4. See the definition of bounded in Assignment Question 3.

- (a) Negate the definition of bounded to state what it means for a subset S of a metric space to be unbounded.
- (b) Is  $\emptyset$  a bounded set?
- (c) Show that any **finite** subset of a metric space is bounded.

5. Give examples of subsets of  $\mathbb{R}$  (with the Euclidean metric) that satisfy the following.

(a) open, and bounded

(c) open, and unbounded

(b) closed, and bounded

(d) closed, and unbounded

6. Consider  $\mathbb{R}$  with the Euclidean metric. For each of the following sets A, find Int(A),  $\overline{A}$ ,  $\partial A$ ,  $\operatorname{Int}(\mathbb{R}\setminus A), \mathbb{R}\setminus A$ , and  $\partial(\mathbb{R}\setminus A)$ . See Assignment Question 4 for the definition of a boundary  $\partial$ .

- (a)  $\mathbb{R}$
- (c) (0,1)
- (e)  $\{0,1\}$

- (b) [0,1]
- (d) (0,1]
- $(f) \{ \frac{1}{n} \mid n \in \mathbb{N} \}$

7. Consider the real numbers  $\mathbb{R}$  with the Euclidean metric. Find examples of subsets A of  $\mathbb{R}$  with the following properties.

- (a)  $\partial(A) = \emptyset$
- (b) A has a nonempty boundary, and A contains its boundary  $\partial A$ .
- (c) A has a nonempty boundary, and A contains no points in its boundary
- (d) A has a nonempty boundary, and A contains some but not all of the points in its boundary.
- (e) A has a nonempty boundary, and  $A = \partial A$ .
- (f) A is a **proper** subset of  $\partial A$ .

8. Let X be a nonempty set with the discrete metric. Let  $A \subseteq X$ . Show that  $A = \operatorname{Int}(A) = \overline{A}$ . Conclude that  $\partial A = \emptyset$ .

## Worksheet problems

(Hand these questions in!)

• Worksheet #5 Problems 1(a), 2, 3.

## Assignment questions

(Hand these questions in!)

- 1. Let  $f: X \to Y$  be a function of sets X and Y. Let  $A, B \subseteq X$ . For each of the following, determine whether you can replace the symbol  $\square$  with  $\subseteq, \supseteq, =$ , or none of the above. Justify your answer by giving a proof of any set-containment or set-equality you claim. If set-equality does not hold in general, give a counterexample.
  - (a)  $f(A \cap B) \square f(A) \cap f(B)$
- (b)  $f(A \cup B) \quad \Box \quad f(A) \cup f(B)$
- (c) For  $A \subseteq B$ ,  $f(B \setminus A) \square f(B) \setminus f(A)$
- 2. Consider the following definition.

**Definition (Homeomorphism.)** Let X and Y be metric spaces. Then a map  $f: X \to Y$  is a homeomorphism if

- f is continuous;
- f has an inverse  $f^{-1}$ ;
- $f^{-1}$  is continuous.

The metric space X is called *homeomorphic* to Y if there exists a homeomorphism  $f: X \to Y$ .

- (a) Show that, if  $f: X \to Y$  is a homeomorphism, then  $f^{-1}: Y \to X$  is a homeomorphism. Conclude that X is homeomorphic to Y if and only if Y is homeomorphic to X. (We simply call the two spaces homeomorphic.)
- (b) Let  $f: X \to Y$  be a homeomorphism of metric spaces. Show that

$$\{U \subseteq Y \mid U \text{ is open}\} \longrightarrow \{V \subseteq X \mid V \text{ is open}\}$$
  
$$U \longmapsto f^{-1}(U)$$

defines a bijection between the collection of all open subsets of Y, and the collection of all open subsets of X.

*Hint:* To show it is a bijection, you could check that the assignment  $V \mapsto f(V)$  is its inverse.

(c) Consider the function

$$f: (\mathbb{R}, \text{ discrete metric}) \to (\mathbb{R}, \text{ Euclidean metric})$$
  
$$f(x) = x$$

Prove that f is continuous and invertible, but its inverse  $f^{-1}$  is not continuous.

*Remark:* Note the contrast to other mathematical fields, such as linear algebra: if a linear map has an inverse, then the inverse is automatically linear. This exercise shows that this is not true for continuous maps!

3. Consider the following definition.

**Definition (Bounded subset.)** Let (X,d) be a metric space. A subset  $S \subseteq X$  is called *bounded* if there is some  $x_0 \in X$  and some  $R \in \mathbb{R}$  with R > 0 such that  $S \subseteq B_R(x_0)$ .

- (a) Let (X, d) be a metric space, and  $A \subseteq X$  a subset. Show that A is bounded if and only if, for every  $x \in A$ , there is some  $R_x > 0$  such that  $A \subseteq B_{R_x}(x)$ .
- (b) Let (X, d) be a metric space. Suppose that  $(a_n)_{n \in \mathbb{N}}$  is a sequence in X converging to an element  $a_{\infty}$ . Show that the set  $\{a_n \mid n \in \mathbb{N}\}$  is a bounded subset of X.
- 4. Consider the following definition.

**Definition (Boundary of a set** A.) Let (X, d) be a metric space, and let  $A \subseteq X$ . Then the boundary of A, denoted  $\partial A$ , is the set  $\overline{A} \setminus \operatorname{Int}(A)$ .

Let (X, d) be a metric space, and let  $A \subseteq X$ .

- (a) Prove that  $Int(A) = \overline{A} \setminus \partial A$ .
- (b) Prove that  $\partial A = \overline{A} \cap (\overline{X \setminus A})$ .
- (c) Conclude from part (b) that  $\partial A$  is closed.
- (d) Additionally conclude from part (b) that  $\partial A = \partial (X \setminus A)$ .
- (e) Prove the following characterization of points in the boundary:

Theorem (An equivalent definition of  $\partial A$ ). Let (X, d) be a metric space, and let  $A \subseteq X$ . Then  $x \in \partial A$  if and only if every ball  $B_r(x)$  about x contains at least one point of A, and at least one point of  $X \setminus A$ .

- (f) Deduce that we can classify every point of X in one of three mutually exclusive categories:
  - (i) interior points of A;
  - (ii) interior points of  $X \setminus A$ ;
  - (iii) points in the (common) boundary of A and  $X \setminus A$ .