

Daniel Vitek

The following is a brief summary of my mathematical history, starting with competitions, and continuing on to research.

I participated in high school Olympiads, taking the USAMO three times and finishing in the top 50 nationwide in my senior year. At Duke, I finished in the top 200 on the Putnam my freshman year, and took 26th place overall my sophomore year. I won the fall 2012 Virginia Tech regional competition. These experiences also got me a job at Awesome Math Summer Program, where I worked during summers 2009-2011 as a teaching assistant and, in summer 2012, as a full instructor in algebra and number theory.

My first taste of mathematical research came when I spent summer 2008 at the Research Science Institute at MIT working on a problem in algebraic graph theory, and presented my work at several high school science fairs. However, my most significant research has taken place at Duke. Since spring semester of 2012 I have been working in algebraic knot theory with Professor Christopher Cornwell. I have been studying an construction called knot contact homology, which is a tensor-algebra-valued invariant of knots. From this tensor algebra it is common to define a two-variable polynomial called the augmentation polynomial; as the name suggests, this polynomial measures when there exists algebra maps from knot contact homology into the complex numbers. I obtained a complete characterization of the augmentation polynomials of $(2, k)$ -torus knots, which was the first calculation of the augmentation polynomial of an infinite family of knots. This calculation was achieved via elimination theory; I analyzed the form of the determinant of the Sylvester matrix of two relations obtained from knot contact homology, and obtained a bound on the number of augmentations of certain forms in the two variables; previous knowledge of augmentations of $(2, k)$ -torus knots established the form.

Currently I'm trying to understand how augmentations of a knot's contact homology arise as images of representations $\rho : \pi_1(\mathbb{R}^3 - K) \rightarrow GL_k \mathbb{C}$ where the peripheral system $\langle l, m \rangle$ of the knot are sent to diagonal matrices of a certain form. In particular the existence or non-existence of such representations in certain dimensions is closely tied to certain classical knot invariants such as meridional rank or bridge number; I'm trying to improve the known bounds and establish the reverse directions of known relations.

In terms of larger-scale interests, I'm most fascinated by arithmetic topology. This project attempts to study the analogies one gets by trying to look at $Spec(\mathbb{Z})$ as a 3-manifold, as is suggested by Artin-Verdier duality. In particular there is a large dictionary (called the Mazur-Kapranov-Reznikov, or MKR, dictionary) that provides an analogy between knots and links in 3-manifolds and ideals in the rings of algebraic integers of a number field; for example, under this dictionary a peripheral system of a knot corresponds to a Frobenius automorphism and a monodromy generating the tame inertial group. This is personally interesting because, while I've gotten into knot theory mostly in college, number theory was easily my favorite and best subject in high school competitions, and the idea of a high-level link between the two is motivating as a higher-level, and perhaps more fruitful, way to view knot theory.