

Useful constants and equations

$$\begin{aligned}
 e &= 1.602 \times 10^{-19} \text{ C} & \frac{1}{4\pi\epsilon_0} &= 9.0 \times 10^9 \text{ N m}^2/\text{C}^2 & \frac{e^2}{4\pi\epsilon_0} &= 1.44 \text{ eV nm} \\
 c &= 3.00 \times 10^8 \text{ m/s} & h &= 6.626 \times 10^{-34} \text{ J s} = 4.136 \times 10^{-15} \text{ eV s} & \hbar &= \frac{h}{2\pi} \\
 hc &= 1240 \text{ eV nm} & \hbar c &= 197.3 \text{ MeV fm} & 1 \text{ eV} &= 1.602 \times 10^{-19} \text{ J} & R_\infty &= 1.097 \times 10^7 \text{ m}^{-1} \\
 m_e &= 9.11 \times 10^{-31} \text{ kg} = 511 \text{ keV}/c^2 = 5.486 \times 10^{-4} \text{ u} & \text{neutral } {}^{12}_6\text{C atom mass} &= 12.0000 \text{ u} \\
 m_p &= 1.673 \times 10^{-27} \text{ kg} = 938.3 \text{ MeV}/c^2 = 1.0073 \text{ u} & 1 \text{ u} &= 931.5 \text{ MeV} \\
 m_n &= 1.675 \times 10^{-27} \text{ kg} = 939.6 \text{ MeV}/c^2 = 1.0087 \text{ u} \\
 a_0 &= 0.0529 \text{ nm} & E_0 &= -13.6 \text{ eV} & \alpha &= 1/137 & \sigma &= 5.670 \times 10^{-8} \text{ W/m}^2\text{K}^4
 \end{aligned}$$

Heisenberg:

$$\Delta p_x \Delta x \sim \hbar \quad \Delta E \Delta t \sim \hbar$$

Atomic Physics:

$$\text{Hydrogen Atom: } E = \frac{-13.6 \text{ eV}}{n^2} \quad \text{with } n = 1, 2, 3, \dots \quad l < n \quad -l \leq m \leq +l$$

$$\text{Generalized Balmer Formula: } \frac{1}{\lambda} = R_\infty \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$\begin{aligned}
 \text{Angular Momentum: } \quad L^2 &= l(l+1) \hbar^2 & l &= 0, 1, 2, 3, \dots \quad (\text{orbital}) \\
 L_z &= |\vec{L}| \cos \theta = m \hbar & -l &\leq m_l \leq l \quad (\text{integer steps})
 \end{aligned}$$

$$\begin{aligned}
 \text{Magnetic Moment: } \quad \text{orbital: } \vec{\mu} &= \frac{q}{2M} \vec{L} \\
 \text{for electron: } \quad |\mu_z| &= 2 \cdot |m_s| \frac{e\hbar}{2m_e} = \mu_B = 5.8 \times 10^{-5} \text{ eV/T} \\
 \text{for proton: } \quad |\mu_z| &= g_p \cdot |m_s| \frac{e\hbar}{2m_p} = 5.6 \frac{1}{2} \mu_N = 8.8 \times 10^{-8} \text{ eV/T}
 \end{aligned}$$

$$\text{Energy of magnetic dipole in B field: } E = -\mu \cdot B$$

$$\text{Energy of particle in 3-dim } \infty \text{ square well: } E_{n_x n_y n_z} = \frac{(n_x^2 + n_y^2 + n_z^2) \hbar^2}{8mL^2}$$

$$\text{Molecular excitations: } \quad \text{vibrational: } E = \left(n + \frac{1}{2}\right) \hbar\omega \quad \text{rotational: } E = \frac{L^2}{2I} = \frac{l(l+1)}{2I} \hbar^2$$

Statistical Physics:

Maxwell Boltzmann distribution: $f_{MB} = A e^{-E/kT}$

Bose-Einstein distribution: $f_{BE} = \frac{1}{B e^{E/kT} - 1}$

Fermi - Dirac distribution: $f_{FD} = \frac{1}{e^{(E-E_F)/kT} + 1}$

Boltzmann constant $k = 1.381 \times 10^{-23} \text{ J/K} = 8.617 \times 10^{-5} \text{ eV/K}$

For "gas" of free fermions: $g(E) = \frac{8\sqrt{2}\pi m^{3/2}}{h^3} \sqrt{E}$

$$E_F = \frac{h^2}{2m} \left(\frac{3N}{8\pi V} \right)^{2/3} \quad E_m = \frac{3}{5} E_F$$

Nuclear Physics:

nuclear radius: $R = (1.2 \times 10^{-15} \text{ m}) A^{1/3}$ average binding energy / nucleon $\approx 8 \text{ MeV}$

range of interaction: $R = \frac{\hbar}{mc} = \frac{\hbar c}{mc^2}$

Decay law: $N = N_0 e^{-\frac{t}{\tau}}$ with $\langle t \rangle = \tau = \frac{1}{\lambda}$ and $T_{1/2} = \frac{\ln 2}{\lambda}$

Binding energy $B(Z,A) = [Z m_p + N m_n - M_{atom}(Z,A)] c^2$

Particle Physics:

Baryon: $Q Q Q$ Meson: $Q \bar{Q}$

Quarks: $\begin{pmatrix} u \\ d \end{pmatrix} \begin{pmatrix} c \\ s \end{pmatrix} \begin{pmatrix} t \\ b \end{pmatrix} \begin{matrix} q = +\frac{2}{3} \\ q = -\frac{1}{3} \end{matrix}$ Leptons: $\begin{pmatrix} e \\ \nu_e \end{pmatrix} \begin{pmatrix} \mu \\ \nu_\mu \end{pmatrix} \begin{pmatrix} \tau \\ \nu_\tau \end{pmatrix} \begin{matrix} q = -1 \\ q = 0 \end{matrix}$

Cosmology:

Luminosity of star: $L = 4\pi r^2 f$ with $f =$ apparent brightness of star

difference in apparent magnitude: $m_1 - m_2 = 2.5 \log(f_1/f_2)$