


Douglass Houghton Workshop, Section 1, Mon 11/26/18
Worksheet May the Road Rise to Meet You


1. We've been working on the problem of finding the shortest road network between three cities in the plane.

In the case we considered, the three cities were at the corners of a 45° - 45° - 90° triangle with legs 50 miles long. The simplest idea is to just build roads along the legs; that makes a network of length 100. But by constructing a Λ -shaped network like the one at the right, we found




- The length of the network is $x + 2\sqrt{2500 - 100x \cos(45) + x^2}$.
- We can improve from the simple 2-road solution ($x = 0$, length = 100) by increasing x . For instance, when $x = 10$, the network has a length of about 97.

- (a) Consider the case where the triangle is still isosceles and the legs still have length 50, but the angle at B is 70° . Write a formula for the length of the network.



- (b) Can you find a value of x which beats the 2-road solution ($x = 0$, length = 100)?

- (c) Now suppose the vertex angle is very obtuse—say 150° . Find a formula for the length of the network.



- (d) Can you beat the 2-road solution in this case?

- (e) Suppose the vertex angle is θ . Write a formula for the length of the network.

2. Write the following sums in sigma (Σ) notation.


- (a) $1 + 2 + 3 + 4 + \cdots + 10$
- (b) $1 + 2 + 3 + 4 + \cdots + n$
- (c) $3 + 5 + 7 + 9 + \cdots + 21$
- (d) $4 + 9 + 16 + 25 + \cdots + 100$
- (e) $2.3 + 2.8 + 3.3 + 3.8 + 4.3 + 4.8 + \cdots + 10.3$
- (f) $f(a_1) + f(a_2) + f(a_3) + \cdots + f(a_n)$

3. Consider the function $f(x) = x^x$.

- (a) It's neither a power function (ax^b) nor an exponential (ab^x). Nevertheless, find its derivative. Hint: rewrite it in the form $e^{u(x)}$ for some function u .
- (b) What is the minimum value that f takes on? (Check with your calculator, but find the answer with calculus.)

4. (This problem appeared on a Winter, 2008 Math 115 exam) A bellows has a triangular frame made of three rigid pieces. Two pieces, each 10 inches long, are hinged at the nozzle. They are attached to the third piece at points A and B which can slide, as shown in the diagrams below. (The figures show a 3D sketch of the bellows and a 2D sketch that may be specifically useful to solve the problem.)


Each piece of the frame is 2 inches wide, so the volume (in cubic inches) of air inside the bellows is equal to the area (in square inches) of the triangular cross-section above, times 2. Suppose you pump the bellows by moving A downward toward the center at a constant speed of 3 in/sec. (So B moves upwards at the same speed.) What is the rate at which air is being pumped out when A and B are 12 inches apart? (So A is 6 inches from the center of the vertical piece of the frame.)



5. The table below gives the expected growth rate, $g(t)$, in ounces per week, of the weight of a baby in its first 54 weeks of life (which is slightly more than a year). Assume for this problem that $g(t)$ is a decreasing function.

week t	0	9	18	27	36	45	54
growth rate $g(t)$	6	6	4.5	3	3	3	2

- (a) Using six subdivisions, find an overestimate and underestimate for the total weight gained by a baby over its first 54 weeks of life.
- (b) How often would you have to weigh the baby to get an estimate guaranteed to be accurate to within $\frac{1}{4}$ pound?



6. Kaleb kicks a soccer ball toward the goal. The ball passes through various air currents along the way, and its horizontal velocity, in miles per hour, is given by the graph to the left. Estimate the distance the ball travels in the first 1.2 seconds.