

### 395 DISCUSSION SECTION 3

WITH NOAH LUNTZLARA

We start with the problems we didn't get to last time:

Let  $G$  be a group, and  $S \subseteq G$  be an arbitrary subset of  $G$ . Recall the definitions of the (*left* and *right*) *cosets* of  $S$  in  $G$ : for each  $g \in G$ , we define

$$gS := \{gx \mid x \in S\}; \quad \text{respectively,} \quad Sg := \{xg \mid x \in S\}.$$

- (Ex. 1) When are right cosets and left cosets the same? When are they different? Give some examples.
- (Ex. 2) Let  $G$  be a finite group and  $H \subseteq G$  be a subgroup of  $G$
- (a) Show that  $g_1H = g_2H$  iff  $g_1g_2^{-1} \in H$
  - (b) Show that the left cosets of  $H$  partition  $G$ ; i.e., the relation  $\sim$  defined by  $g_1 \sim g_2$  provided that  $g_1H = g_2H$  is an equivalence relation.
  - (c) Show that for any  $g_1, g_2 \in G$  there is a bijection  $g_1H \rightarrow g_2H$ , and hence all left cosets of  $H$  in  $G$  have the same cardinality.
  - (d) The number of left cosets of  $H$  in  $G$  is called the *index* of  $H$  in  $G$ , denoted  $[G : H]$ . Prove *Lagrange's Theorem*:  $|G| = |H| \cdot [G : H]$ .
  - (e) Explain why the order of a subgroup of a finite group must divide the order of the group (this is also called Lagrange's Theorem).
- (Ex. 3) Show that when we do everything above with right cosets instead of left cosets, we get the same number for  $[G : H]$ .
- (Ex. 4) Classify all groups with no proper nontrivial subgroups (i.e., subgroups that are neither the group itself nor the trivial subgroup  $\{e\}$ ).

A subgroup  $H$  of a group  $G$  is called *normal* provided that  $gHg^{-1} = H$  for every  $g \in G$  (where  $gHg^{-1} = \{ghg^{-1} \mid h \in H\}$ ). If  $H$  is a normal subgroup of  $G$ , we write  $H \trianglelefteq G$ .

- (Ex. 5) Show that  $H \trianglelefteq G$  iff  $gH = Hg$  for every  $g \in G$ .
- (Ex. 6)
- (a) Show that for every group  $G$ ,  $G \trianglelefteq G$  and  $\{e\} \trianglelefteq G$ .
  - (b) Exhibit three groups with non-normal subgroups.
  - (c) Exhibit three groups with proper nontrivial normal subgroups.
- (Ex. 7) Show that if  $H \trianglelefteq G$  and  $K$  is a subgroup of  $G$  which contains  $H$ , then  $H \trianglelefteq K$ .
- (Ex. 8) *Bonus*: Try and fail to prove  $H \trianglelefteq K \trianglelefteq G \Rightarrow H \trianglelefteq G$ . Then find a group  $G$  with subgroups  $H$  and  $K$  such that  $K \trianglelefteq G$  and  $H \trianglelefteq K$ , but  $H$  is not normal in  $G$ .
- (Ex. 9) Let  $G$  be a group and  $H$  be an index-2 subgroup, so that  $[G : H] = 2$ . Show that  $H \trianglelefteq G$ .
- (Ex. 10) Recall that if  $\phi : G \rightarrow H$  is a group homomorphism,  $\ker(\phi)$  denotes  $\phi^{-1}(e_H)$ .
- (a) Show that  $\ker(\phi) \trianglelefteq G$ .
  - (b) State and prove a converse for the previous statement.