PROBLEM SET 10 (DUE ON THURSDAY, NOV 19)

(All Exercises are references to the November 18, 2017 version of Foundations of Algebraic *Geometry* by R. Vakil.)

- **Problem 1.** Exercise 10.1.M (sections of morphisms you may want to read section 8.1.2) on locally closed embeddings)
- Problem 2. Exercise 10.2.B (when are morphisms determined by where they send closed points?)
- Problem 3. Exercise 10.2.E (graphs of rational maps)
- Describe the graph of the rational map $\mathbb{A}^2_{\mathbb{C}} \dashrightarrow \mathbb{A}^1_{\mathbb{C}}$ given by x/y. Can you Problem 4. tell from this graph that this rational map cannot extend to a morphism $\mathbb{A}^2_{\mathbb{C}} \to \mathbb{P}^1_{\mathbb{C}}$? Let $n \ge 2$ be an integer. Compute the (maximal) domain of definition of the
- Problem 5. generalized Cremona transformation

$$C: \mathbb{P}^n_{\mathbb{C}} \dashrightarrow \mathbb{P}^n_{\mathbb{C}},$$

a rational map given by $[x_0 : \cdots : x_n] \mapsto [x_0^{-1} : \cdots : x_n^{-1}]$ (on closed points with $x_0 \cdots x_n \neq 0$).