

## PROBLEM SET 4 (DUE ON THURSDAY, APRIL 4)

(All Exercises are references to the November 18, 2017 version of *Foundations of Algebraic Geometry* by R. Vakil.)

- Problem 1.** Exercise 18.4.K (ample line bundles on projective curves have positive degree)
- Problem 2.** Exercise 18.6.K (Bezout's Theorem - if you want, you can just consider the case where  $X$  is reduced)
- Problem 3.** Suppose  $Z_1, Z_2$  are closed subschemes of a projective  $k$ -scheme  $X$ . Show that
$$\chi(Z_1 \cup Z_2, \mathcal{O}_{Z_1 \cup Z_2}) = \chi(Z_1, \mathcal{O}_{Z_1}) + \chi(Z_2, \mathcal{O}_{Z_2}) - \chi(Z_1 \cap Z_2, \mathcal{O}_{Z_1 \cap Z_2}),$$
where  $Z_1 \cup Z_2$  and  $Z_1 \cap Z_2$  are the scheme-theoretic union and intersection of closed subschemes. (In fact, Hilbert polynomials computed with respect to an ample line bundle on the ambient scheme  $X$  will satisfy the same identity - compare with Exercise 18.6.P.)
- Problem 4.** Exercise 18.6.R (arithmetic genus of complete intersection of surfaces in  $\mathbb{P}^3$  - in this context complete intersection just means that the surfaces being intersected do not share an irreducible component)
- Problem 5.** Let  $m, n$  be positive integers and let  $f \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(m, n))$  be a bihomogeneous polynomial of degree  $(m, n)$ . Compute the arithmetic genus of the curve  $X = V(f)$ . Your answer for  $(m, n) = (d, d)$  should agree with your answer to the previous problem when  $(m, n) = (2, d)$  - why is this? (Hint: use the ideal sheaf sequence for  $X$  inside  $\mathbb{P}^1 \times \mathbb{P}^1$ . Also, the notation  $\mathcal{O}(m, n)$  is defined in section 16.4.8.)
- Problem 6.** Show that for  $n \geq 3$ , the intersection of any two hypersurfaces in  $\mathbb{P}_k^3$  is connected. (Hint: if  $X$  is the intersection, compute  $h^0(X, \mathcal{O}_X) = 1$  using two long exact sequences of cohomology groups).