

DIEUDONNÉ THEORY VIA COHOMOLOGY OF CLASSIFYING STACKS

SHUBHODIP MONDAL

Abstract

We prove that if G is a finite flat group scheme of p -power rank over a perfect field of characteristic p , then the second crystalline cohomology of its classifying stack $H_{crys}^2(BG)$ recovers the Dieudonné module of G . We also provide a calculation of crystalline cohomology of the classifying stack of an abelian variety. We use this to prove that crystalline cohomology of the classifying stack of a p -divisible group is a symmetric algebra (in degree 2) on its Dieudonné module. We also prove mixed characteristic analogues of some of these results using prismatic cohomology.

Contents

1	Introduction	2
2	Crystalline cohomology	4
2.1	Crystalline cohomology for stacks	4
2.2	The crystalline site	4
3	Application to crystalline Dieudonné theory	7
3.1	Dieudonné module of finite group schemes	8
3.2	Dieudonné theory of p -divisible groups	11
3.3	Cohomology of the classifying stack of an abelian variety	11
3.4	Cohomology of the classifying stack a p -divisible group	13
4	Prismatic cohomology	16
4.1	Prismatic cohomology for stacks	16
4.2	Application to prismatic Dieudonné theory	18

1 Introduction

Let p be a prime which will be fixed throughout. Our beginning point is the classification of finite commutative group schemes of p -power rank over a perfect field k of characteristic p in terms of contravariant Dieudonné theory. This is achieved by defining an algebraic invariant associated to any given finite group scheme G (assumed to be commutative and p power rank unless otherwise mentioned) which is called the Dieudonné module of G and will be denoted by $M(G)$. We will recall the classical construction of $M(G)$ in Section 3. The basic object of interest is the Dieudonné ring \mathcal{D}_k which is an associative algebra over the ring of Witt vectors $W(k)$. The Dieudonné functor M which takes G to $M(G)$ transforms the study of finite group schemes over k to the study of semi-linear algebraic objects. More precisely,

Theorem 1.1 (Dieudonné). *The Dieudonné functor M induces an anti-equivalence between the category of finite commutative group schemes over k of p -power rank and left \mathcal{D}_k -modules with finite $W(k)$ length.*

Our primary goal in this paper is to provide a geometric construction of $M(G)$. In particular, we attempt to understand if $M(G)$ can be constructed in terms of some *cohomology* theory associated to some “space.” Since $M(G)$ is a module over $W(k)$, it could be natural to take the naive candidate, which is the i -th crystalline cohomology of G for some i denoted as $H_{crys}^i(G)$. However, one can see that this cannot not work for any i simply by taking discrete group schemes as input for $M(\cdot)$. Moreover, any reasonable cohomology theory will suffer from exactly the same problems. The issue is that a cohomology theory will view any discrete group scheme just as a discrete space and will be unable to detect its group structure. This suggests that we look for other spaces naturally associated to G instead of just G itself. In topology, one can get around this issue by considering the classifying space BG as explained below in Example 1.3. In category theory, the analogue of this would be to simply view a group G as a groupoid with one object and morphism set isomorphic to G . In algebraic geometry, for a group scheme G , a model of these constructions would be to consider the classifying *stack* BG . Our main goal is to prove the following

Theorem 1.2. *For a finite commutative group scheme G over k which is of p -power rank, the Dieudonné module $M(G)$ is naturally isomorphic (upto a Frobenius twist) to $H_{crys}^2(BG)$, where BG denotes the classifying stack of G . (cf. Section 3.1)*

Example 1.3. We explain the topological analogue of Theorem 1.1 and Theorem 1.2. In the case where G is a finite abelian group, one can consider the classifying space BG of G which has the property that $\pi_1(BG) \simeq G$ and $\pi_i(BG) = 0$ for $i > 1$. In this case, $H_{singular}^2(BG) \simeq \text{Ext}^1(G, \mathbb{Z})$, which is non-canonically isomorphic to the group G . By using the short exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$ we obtain that $\text{Ext}^1(G, \mathbb{Z}) \simeq \text{Hom}(G, \mathbb{Q}/\mathbb{Z}) \simeq \varinjlim \text{Hom}(G, \mathbb{Z}/n\mathbb{Z}) \simeq \text{Hom}(G, \mathbb{S}^1)$. By Pontryagin duality, one obtains that sending $G \rightarrow H_{singular}^2(BG)$ gives an anti-equivalence from the category of finite abelian groups to itself.

We also define the classifying stack BG of a p -divisible group G and prove the following

Theorem 1.4. *Let G be a p -divisible group over k . Then $H_{crys}^*(BG) \simeq \text{Sym}^*(M(G)[-2])$, where $M(G)$ denotes the Dieudonné module of G . Here $M(G)$ is considered to be in degree 2. In particular, $H_{crys}^2(BG)$ recovers the Dieudonné module $M(G)$ and $H_{crys}^i(BG) = 0$ for odd i . (cf. Section 3.4)*

In [SW14], Scholze and Weinstein defined a mixed characteristic analogue of (covariant) Dieudonné modules for p -divisible groups over a perfectoid ring. More recently, in [ALB19], Anschütz and Le Bras defined a mixed characteristic analogue of contravariant Dieudonné modules over more general base rings. In Section 4.2 we briefly recall their work and the definition of the filtered prismatic Dieudonné module $\underline{M}_\Delta(G) = (M_\Delta(G), \text{Fil } M_\Delta(G), \varphi_{M_\Delta(G)})$ associated to a p -divisible group G as defined by them. Their main theorem is the following classification result.

Theorem 1.5 ([ALB19]). *Let R be a quasi-regular semiperfectoid ring. The filtered prismatic Dieudonné module functor*

$$G \rightarrow \underline{M}_\Delta(G)$$

defines an anti-equivalence between the category p -divisible groups over R and the category of filtered prismatic Dieudonné modules over R .

Using the classifying stack BG of a p -divisible group G we prove the following

Theorem 1.6. *Let G be a p -divisible group over a quasi-regular semiperfectoid ring R . Then the prismatic cohomology $H_{\Delta}^2(BG)$ is naturally isomorphic to the prismatic Dieudonné module $M_{\Delta}(G)$. This isomorphism identifies the natural Frobenius on $H_{\Delta}^2(BG)$ with the endomorphism $\varphi_{M_{\Delta}(G)}$ on $M_{\Delta}(G)$. Further, the Nygaard filtration $\mathcal{N}^{\geq 1}H_{\Delta}^2(BG)$ on prismatic cohomology of the stack BG is isomorphic to $\text{Fil } M_{\Delta}(G)$. (cf. Section 4.2)*

On the other hand, there has been some recent work involving computation of p -adic cohomology theories associated to classifying stack of group schemes over p -adic base rings. For example, we refer to the work of Totaro [Tot18] and the work of Antieau, Bhatt and Mathew [ABM19]. Our results can also be viewed as some computations in this direction.

Outline of the paper. The statement of Theorem 1.2 presents the need for having a theory of crystalline cohomology for stacks. This theory has already been developed by Olsson in [Ol07] using the lisse-étale crystalline site of an algebraic stack. In Section 2, we provide another definition of crystalline cohomology of algebraic stacks through syntomic descent relying on the work of Fontaine and Messing [FM87]. In Proposition 2.16 we prove that the definition via syntomic descent and the definition via the lisse-étale crystalline site are equivalent.

In Section 3.1, we provide a brief review of crystalline Dieudonné theory and prove Theorem 1.2. Our proof uses a description of $M(G)$ obtained in the work of Berthelot, Messing and Breen [BBM82] which expresses $M(G)$ as a certain Ext group in the crystalline topos which is recalled in Theorem 3.8. The main ingredient of our proof of Theorem 1.2 is to obtain a suitable spectral sequence computing $H_{crys}^*(BG)$ which is done in Proposition 3.13. In order to obtain the spectral sequence in Proposition 3.13, we use Čech descent along the syntomic map $* \rightarrow BG$. In Section 3.2, we deduce the analogue of Theorem 1.2 for p -divisible groups. In Section 3.3, we provide a complete calculation of crystalline cohomology of the classifying stack BA of an abelian variety A . This is done in Proposition 3.28, where we prove that $H_{crys}^{2*}(BA) \cong \text{Sym}^*(H_{crys}^1(A))$ and $H_{crys}^i(BA) = 0$ for odd i . An analogue of this result in ℓ -adic cohomology was proved by Behrend [Beh03, Thm. 6.1.6]. Our proof of Proposition 3.28 uses different techniques. We rely on some explicit simplicial constructions and some computations from the theory of non-abelian derived functors. Using these calculations, in Section 3.4, we completely calculate the crystalline cohomology of BG for a p -divisible group G .

In Section 4, we enter the mixed characteristic situation. We begin by recalling the work of Anschütz and Le Bras and define the contravariant prismatic Dieudonné module $M_{\Delta}(G)$ for a p -divisible group G over a quasi-regular semiperfectoid base ring R as an Ext^1 group in the prismatic topos. This definition is compatible with the definition in [SW14] as proven in [ALB19, Prop. 4.3.7.]. We will then define quasi-syntomic stacks and define the classifying stack BG of G as a quasi-syntomic stack in Definition 4.27. In Definition 4.13, we extend the notion of prismatic cohomology developed by Bhatt and Scholze [BS19] to quasi-syntomic stacks via quasi-syntomic descent. Then analogous to Proposition 3.16, in Proposition 4.33 we prove that $M_{\Delta}(G) \simeq H_{\Delta}^2(BG)$, where the latter denotes the prismatic cohomology of BG . Similar to the crystalline case, in Proposition 4.40 we obtain a complete calculation of prismatic cohomology of the classifying stack $B\widehat{A}$, where \widehat{A} is the p -adic completion of an abelian scheme A .

Acknowledgements. I am grateful to Bhargav Bhatt for suggesting that Dieudonné modules could be recoverable via crystalline cohomology of classifying stacks as well as numerous helpful conversations regarding various aspects of this paper. I am also thankful to Igor Kriz, Arthur-César Le Bras, Shizhang Li and Martin Olsson for helpful conversations. Special thanks to the anonymous referee for offering many suggestions on an earlier version of this paper. During the preparation of this paper, I was partially supported by the Rackham International Students Fellowship.

2 Crystalline cohomology

2.1 Crystalline cohomology for stacks

We gather some notations for this section. We fix a prime p . Let k be a perfect field of characteristic $p > 0$ and let $W(k)$ be the ring of Witt vectors of k . Let SYNSch_k denote the big Grothendieck site of schemes over k with the syntomic topology. Let $D(\text{SYNSch}_k, W(k))$ denote the unbounded derived category of sheaves of $W(k)$ -modules and let $D(W(k))$ denote the unbounded derived category of $W(k)$ -modules (both can be equipped with natural ∞ -catgoerical enhancements). For a given scheme $X \in \text{SYNSch}_k$, there is a functor $R\Gamma(X, \cdot) : D(\text{SYNSch}_k, W(k)) \rightarrow D(W(k))$. Let $R\Gamma_{\text{crys}}(X) \in D(W(k))$ denote the crystalline cohomology of X .

Proposition 2.1. *There is an object $R\mathcal{O}^{\text{crys}} \in D(\text{SYNSch}_k, W(k))$ such that $R\Gamma(X, R\mathcal{O}^{\text{crys}}) \simeq R\Gamma_{\text{crys}}(X)$.*

Proof. This is a formal corollary of a result of Fontaine and Messing. Using [FM87, Prop. 1.3], we get that for any n , there is a sheaf $\mathcal{O}^{\text{crys}}/W_n$ on SYNSch_k such that the n -truncated crystalline cohomology $(R\Gamma_{\text{crys}}/W_n)(X) \simeq R\Gamma(X, \mathcal{O}^{\text{crys}}/W_n)$. Now we define $R\mathcal{O}^{\text{crys}} := R\varprojlim_n \mathcal{O}^{\text{crys}}/W_n$. Then it follows that $R\Gamma(X, R\mathcal{O}^{\text{crys}}) \simeq \varprojlim_n (R\Gamma_{\text{crys}}/W_n)(X) \simeq R\Gamma_{\text{crys}}(X)$. \square

Definition 2.2. Let \mathcal{X} be an algebraic stack over k . Let $C_{\mathcal{X}}$ denote the category of schemes X over \mathcal{X} . We define the *crystalline cohomology of the algebraic stack \mathcal{X}* as

$$R\Gamma_{\text{crys}}(\mathcal{X}) := R\varprojlim_{X \in C_{\mathcal{X}}} R\Gamma_{\text{crys}}(X) \in D(W(k)).$$

Remark 2.3. Alternatively, given an algebraic stack \mathcal{X} one can consider the slice site $\mathcal{X}_{\text{syn}} := \text{SYNSch}_{k/\mathcal{X}}$. Then the object $R\mathcal{O}^{\text{crys}}$ induces an object in the derived category of sheaves of $W(k)$ -modules on \mathcal{X}_{syn} which by abuse of notation we again write as $R\mathcal{O}^{\text{crys}}$. Then $R\Gamma_{\text{crys}}(\mathcal{X}) \simeq R\Gamma(\mathcal{X}_{\text{syn}}, R\mathcal{O}^{\text{crys}})$.

Remark 2.4. By Definition 2.2 and Zariski descent, we observe that $R\Gamma_{\text{crys}}(\mathcal{X}) \simeq R\varprojlim_{X \in C_{\mathcal{X}}^{\text{aff}}} R\Gamma_{\text{crys}}(X)$ where $C_{\mathcal{X}}^{\text{aff}}$ denotes the category of affine schemes over \mathcal{X} .

Remark 2.5. Let $X \rightarrow \mathcal{X}$ be a syntomic cover of the algebraic stack \mathcal{X} by a scheme X . Let X_{\bullet} be the simplicial algebraic space given by the Čech nerve of $X \rightarrow \mathcal{X}$ (so that X_n is the n -fold fibre product of X over \mathcal{X} , which is, in general, only an algebraic space.) Then we have that $R\Gamma_{\text{crys}}(\mathcal{X}) \simeq R\varprojlim R\Gamma_{\text{crys}}(X_{\bullet})$.

2.2 The crystalline site

In this section, we provide a brief reminder of definitions of the (big) crystalline site and the associated topos. We prefer to work with the big site because it is functorial and still computes the same cohomology groups. Our exposition roughly follows [BBM82].

We fix a prime p as before. Let k be a fixed field of characteristic p and let $W(k)$ be the ring of Witt vectors. Let $(W(k), p, \gamma)$ be the usual divided power structure on $W(k)$.

Definition 2.6. We denote by $\text{Crys}(k/W(k))$ a category whose objects are given by the data of an k -scheme U ; a $W(k)$ -scheme T on which p is locally nilpotent; a closed $W(k)$ -immersion $i : U \rightarrow T$ and a divided power structure δ on the ideal sheaf corresponding to the closed immersion i , which we require to be compatible with the divided power structure γ . We will denote an object of $\text{Crys}(k/W(k))$ by (U, T, δ) or simply by (U, T) when no confusion is likely to arise. A morphism in this category is given by a pair of morphisms $u : U' \rightarrow U$ and $v : T' \rightarrow T$, where u is a k -morphism and v is a $W(k)$ -morphism compatible with divided powers such that $v \circ i' = i \circ u$.

Definition 2.7. A family $(U_i, T_i) \rightarrow (U, T)$ of maps in $\text{Crys}(k/W(k))$ is a τ -covering if $U_i = U \times_T T_i$ for all i and $\{T_i \rightarrow T\}$ is a τ -covering. Here τ could be Zariski, étale, smooth, syntomic or fppf. This equips the category $\text{Crys}(k/W(k))$ with a Grothendieck topology and we denote the site we obtain this way by $\text{Crys}(k/W(k))_{\tau}$. We let $(k/W(k))_{\text{Crys}, \tau}$ denote the associated topos. We define a presheaf $\mathcal{O}^{\text{crys}}(U, T) := \Gamma(T, \mathcal{O}_T)$. This is a sheaf of rings on the site $\text{Crys}(k/W(k))_{\tau}$.

Definition 2.8. Let X be a scheme over k . By \underline{X} we denote a sheaf on $\text{Crys}(k/W(k))_\tau$ defined by $\underline{X}(U, T) := X(U) = \text{Hom}_k(U, X)$. cf. [BBM82, 1.1.4.5]. This is a sheaf for any τ , where τ is from the list mentioned in Definition 2.7.

Remark 2.9. We will write $\tau \leq \tau'$ while comparing different topologies to mean that τ' is finer than τ .

Proposition 2.10. *Let G be an abelian sheaf on $\text{Crys}(k/W(k))_{\tau'}$. Then*

$$R\text{Hom}_{\text{Crys}(k/W(k))_\tau}(G, \mathcal{O}^{\text{crys}}) \simeq R\text{Hom}_{\text{Crys}(k/W(k))_{\tau'}}(G, \mathcal{O}^{\text{crys}})$$

for $\tau \leq \tau'$.

Proof. This is [BBM82, Prop. 1.3.6]. The proof relies on [BBM82, Prop. 1.1.19]. \square

Proposition 2.11. *Let X be a scheme over k . Then*

$$R\Gamma_{\text{crys}}(X) \simeq R\text{Hom}_{\text{Crys}(k/W(k))_\tau}(\mathbb{Z}[\underline{X}], \mathcal{O}^{\text{crys}}),$$

where one can choose any topology τ from the list mentioned in Definition 2.7. Here $\mathbb{Z}[\underline{X}]$ denotes the “free abelian group” on \underline{X} in the topos $(k/W(k))_{\text{Crys}, \tau}$.

Proof. When τ is Zariski topology, this follows from [BBM82, Prop. 1.3.4] and by definition of crystalline cohomology. The rest follows from Proposition 2.10. \square

Remark 2.12. Therefore, we see that it does not matter which topology τ we choose while computing crystalline cohomology. We will often choose $\tau = \text{Zariski}$ and in this case we will omit τ from the notation and write the relevant site as $\text{Crys}(k/W(k))$ and the topos as $(k/W(k))_{\text{Crys}}$.

Remark 2.13. One can also define the “truncated crystalline sites” by replacing $W(k)$ by $W_n(k)$ in Definition 2.6 and Definition 2.7 for each $n \geq 1$. We denote the corresponding site (equipped with the Zariski topology for simplicity) by $\text{Crys}(k/W_n(k))$ and the associated topos by $(k/W_n(k))_{\text{Crys}}$. This has a sheaf of rings $\mathcal{O}^{\text{crys}}$ defined in a way similar to Definition 2.7. Analogous to Proposition 2.11, one has that for a scheme X over k , $R\Gamma_{\text{crys}}(X/W_n) \simeq R\text{Hom}_{\text{Crys}(k/W_n(k))}(\mathbb{Z}[\underline{X}], \mathcal{O}^{\text{crys}})$, where \underline{X} is the sheaf on $\text{Crys}(k/W_n(k))$ defined in a way similar to Definition 2.8.

Now we provide a different definition of crystalline cohomology of an algebraic stack following [Ol07, §2.7.1]. Then we will prove that this definition is equivalent to Definition 2.2.

Definition 2.14. Let \mathcal{X} be an algebraic stack over k . We define the *lisse-étale crystalline* site of \mathcal{X} first as the category $\text{Crys}(\mathcal{X})_{\text{lisse-ét}}$ whose objects are the data of a k -schemes U with a smooth map $U \rightarrow \mathcal{X}$ over k ; a $W(k)$ -scheme T on which p is locally nilpotent; a closed $W(k)$ -immersion $i : U \rightarrow T$ and a divided power structure δ on the ideal sheaf corresponding to the closed immersion i , which we require to be compatible with $(W(k), p, \gamma)$. We will denote an object of $\text{Crys}(\mathcal{X})_{\text{lisse-ét}}$ by (U, T, δ) or simply by (U, T) when no confusion is likely to arise. The morphisms of this category is defined in the obvious way. A family $(U_i, T_i) \rightarrow (U, T)$ of maps in $\text{Crys}(\mathcal{X})_{\text{lisse-ét}}$ is covering if $U_i = U \times_T T_i$ for all i and $\{T_i \rightarrow T\}$ is an étale covering. This equips $\text{Crys}(\mathcal{X})_{\text{lisse-ét}}$ with a Grothendieck topology and the resulting site is called the *lisse étale crystalline* site. Let $\mathcal{O}^{\text{crys}}(U, T) := \Gamma(T, \mathcal{O}_T)$ be a sheaf of rings on $\text{Crys}(\mathcal{X})_{\text{lisse-ét}}$.

Definition 2.15. We define $R\Gamma_{\text{lisse-ét-crys}}(\mathcal{X}) := R\Gamma(\text{Crys}(\mathcal{X})_{\text{lisse-ét}}, \mathcal{O}^{\text{crys}})$.

Proposition 2.16. $R\Gamma_{\text{lisse-ét-crys}}(\mathcal{X}) \simeq R\Gamma_{\text{crys}}(\mathcal{X})$, i.e., Definition 2.15 is consistent with Definition 2.2.

Proof. We break down the proof in a few steps.

Step 1. First we fix some notations for the proof. We let $\text{Crys}(\mathcal{X})_{\text{ét}}$ denote the big variant of the site in Definition 2.14 where we remove the hypothesis that for a pair (U, T) the map $U \rightarrow \mathcal{X}$ had to be smooth and the covers are still given by étale covers. Sending $(U, T) \rightarrow \Gamma(T, \mathcal{O}_T)$ defines a sheaf of rings on both of these sites which we will denote by $\mathcal{O}^{\text{crys}}$ in both cases when no confusion is likely to occur. Let $\text{Crys}(\mathcal{X})_{\text{syn}}$ denote the variant of $\text{Crys}(\mathcal{X})_{\text{ét}}$ with the same underlying category but now the covers are given by syntomic

covers. Lastly we denote the big syntomic site of \mathcal{X} by \mathcal{X}_{syn} which is the full subcategory of schemes over \mathcal{X} and the topology is given by syntomic coverings.

Step 2. In this step, we show that crystalline cohomology can be computed in the big site $\text{Crys}(\mathcal{X})_{\acute{e}t}$. We have an inclusion functor

$$\text{Crys}(\mathcal{X})_{lis-\acute{e}t} \rightarrow \text{Crys}(\mathcal{X})_{\acute{e}t}$$

which is continuous and cocontinuous and therefore gives a map of topoi

$$i : \text{ShvCrys}(\mathcal{X})_{lis-\acute{e}t} \rightarrow \text{ShvCrys}(\mathcal{X})_{\acute{e}t}.$$

We want to prove that there is a natural isomorphism $R\Gamma_{lis-\acute{e}t}(\mathcal{X}, i^{-1}\mathcal{O}^{crys}) \simeq R\Gamma_{\acute{e}t}(\mathcal{X}, \mathcal{O}^{crys})$. Thus, it is enough to prove that i^{-1} has a left adjoint $i_!$ which is exact. Following the method of proof in [SP, Tag 07IJ], it is enough to show that $i_! * = *$ where $*$ denotes the final object.

Let $Y \rightarrow \mathcal{X}$ be a smooth cover of \mathcal{X} by a scheme Y . First we prove that for any given object $(U, T) \in \text{Crys}(\mathcal{X})_{\acute{e}t}$, there exists a cover $(U_i, T_i) \rightarrow (U, T)$ in $\text{Crys}(\mathcal{X})_{\acute{e}t}$ such that for all i there are maps $U_i \rightarrow Y$ over \mathcal{X} . In order to show this, first we base change to get a smooth map $U \times_{\mathcal{X}} Y \rightarrow U$ which admits a projection map $U \times_{\mathcal{X}} Y \rightarrow Y$. Now we pick an étale cover $U' \rightarrow U \times_{\mathcal{X}} Y$ where U' is a scheme and thus we get a smooth map $U' \rightarrow U$. This can be refined further to obtain maps $U_i \rightarrow U'$ such that the composition $\{U_i \rightarrow U' \rightarrow U\}$ is an étale cover. Now $i : U \rightarrow T$ is a divided power thickening of U with p being locally nilpotent on T . Therefore $i : U \rightarrow T$ is a universal homeomorphism. By invariance of étale sites [SP, Tag 04DY] for universal homeomorphisms, we get étale maps $T_i \rightarrow T$ such that $T_i \times_T U \simeq U_i$. This gives a thickening $U_i \rightarrow T_i$. Since the map $T_i \rightarrow T$ is étale, by [SP, Tag 07H1], the ideal of the thickening $U_i \rightarrow T_i$ has divided powers. Further, since $\{U_i \rightarrow U\}$ was an étale cover, $\{T_i \rightarrow T\}$ is also an étale cover by construction. Thus we have constructed a refinement $(U_i, T_i) \rightarrow (U, T)$ in the site $\text{Crys}(\mathcal{X})_{\acute{e}t}$. For all i , we have also constructed maps $U_i \rightarrow Y$ over \mathcal{X} .

Now by using the proof in [SP, Tag 07IJ], we fix a set of divided power thickenings $(U_s, T_s)_{s \in S}$ by picking $(U_s)_{s \in S} = (\text{Spec } C_s)_{s \in S}$ to be an affine Zarisky open cover of the scheme Y , and for each $s \in S$, building T_s by using the algebra C_s as in [SP, Tag 07HP]. Since the map $Y \rightarrow \mathcal{X}$ was smooth, it follows that $(U_s, T_s) \in \text{Crys}(\mathcal{X})_{lis-\acute{e}t}$. Using the discussion from the previous paragraph and by the construction of $(U_s, T_s)_{s \in S}$, we get an epimorphism $\coprod_{s \in S} h_{(U_s, T_s)} \rightarrow *$ in $\text{ShvCrys}(\mathcal{X})_{\acute{e}t}$, which implies $i_! * = *$ proving our main claim in Step 2.

Step 3. In this step, we show that crystalline cohomology can also be computed in the finer site $\text{Crys}(\mathcal{X})_{syn}$. We have a map of topoi

$$v : \text{ShvCrys}(\mathcal{X})_{syn} \rightarrow \text{ShvCrys}(\mathcal{X})_{\acute{e}t}.$$

Let v_* denote the corresponding right adjoint. The left adjoint v^{-1} is given by sheafification. Thus we obtain a natural isomorphism $R\Gamma_{syn}(\mathcal{X}, \mathcal{O}^{crys}) \simeq R\Gamma_{\acute{e}t}(\mathcal{X}, Rv_*\mathcal{O}^{crys})$. We claim that $Rv_*\mathcal{O}^{crys} \simeq v_*\mathcal{O}^{crys} = \mathcal{O}^{crys}$. This statement can be checked locally and thus follows from the scheme case where it is already known. We refer to [BBM82, Prop. 1.1.19] for a proof in the case of schemes.

Step 4. In this step, we show that crystalline cohomology can be computed in \mathcal{X}_{syn} as well and conclude the proof. We have a functor

$$h : \text{Crys}(\mathcal{X})_{syn} \rightarrow \mathcal{X}_{syn}$$

which sends $(V, T) \rightarrow V$. By the proof of [FM87, Prop. 1.3], this functor is cocontinuous. By [SP, Tag 00XI], there is a morphism of topoi

$$u : \text{ShvCrys}(\mathcal{X})_{syn} \rightarrow \text{Shv}\mathcal{X}_{syn},$$

where $u_*(\mathcal{F})(U) := \varinjlim_{(V, T)/U} \mathcal{F}(V, T)$. We write the left adjoint by u^{-1} which is exact by [SP, Tag 00XL]. By adjunction, it follows that u_* commutes with the global section functor: indeed, we obtain $\text{Hom}(*, u_*\mathcal{F}) \simeq \text{Hom}(*, \mathcal{F})$. Therefore it follows that $R\Gamma_{syn}(\mathcal{X}, \mathcal{O}^{crys}) \simeq R\Gamma R u_*\mathcal{O}^{crys}$. Now the right hand side is equivalent to Definition 2.2. Therefore we are done. \square

3 Application to crystalline Dieudonné theory

In this section, we apply the notion of crystalline cohomology of stacks to describe the Dieudonné module of a finite group scheme of p -power order or a p -divisible group as the crystalline cohomology of the classifying stack. Before we do that, we briefly remind the reader the main theorem of contravariant Dieudonné theory and the related definitions.

Definition 3.1. Let k be a perfect field and $W(k)$ the ring of Witt vectors of k . Let σ denote the Witt vector Frobenius which is induced from the Frobenius in k . The Dieudonné ring \mathcal{D}_k is defined to be the free non-commutative polynomial ring in two generators F, V over $W(k)$ subjected to the relations $FV = VF = p$, $Fc = \sigma(c)F$ for $c \in W(k)$, and $cV = V\sigma(c)$ for $c \in W(k)$. \mathcal{D}_k is non-commutative when $k \neq \mathbb{F}_p$, and is $\mathbb{Z}_p[x, y]/(xy - p)$ when $k = \mathbb{F}_p$.

Definition 3.2. We let W_n denote the group scheme that corepresents the functor that sends a k -algebra A to the ring of length n Witt vectors $W_n(A)$. These group schemes are naturally induced with a Frobenius endomorphism F on them and we define W_n^m to be the group scheme which is the kernel of F^m on W_n . We also have a map $V : W_n \rightarrow W_{n+1}$ induced by the Witt vector Verschiebung which turns $\{W_n\}_{n \geq 1}$ into a directed system of group schemes. We define $CW^u := \varinjlim W_n$. One defines the formal p -group CW of Witt covectors as a completion of CW^u in a suitable sense. We refer to [Fo77] for details.

Example 3.3. By definition, W_1 is the additive group scheme \mathbb{G}_a . Also, W_1^1 is the finite additive group scheme with the underlying scheme given by $\text{Spec } k[x]/x^p$ which is usually denoted as α_p . Similarly, W_1^n is the finite additive group scheme with the underlying scheme given by $\text{Spec } k[x]/x^{p^n}$, which is denoted as α_{p^n} . The Cartier dual of α_{p^n} is given by the group scheme W_n^1 . More generally, the Cartier dual of W_n^m is isomorphic to W_n^m [Dem72, Chapter III, §4].

Definition 3.4. For a finite group scheme G over k , we define the *Dieudonné module* of G to be

$$M(G) := \text{Hom}(G, CW)$$

where the Hom is being taken in the category of formal groups.

Now we are ready to state the classical theorem of contravariant Dieudonné theory, which can be found in [Fo77, Chapitre III, §1, Thm. 1].

Theorem 3.5. *The functor $G \rightarrow M(G)$ induces an anti-equivalence between the category of finite group schemes over k and left \mathcal{D}_k -modules with finite $W(k)$ length.*

Example 3.6. Let us give some examples of Dieudonné modules associated to certain finite flat group schemes. For the group schemes W_n^m from Example 3.3, we have $M(W_n^m) \simeq \mathcal{D}_k/\mathcal{D}_k \cdot (F^m, V^n)$ [Oor66, §15.4]. In particular, $M(\alpha_{p^n}) \simeq \mathcal{D}_k/\mathcal{D}_k \cdot (F^n, V)$. For the multiplicative group scheme underlying p^n -th roots of unity denoted as μ_{p^n} , we have $M(\mu_{p^n}) \simeq \mathcal{D}_k/\mathcal{D}_k \cdot (F^n, (V - 1))$. For the constant group scheme $\mathbb{Z}/p^n\mathbb{Z}$, we have $M(\mathbb{Z}/p^n\mathbb{Z}) \simeq \mathcal{D}_k/\mathcal{D}_k \cdot ((F - 1), V^n)$.

Example 3.7. Let $E[p]$ denote the p -torsion group scheme of a supersingular elliptic curve E over k . Then $M(E[p]) \simeq \mathcal{D}_k/\mathcal{D}_k \cdot (F^2, F - V, V^2)$. We refer to [Oor66, §15.5] for more details.

In [BBM82], Berthelot, Breen and Messing obtained an alternative description of $M(G)$. Given a finite group scheme G , using Definition 2.8, we obtain an abelian group object \underline{G} of the topos $(k/W(k))_{\text{Crys}}$. They prove the following theorem which expresses $M(G)$ as a certain Ext group in the crystalline topos.

Theorem 3.8 ([BBM82, Thm. 4.2.14]). *For a finite group scheme G for p power rank, we have $\sigma^*M(G) \simeq \text{Ext}_{(k/W(k))_{\text{Crys}}}^1(\underline{G}, \mathcal{O}^{\text{crys}})$, where $\sigma^*M(G)$ denotes the extension of scalars along the Witt vector Frobenius σ ; i.e., $\sigma^*M(G) := M(G) \otimes_{W(k), \sigma} W(k)$.*

3.1 Dieudonné module of finite group schemes

In this subsection, we work with finite group schemes G of p -power rank over a perfect field k . Let $BG := [\mathrm{Spec} k/G]$ be the classifying stack of G [SP, Tag 044O]. Any such group scheme is a local complete intersection and therefore it follows that the map $\mathrm{Spec} k \rightarrow BG$ is a syntomic cover. Its Čech nerve is given by the following simplicial scheme:

$$\cdots G \times G \times G \rightrightarrows G \times G \rightrightarrows G \rightrightarrows \mathrm{Spec} k .$$

As in Definition 2.8, we can attach a sheaf \underline{G} on $\mathrm{Crys}(k/W(k))$ which can be viewed as an object of the topos $(k/W(k))_{\mathrm{Crys}}$. Corresponding to the Čech nerve above, we obtain by functoriality, a simplicial object of the topos:

$$B\underline{G} := \cdots \underline{G} \times \underline{G} \times \underline{G} \rightrightarrows \underline{G} \times \underline{G} \rightrightarrows \underline{G} \rightrightarrows * .$$

where $*$ is the final object. With this simplicial object, we can attach the free simplicial abelian group object:

$$\cdots \mathbb{Z}[\underline{G} \times \underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G}] \rightrightarrows \mathbb{Z} .$$

The alternating face map complex associated to the above simplicial object is as an object in the category of complexes of abelian objects of the topos $(k/W(k))_{\mathrm{Crys}}$ which can also be viewed as an object in the derived category of abelian objects of $(k/W(k))_{\mathrm{Crys}}$ denoted as $D(k/W(k))$. This object living in $D(k/W(k))$ is also isomorphic to the homotopy colimit of

$$\cdots \mathbb{Z}[\underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G}] \rightrightarrows \mathbb{Z} .$$

viewed as a functor from Δ^{op} to $D(k/W(k))$. We denote this object by $\mathbb{Z}[B\underline{G}] \in D(k/W(k))$. If we work with the truncated crystalline site $\mathrm{Crys}(k/W_n(k))$, all of these constructions remain valid and one can associate $\mathbb{Z}[B\underline{G}] \in D(k/W_n(k))$ as well.

Lemma 3.9. $H^0(\mathbb{Z}[B\underline{G}]) \simeq \mathbb{Z}$ and $H^{-1}(\mathbb{Z}[B\underline{G}]) \simeq \underline{G}$.

Proof. First we work with the presheaf simplicial abelian group object

$$\cdots \mathbb{Z}^{\mathrm{pre}}[\underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}^{\mathrm{pre}}[\underline{G}] \rightrightarrows \mathbb{Z} .$$

which is obtained by applying the free abelian presheaf functor to $B\underline{G}$ [SP, Tag 03CP]. Let $\mathbb{Z}^{\mathrm{pre}}[B\underline{G}]$ be the associated complex: $K^\bullet := \cdots \mathbb{Z}^{\mathrm{pre}}[\underline{G} \times \underline{G}] \rightarrow \mathbb{Z}^{\mathrm{pre}}[\underline{G}] \rightarrow \mathbb{Z} \rightarrow 0$. We see that H^0 of this complex is \mathbb{Z} since the last differential is zero. One also notes that K^\bullet computes the (presheaf) group cohomology of the (presheaf) abelian group \underline{G} with constant coefficients in \mathbb{Z} and hence, $H^{-1}(K^\bullet) \simeq \underline{G}$. Now since sheafification is exact, we obtain the required statements. \square

Lemma 3.10. *Let G be a group scheme of order p^m . Then $H^{-i}(\mathbb{Z}[B\underline{G}])$ is killed by p^m for $i > 0$.*

Proof. We start by recalling some definitions for this proof. Let A be an ordinary n torsion abelian group. Let us define $\mathbb{Z}[BA]$ to be the alternating face complex associated to the simplicial abelian group

$$\cdots \mathbb{Z}[A \times A] \rightrightarrows \mathbb{Z}[A] \rightrightarrows \mathbb{Z} .$$

Then as noted previously, $H^{-i}(\mathbb{Z}[BA]) \simeq H_i(A)$, where the latter denotes group cohomology with coefficients in \mathbb{Z} . This is also the homology of the Eilenberg-MacLane space $K(A, 1)$. We will show that $H_i(A)$ is n -torsion (*). First, from the complex $\mathbb{Z}[BA]$, one notes that $H_i(A)$ commutes with filtered colimits as a functor defined on abelian groups. Since A is an n -torsion abelian group, it can be expressed as filtered colimit of finite n -torsion abelian groups. Therefore, it is enough to check the statement for an n -torsion finite abelian group A . Further, one notes that $K(A_1, 1) \times K(A_2, 1) \simeq K(A_1 \times A_2, 1)$, therefore, by using the Künneth formula, we are reduced to checking this for n -torsion cyclic groups, where it follows from the well-known computation of group homology of finite cyclic groups using the Tate complex. The statement in the lemma now follows from applying (*) to $\mathbb{Z}^{\mathrm{pre}}[B\underline{G}]$ and noting that since G has order p^m , one has $p^m \cdot G = 0$. \square

Remark 3.11. These lemmas clearly remain valid even if we were working with the truncated crystalline sites $\text{Crys}(k/W_n(k))$ mentioned in Remark 2.13.

Proposition 3.12. $R\Gamma_{\text{crys}}(BG) \simeq R\text{Hom}_{D(k/W(k))}(\mathbb{Z}[B\mathcal{G}], \mathcal{O}^{\text{crys}})$.

Proof. Since $\text{Spec } k \rightarrow BG$ is a syntomic map, we can apply Remark 2.5 which gives that

$$R\Gamma_{\text{crys}}(BG) \simeq R\varprojlim (R\Gamma_{\text{crys}}(\text{Spec } k) \rightrightarrows R\Gamma_{\text{crys}}(G) \rightrightarrows R\Gamma_{\text{crys}}(G \times G) \cdots).$$

Which by Proposition 2.11 is

$$\simeq R\varprojlim (R\text{Hom}_{D(k/W(k))}(\mathbb{Z}, \mathcal{O}^{\text{crys}}) \rightrightarrows R\text{Hom}_{D(k/W(k))}(\mathbb{Z}[\mathcal{G}], \mathcal{O}^{\text{crys}}) \rightrightarrows R\text{Hom}_{D(k/W(k))}(\mathbb{Z}[\mathcal{G} \times \mathcal{G}], \mathcal{O}^{\text{crys}}) \cdots).$$

We can take the $R\varprojlim$ inside as a homotopy colimit, which gives us that the above is $\simeq R\text{Hom}_{D(k/W(k))}(\mathbb{Z}[B\mathcal{G}], \mathcal{O}^{\text{crys}})$, as desired. \square

Proposition 3.13. *There is a spectral sequence with E_2 -page*

$$E_2^{i,j} = \text{Ext}_{(k/W(k))_{\text{crys}}}^i(H^{-j}(\mathbb{Z}[B\mathcal{G}]), \mathcal{O}^{\text{crys}}) \implies H_{\text{crys}}^{i+j}(BG),$$

and another spectral sequence with E_1 -page

$$E_1^{i,j} = H_{\text{crys}}^j(G^i) \implies H_{\text{crys}}^{i+j}(BG),$$

where G^i denotes the i fold fibre product of G with itself. By convention, $G^0 = *$.

Proof. This is a consequence of [SP, Tag 07A9], Proposition 3.12 and Proposition 2.11. \square

Proposition 3.14. $H_{\text{crys}}^1(BG) = 0$.

Proof. We can use the E_2 spectral sequence from Proposition 3.13. to compute $H_{\text{crys}}^i(BG)$. We note that $\text{Ext}^i(\mathbb{Z}, \mathcal{O}^{\text{crys}}) = 0$ for $i > 0$ as it computes the cohomology of $\text{Spec } k$ for a perfect field k by Proposition 2.11. Also, by [BBM82, Prop. 4.2.6], we have $\text{Hom}(\mathcal{G}, \mathcal{O}^{\text{crys}}) = 0$. These calculations along with the spectral sequence and Lemma 3.9 imply $H_{\text{crys}}^1(BG) = 0$. \square

Proposition 3.15. *If G is a finite group scheme of p -power order, then for any $i > 0$, $H_{\text{crys}}^i(BG)$ is killed by a power of p as an abelian group.*

Proof. This follows from the spectral sequence in Proposition 3.13. Indeed, already in the E_2 -page of the spectral sequence, all but $E_2^{0,0}$ is p -power torsion by Lemma 3.10. Hence, all but $E_\infty^{0,0}$ is p -power torsion as well. Therefore, for any $i > 0$, $H_{\text{crys}}^i(BG)$ has a finite filtration whose successive quotients are p -power torsion and hence is p -power torsion itself. \square

Proposition 3.16. *If G is a finite group scheme of p -power order, then $H_{\text{crys}}^2(BG) \simeq \text{Ext}_{(k/W(k))_{\text{crys}}}^1(\mathcal{G}, \mathcal{O}^{\text{crys}})$.*

Proof. First we note that there is a natural map $\text{Ext}_{(k/W(k))_{\text{crys}}}^1(\mathcal{G}, \mathcal{O}^{\text{crys}}) \rightarrow H_{\text{crys}}^2(BG)$, which is injective. Indeed, from the E_2 -spectral sequence (which is natural in G), we note that $\text{Fil}^0 H_{\text{crys}}^2(BG) = H_{\text{crys}}^2(BG)$, $\text{Fil}^i H_{\text{crys}}^2(BG) = 0$ for $i \geq 2$ and therefore, $\text{Fil}^1 H_{\text{crys}}^2(BG) = \text{Ext}_{(k/W(k))_{\text{crys}}}^1(\mathcal{G}, \mathcal{O}^{\text{crys}})$, which gives the required injective map. We proceed to proving that this natural map is an isomorphism.

Let $R\Gamma_{\text{crys}}(BG/W_n) := R\Gamma_{\text{crys}}(BG) \otimes_{W(k)}^L W_n(k)$ and $H_{\text{crys}}^i(BG/W_n) := H^i(R\Gamma_{\text{crys}}(BG/W_n))$. Then we have the following exact sequence:

$$0 \rightarrow H_{\text{crys}}^1(BG)/p^n \rightarrow H_{\text{crys}}^1(BG/W_n) \rightarrow H_{\text{crys}}^2(BG)[p^n] \rightarrow 0.$$

Now we choose n large enough such that $p^n G = 0$ and $H_{\text{crys}}^2(BG)$ is p^n -torsion. This is possible by Proposition 3.15. Then the above exact sequence along with Proposition 3.14 gives $H_{\text{crys}}^1(BG/W_n) \simeq H_{\text{crys}}^2(BG)$ for such an n . By [BBM82, Prop. 4.2.17], we have $\text{Ext}_{(k/W(k))_{\text{crys}}}^1(\mathcal{G}, \mathcal{O}^{\text{crys}}) \simeq \text{Hom}_{(k/W_n(k))_{\text{crys}}}(\mathcal{G}, \mathcal{O}^{\text{crys}})$.

Therefore, it is enough to show that $H_{crys}^1(BG/W_n) \simeq \text{Hom}_{(k/W_n(k))_{Crys}}(\underline{G}, \mathcal{O}^{crys})$.

Since

$$R\Gamma_{crys}(BG) \simeq R\varprojlim (R\Gamma_{crys}(\text{Spec } k) \rightrightarrows R\Gamma_{crys}(G) \rightrightarrows R\Gamma_{crys}(G \times G) \cdots),$$

we obtain

$$R\Gamma_{crys}(BG/W_n) \simeq R\varprojlim (R\Gamma_{crys}(\text{Spec } k/W_n) \rightrightarrows R\Gamma_{crys}(G/W_n) \rightrightarrows R\Gamma_{crys}(G \times G/W_n) \cdots).$$

In order to justify the last step, we note that $W_n(k)$ is quasi-isomorphic to the complex $W(k) \xrightarrow{p^n} W(k)$; thus the functor $(\cdot) \otimes_{W(k)}^L W_n(k)$ commutes with the $R\varprojlim$ as required.

Therefore, by Remark 3.11 and the proof of Proposition 3.12, one obtains that

$$R\Gamma_{crys}(BG/W_n) \simeq R\text{Hom}_{D(k/W_n(k))}(\mathbb{Z}[B\underline{G}], \mathcal{O}^{crys}),$$

and analogous to Proposition 3.13 a spectral sequence with E_2 -page

$$E_2^{i,j} = \text{Ext}_{(k/W_n(k))_{Crys}}^i(H^{-j}(\mathbb{Z}[B\underline{G}], \mathcal{O}^{crys}) \implies H_{crys}^{i+j}(BG/W_n).$$

Now applying Lemma 3.9 yields the desired isomorphism. To check this, we see that $E_2^{1,0} = 0$ and $E_2^{0,1} = \text{Hom}_{(k/W_n(k))_{Crys}}(\underline{G}, \mathcal{O}^{crys})$. Also we have $E_2^{2,0} = 0$. Since $E_2^{i,j} = 0$ for $i < 0$ or $j < 0$, we obtain that $E_\infty^{0,1} = \text{Hom}_{(k/W_n(k))_{Crys}}(\underline{G}, \mathcal{O}^{crys})$. Therefore, we indeed obtain the required isomorphism $H_{crys}^1(BG/W_n) \simeq \text{Hom}_{(k/W_n(k))_{Crys}}(\underline{G}, \mathcal{O}^{crys})$. Relevant part of the E_2 page of the above spectral sequence is depicted in the diagram below.

$$\begin{array}{ccccccc}
 0 & & E_2^{0,2} & & E_2^{1,2} & & E_2^{2,2} \\
 & \searrow & & \searrow & & \searrow & \\
 0 & & \text{Hom}(\underline{G}, \mathcal{O}^{crys}) & & E_2^{1,1} & & E_2^{2,1} \\
 & \searrow & & \searrow & & \searrow & \\
 0 & & E_2^{0,0} = W_n(k) & & E_2^{1,0} = 0 & & E_2^{2,0} = 0 \\
 & \searrow & & \searrow & & \searrow & \\
 0 & & 0 & & 0 & & 0
 \end{array}$$

□

Proof of Theorem 1.2. Follows from Proposition 3.16 and Theorem 3.8. □

Remark 3.17. One can develop the theory of crystalline cohomology for higher stacks and study the cohomology of the n -stack $K(G, n)$ [To06]. For a group scheme G , the n -stack $K(G, n)$ is supposed to be analogue of the Eilenberg-MacLane space $K(G, n)$ for any discrete abelian group G which has the property that $\pi_i(K(G, n)) = G$ for $i = n$ and $\pi_i(K(G, n)) = 0$ for $i \neq \{n, 0\}$. In the topological case, there exists a chain complex of abelian groups $\mathbb{Z}[B^n G]$ such that $H_i(\mathbb{Z}[B^n G])$ computes the singular homology of the CW complex $K(G, n)$. Here, we do not define crystalline cohomology for higher stacks in general, however, for a finite group scheme G , we can work with an ad hoc definition generalizing Proposition 3.12. Similar to the definition of $\mathbb{Z}[B\underline{G}]$ as an object of the crystalline topos, one can also define $\mathbb{Z}[B^n \underline{G}]$. Then we define $R\Gamma_{crys}(K(G, n)) := R\text{Hom}_{(k/W(k))}(\mathbb{Z}[B^n \underline{G}], \mathcal{O}^{crys})$. Then from the analogue of the spectral sequence in Proposition 3.13, we obtain that $H_{crys}^i(K(G, n)) = 0$ for $0 < i < n + 1$ and $H_{crys}^{n+1}(K(G, n)) = M(G)$. In order to prove this for $n \geq 2$, the computation with spectral sequence relies on the fact that for an abelian group A , $H_i(K(A, n), \mathbb{Z}) = 0$ for $0 < i < n$, $H_n(K(A, n), \mathbb{Z}) = A$ and $H_{n+1}(K(A, n), \mathbb{Z}) = 0$. First two of these computations follow from Hurewicz theorem and the last one follows from applying the Serre fibration spectral sequence for the homotopy fibration sequence $K(A, n) \rightarrow * \rightarrow K(A, n + 1)$.

3.2 Dieudonné theory of p -divisible groups

At first, we recall the definition of a p -divisible group.

Definition 3.18. Let p be a prime and $h > 0$ an integer. A p -divisible group (or Barsotti Tate group) of height h over a scheme S is a directed system $G = \{G_n\}$ of finite flat group schemes over S such that each G_n is p^n -torsion of order p^{nh} and the transition map $i_n : G_n \rightarrow G_{n+1}$ is an isomorphism of G_n onto $G_{n+1}[p^n]$ for all $n \geq 1$. A morphism $f : G \rightarrow H$ between p -divisible groups is a compatible system of S -group maps $f_n : G_n \rightarrow H_n$ for all $n \geq 1$. If $S' \rightarrow S$ is a map of schemes then $G \times_S S' := G_n \times_S S'$ is the p -divisible group of height h over S' obtained by base change.

Example 3.19. If $A \rightarrow S$ is an abelian scheme with fibers of constant dimension $g > 0$ and p is a prime, then $G_n := A[p^n]$ is a finite flat p^n -torsion group scheme of order p^{2gn} for all $n \geq 1$ and $\{G_n\}$ is a directed system via isomorphisms $G_n \cong G_{n+1}[p^n]$ for all $n \geq 1$. For an abelian variety A , we denote this p -divisible group by $A[p^\infty]$. It has height $2g$ where $g = \dim A$.

We also recall the following theorem on Dieudonné theory of p -divisible groups, which can be found in [Fo77, Chapitre III, §6, Prop. 6.1].

Theorem 3.20. *The functor $G \rightarrow M(G) := \varprojlim M(G_n)$ is an anti-equivalence of categories between the category of p -divisible groups over k and the category of finite free $W(k)$ -modules D equipped with a Frobenius semilinear endomorphism $F : D \rightarrow D$ such that $pD \subseteq F(D)$. The height of G is the $W(k)$ -rank of $M(G)$.*

Now in order to formulate the analogue of Theorem 1.2 for p -divisible groups, we begin by making a definition.

Definition 3.21. Let G be a p -divisible group over k . We define the classifying stack of G to be the $BG := \varinjlim BG_n$ where the colimit is being taken in the category of stacks.

Proposition 3.22. *For a p -divisible group $G = \{G_n\}$ we have $R\Gamma_{\text{crys}}(BG) \simeq R\varprojlim R\Gamma_{\text{crys}}BG_n$.*

Proof. We write $\mathcal{F}_n := BG_n$ and $\mathcal{F} := BG$ for this proof. Using Remark 2.4, we see that crystalline cohomology of a stack \mathcal{Y} only depends on \mathcal{Y} viewed as a presheaf of groupoids on the category of affine k -schemes Aff_k . If $\mathcal{Y} := \text{colim} \mathcal{Y}_\alpha$ as presheaves of groupoids, then by the alternative definition in Remark 2.4 it follows that $R\Gamma_{\text{crys}}(\mathcal{Y}) \simeq R\lim_\alpha R\Gamma_{\text{crys}}(\mathcal{Y}_\alpha)$. Therefore, in order to show that $R\Gamma_{\text{crys}}(\mathcal{F}) \simeq R\varprojlim_n R\Gamma_{\text{crys}}(\mathcal{F}_n)$, it is enough to prove that \mathcal{F} is the colimit of \mathcal{F}_n in the category of presheaves of groupoids on Aff_k . This follows since affine schemes in the fpqc site of all schemes satisfy the property that any fpqc covering of an affine scheme has a finite subcovering by affine schemes. Indeed, by the property we mentioned, checking descent over an affine scheme is essentially a finite limit condition. Thus our claim follows since filtered colimit commutes with finite limits. \square

Proposition 3.23. *We have a natural isomorphism $H_{\text{crys}}^2(BG) \simeq M(G)$.*

Proof. We are interested to compute $H_{\text{crys}}^2(BG)$ which is $H^2(R\varprojlim R\Gamma_{\text{crys}}BG_n)$ by Proposition 3.22. We compute the cohomology of this \mathbb{N} -indexed derived limit by using [SP, Tag 07KY], which gives us the following short exact sequence

$$0 \rightarrow R^1 \varprojlim H_{\text{crys}}^1(BG_n) \rightarrow H_{\text{crys}}^2(BG) \rightarrow \varprojlim H_{\text{crys}}^2(BG_n) \rightarrow 0.$$

Therefore our claim follows from Proposition 3.14, Proposition 3.16, and Theorem 3.20. \square

3.3 Cohomology of the classifying stack of an abelian variety

Now we consider an abelian variety A over the field k . Let BA denote the classifying stack of A . In this section, in Proposition 3.28, we prove that $H_{\text{crys}}^{2*}(BA) \cong \text{Sym}^*(H_{\text{crys}}^1(A))$, and $H_{\text{crys}}^i(BA) = 0$ for odd i . In other words, $H_{\text{crys}}^*(BA) \cong \text{Sym}^*(H_{\text{crys}}^1(A))$, where $H_{\text{crys}}^1(A)$ is considered to be in degree 2.

Remark 3.24. An analogue of this proposition was proven by Borel in topology [Bo53], and by Behrend in ℓ -adic cohomology using fibration spectral sequences [Beh03, Thm. 6.1.6]. Here we take a different approach based on descent theory. One knows that there is a functorial isomorphism $H_{\text{crys}}^*(A) \cong \bigwedge^* H_{\text{crys}}^1(A)$. We show that Proposition 3.28 is a formal consequence of this isomorphism and the Künneth formula.

In this section, we will crucially use the theory of derived functors of non-additive functors which appears in [DP61], [II71]. Note that, analogously to Proposition 3.13, there is an E_1 spectral sequence with $E_1^{i,j} = H_{crys}^j(A^i) \implies H_{crys}^{i+j}(BA)$, where A^i denotes the i -fold fibre product of A with itself. We note that by definition, for $j \geq 1$, $E_1^{\bullet,j}$ is the alternating face complex associated to the cosimplicial object given by

$$(H_{crys}^j(*) \rightrightarrows H_{crys}^j(A) \rightrightarrows H_{crys}^j(A \times A) \cdots);$$

which is naturally isomorphic to the cosimplicial object

$$\bigwedge^j (H_{crys}^1(*) \rightrightarrows H_{crys}^1(A) \rightrightarrows H_{crys}^1(A \times A) \cdots).$$

Lemma 3.25. *The complex $E_1^{\bullet,1}$ is homotopy equivalent to $H_{crys}^1(A)[-1]$.*

Proof. As noted above, $E_1^{\bullet,1}$ is the alternating face complex associated to

$$(H_{crys}^1(*) \rightrightarrows H_{crys}^1(A) \rightrightarrows H_{crys}^1(A \times A) \cdots).$$

By Kunneth formula, one has a natural isomorphism $H_{crys}^1(A^i) \cong \bigoplus_{s=1}^i H_{crys}^1(A)$, and $H^1(*) = 0$. Writing $H_{crys}^1(A) = V$, we note that $E_1^{\bullet,1}$ is naturally isomorphic to the following explicit complex (where V sits in cohomological degree 1)

$$K_1^\bullet : \dots \rightarrow 0 \rightarrow V \rightarrow V^{\oplus 2} \rightarrow V^{\oplus 3} \rightarrow \dots$$

Here the first differential $d_1 : V \rightarrow V^{\oplus 2}$ is zero. In general, for $n \geq 2$, the differential $d_n : V^{\oplus n} \rightarrow V^{\oplus n+1}$ is given by

$$d_n(v_1, v_2, \dots, v_n) = (-v_1, 0, v_2 - v_3, 0, v_4 - v_5, 0, \dots, v_n)$$
 for even n ,

and

$$d_n(v_1, v_2, \dots, v_n) = (0, v_2, v_2, v_4, v_4, \dots, 0)$$
 for odd n .

One can check the above by induction. Now we consider the complex $K_2^\bullet : \dots \rightarrow 0 \rightarrow V \rightarrow 0 \rightarrow 0 \dots$ where V sits in cohomological degree 1. There are obvious maps from $K_1^\bullet \rightarrow K_2^\bullet$ and $K_2^\bullet \rightarrow K_1^\bullet$ (induced by id_V on cohomological degree 1) and we prove that they induce the desired homotopy equivalence. There is nothing to prove for the composite map from $K_2^\bullet \rightarrow K_2^\bullet$. For the other composite map $K_1^\bullet \rightarrow K_1^\bullet$, our task is to prove that it is homotopic to $\text{id}_{K_1^\bullet}$. We construct the required homotopy h_i . We set $h_i = 0$ for $i \leq 2$. For $i \geq 3$, now we define $h_i : V^{\oplus i} \rightarrow V^{\oplus i-1}$. We let $h_3(v_1, v_2, v_3) = (-v_1, v_3)$ and $h_4(v_1, \dots, v_4) = (0, v_2, 0)$. In general, we set

$$h_i(v_1, \dots, v_i) = (0, v_2, 0, \dots, 0, v_{i-2}, 0)$$
 for even $i \geq 6$,

and

$$h_i(v_1, \dots, v_i) = (-v_1, 0, -v_3, 0, \dots, 0, -v_{i-2}, v_i)$$
 for odd $i \geq 5$.

One easily checks that this gives the required homotopy. \square

Remark 3.26. For the cohomology of the complex K_1^\bullet , one can also use a less explicit argument. Writing $V = H_{crys}^1(A)$ as before, the addition $A \times A \rightarrow A$ induces a map $V \rightarrow V \oplus V$ which is dual to the addition on V^\vee . Therefore, one sees that the complex K_1^\bullet is the dual complex to BV^\vee (whose cohomology is concentrated in cohomological degree -1 and is equal to V^\vee). Applying duals back, one sees that cohomology of K_1^\bullet is concentrated in cohomological degree 1 and is equal to V . In fact, one can also prove that K_1^\bullet is homotopy equivalent to $V[-1]$ in this manner by noting that BV^\vee is homotopy equivalent to $V^\vee[1]$. We thank the referee for pointing this out.

Lemma 3.27. *The complex $E_1^{\bullet,n}$ is homotopy equivalent to $\text{Sym}^n(H_{crys}^1(A))[-n]$.*

Proof. Writing $H_{crys}^1(A) = V$ again, we note that by Lemma 3.25, $E_1^{\bullet,1}$ is homotopy equivalent to $V[-1]$. Therefore, the (derived) n -th exterior power $\wedge^n(V[-1])$ is homotopy equivalent to $\wedge^n(E_1^{\bullet,1})$ by [Il71, Def. 4.1.3.2] and Lemma 4.1.3.5 from loc. cit. It also follows from Definition 4.1.3.2 and Section 1.3.4 in loc. cit. that $\wedge^n(E_1^{\bullet,1})$ is homotopy equivalent to the alternating face complex associated to

$$\bigwedge^n (H_{crys}^1(*) \rightrightarrows H_{crys}^1(A) \rightrightarrows H_{crys}^1(A \times A) \cdots),$$

which is isomorphic to $E_1^{\bullet,n}$. Now one also notes that $\wedge^n(V[-1])$ is homotopy equivalent to $\text{Sym}^n(V)[-n]$, by the formula of décalage (i) in Proposition 4.3.2.1 loc. cit. This proves the lemma. \square

Proposition 3.28. *We have a natural isomorphism $H_{crys}^{2*}(BA) \cong \text{Sym}^*(H_{crys}^1(A))$, and $H_{crys}^i(BA) = 0$ for odd i .*

Proof. This follows from the existence of the E_1 spectral sequence analogous to Proposition 3.13 considered above (which degenerates at the E_2 page by Lemma 3.27), and the calculations from Lemma 3.25 and Lemma 3.27. The spectral sequence also guarantees the naturality of the isomorphisms. \square

Corollary 3.29. *For an abelian variety A over k , $H_{crys}^2(BA)$ is naturally isomorphic to the Dieudonné module associated to the p -divisible group $A[p^\infty]$.*

Proof. This follows from Proposition 3.28 and the fact that $H_{crys}^1(A)$ is isomorphic to the Dieudonné module associated to the p -divisible group $A[p^\infty]$. \square

3.4 Cohomology of the classifying stack a p -divisible group

Let G be a p -divisible group over a perfect field k . Using the calculation in Section 3.3, we are able to fully compute the cohomology ring $H_{crys}^*(BG)$. In this section, we prove that $H_{crys}^*(BG) \simeq \text{Sym}^*(M(G))$, where $M(G)$ is the Dieudonné module of G and it is considered to be in degree 2. Our strategy is to relate the p -divisible group to a suitable abelian variety and deduce the result from Proposition 3.28. First, we will record two lemmas.

Lemma 3.30. *Let G be a uniquely p -divisible abelian group. Then $H_i(G, k) = 0$ for $i > 0$ where $H_i(G, k)$ denotes the group homology with coefficients in the field k equipped with trivial G action.*

Proof. Since G is an abelian group there is an exact sequence $0 \rightarrow \mathbb{Z}^{\oplus I} \rightarrow \mathbb{Z}^{\oplus J} \rightarrow G \rightarrow 0$. By taking colimits over multiplication by p , we obtain an exact sequence $0 \rightarrow \mathbb{Z}[1/p]^{\oplus I} \rightarrow \mathbb{Z}[1/p]^{\oplus J} \rightarrow G \rightarrow 0$. Using the Hochschild–Serre spectral sequence, we are reduced to checking the claim for $G = \mathbb{Z}[1/p]^{\oplus I}$. By taking filtered colimits, we can assume that I is finite. Then by Kunneth formula it is enough to check the claim for $\mathbb{Z}[1/p]$. By taking filtered colimits over multiplication by p , it follows from the fact that $H_1(\mathbb{Z}, k) = k$ and $H_i(\mathbb{Z}, k) = 0$ for $i > 0$. \square

Lemma 3.31. *Let G be an abelian group such that multiplication by p is surjective on G . Then it follows that $W_n(k)[BG[p^\infty]] \rightarrow W_n(k)[BG]$ is an isomorphism in the derived category $D(W_n(k))$ of $W_n(k)$ -modules. Here $G[p^\infty] := \varinjlim G[p^n]$.*

Proof. By going modulo p , it would be enough to show that $k[BG[p^\infty]] \rightarrow k[BG]$ is quasi-isomorphism. For that, it would be enough to prove that the map induced on group homology $H_i(G[p^\infty], k) \rightarrow H_i(G, k)$ is an isomorphism. Here, in both cases group cohomology is taken with constant coefficients in k . We have the following exact sequence

$$0 \rightarrow G[p^\infty] \rightarrow G \rightarrow \varinjlim_p G \rightarrow 0.$$

We will write $Q := \varinjlim_p G$. By the Hochschild–Serre spectral sequence we have a spectral sequence

$$E_2 = H_p(Q, H_q(G[p^\infty], k)) \implies H_{p+q}(G, k).$$

Since G is abelian and G acts trivially on k , it follows that Q acts trivially on $H_q(G[p^\infty], k)$. Since k is a field $H_q(G[p^\infty], k)$ is a direct sum of copies of k and therefore, we are done by Lemma 3.30. \square

Proposition 3.32. *Let A be an abelian variety over k . Let $A[p^\infty]$ be the associated p -divisible group. Then $H_{crys}^*(B(A[p^\infty])) \simeq \text{Sym}^*(H_{crys}^2(BA[p^\infty]))$, where $H_{crys}^2(BA[p^\infty])$ is considered to be in degree 2.*

Proof. We let \mathcal{X} denote the topos $\text{Shv}(\text{SYNSch}_k)$. Given the abelian variety A , we can associate an object in \mathcal{X} (by considering functor of points) which will also be denoted by A . Similarly, by taking direct limits, we can associate an object in \mathcal{X} corresponding to $A[p^\infty]$ which will again be denoted by $A[p^\infty]$. Since they both are group objects, we can consider the associated classifying objects as simplicial objects in \mathcal{X} denoted respectively by BA and $BA[p^\infty]$. One can also consider the free $W_n(k)$ -module on these objects and take the associated alternating face complex which will be denoted by $W_n(k)[BA]$ and $W_n(k)[BA[p^\infty]]$ respectively which can both be viewed as objects in $D(W_n(k))_{\mathcal{X}}$; the derived category of $W_n(k)$ -modules in \mathcal{X} . We let $R\Gamma_{crys}(BA[p^\infty]/W_n) := R\text{Hom}_{D(W_n(k))_{\mathcal{X}}}(W_n(k)[BA[p^\infty]], \mathcal{O}^{crys}/W_n)$ and $R\Gamma_{crys}(BA/W_n) := R\text{Hom}_{D(W_n(k))_{\mathcal{X}}}(W_n(k)[BA], \mathcal{O}^{crys}/W_n)$.

Lemma 3.33. *There is a natural isomorphism $W_n(k)[BA[p^\infty]] \simeq W_n(k)[BA]$ in $D(W_n(k))_{\mathcal{X}}$. Thus*

$$R\Gamma_{crys}(BA[p^\infty]/W_n) \simeq R\Gamma_{crys}(BA/W_n).$$

Proof. Since multiplication by p map on the abelian variety A is a syntomic cover, it follows that as an abelian group object of \mathcal{X} , multiplication by p is surjective on A . Using the map $A[p^\infty] \rightarrow A$, we obtain a natural map $W_n(k)[BA[p^\infty]] \rightarrow W_n(k)[BA]$ and in order to prove that this map is a quasi-isomorphism, we need to check that $H^i(W_n(k)[BA[p^\infty]]) \rightarrow H^i(W_n(k)[BA])$ is isomorphism. Note that since the topos \mathcal{X} can be generated by affine schemes, it is (locally) coherent, and by Deligne's theorem [AGV72, exposé VI, p.336], it has enough points. Therefore, in order to check that the map $H^i(W_n(k)[BA[p^\infty]]) \rightarrow H^i(W_n(k)[BA])$ is an isomorphism, it is enough to check it by taking stalks at a (geometric) point $x : \text{Sets} \rightarrow \mathcal{X}$. But since taking stalk is an exact functor, it commutes with taking cohomology. Therefore, we can take stalks levelwise on the complexes associated to $W_n(k)[BA[p^\infty]]$ and $W_n(k)[BA]$ and then check that it is a quasi-isomorphism. Again, by noting that taking stalks commutes with the free $W_n(k)$ -module object construction, we are reduced to proving that there is a natural quasi-isomorphism $W_n(k)[BA_x[p^\infty]] \rightarrow W_n(k)[BA_x]$ of complexes of $W_n(k)$ -modules. This follows from Lemma 3.31. \square

By taking inverse limits over n , we obtain $R\Gamma_{crys}(BA[p^\infty]) \simeq R\Gamma_{crys}(BA)$. Now Proposition 3.32 follows from the fact that $H_{crys}^*(BA)$ is a symmetric algebra in $H_{crys}^2(BA)$ by Proposition 3.28. \square

Proposition 3.34. *Let G be any p -divisible group over k . Then $H_{crys}^*(BG) = \text{Sym}^*(H_{crys}^2(BG))$, where $H_{crys}^2(BG)$ is considered to be in degree 2.*

Proof. At first, we assume that the field k is algebraically closed. By [Oor00], there exists a p -divisible group G' such that there is an isomorphism $G \times G' \simeq A[p^\infty]$ for some abelian variety A . Indeed, by taking G' to be the dual of G , we get a p -divisible group $G \times G'$ which has a symmetric Newton polygon; therefore, by Section 5, loc. cit. there exists some abelian variety A' such that $A'[p^\infty]$ and $G \times G'$ have the same Newton Polygon. By the Dieudonné-Manin classification, A' is isogenous to $G \times G'$. Quotienting A' by a finite flat group scheme if necessary, one obtains an abelian variety A as desired.

Using the zero section of G' we have a map $G \rightarrow G \times G'$ which when composed with projection to G gives the identity map. Thus BG is a retract of $BA[p^\infty]$. Since $H_{crys}^*(BA[p^\infty])$ is generated as a (commutative) symmetric algebra in degree 2 classes, it follows that H_{crys}^*BG is generated by degree 2 classes, i.e., there is a surjection $\text{Sym}^*(H_{crys}^2(BG)) \rightarrow H_{crys}^*(BG)$. Doing the same for G' we arrive at the following commutative diagram.

$$\begin{array}{ccc} \text{Sym}^* H_{crys}^2(BG) \otimes \text{Sym}^* H_{crys}^2(BG') & \xrightarrow{\simeq} & \text{Sym}^*(H_{crys}^2(BG \times BG')) \\ \downarrow & & \downarrow \simeq \\ H_{crys}^*(BG) \otimes H_{crys}^*(BG') & \longrightarrow & H_{crys}^*(BG \times BG') \end{array}$$

The upper horizontal map is an isomorphism since $H_{crys}^2(BG)$ is isomorphic to the Dieudonné module (Proposition 3.23) and therefore, $H_{crys}^2(BG \times BG') \simeq H_{crys}^2(BG) \oplus H_{crys}^2(BG')$. By using Proposition 3.32,

the right vertical map is an isomorphism since $BG \times BG' \simeq BA[p^\infty]$. This shows that the surjection $\mathrm{Sym}^*(H_{crys}^2(BG)) \rightarrow H_{crys}^*(BG)$ must be an isomorphism, yielding the assertion of the proposition when k is algebraically closed.

We will now remove the assumption that k is algebraically closed. Let k be any perfect field and \bar{k} denote the algebraic closure of k . Let $G_{\bar{k}}$ denote the base change of G to $\mathrm{Spec} \bar{k}$. Our goal is to show that the natural map $\mathrm{Sym}^*(H_{crys}^2(BG)) \rightarrow H_{crys}^*(BG)$ is an isomorphism. By Proposition 3.23 and taking inverse limits over n , it would be enough to show that $H_{crys}^*(BG/W_n) = \mathrm{Sym}^*(H_{crys}^2(BG/W_n))$. Since we already know that $H_{crys}^*(BG_{\bar{k}}/W_n) = \mathrm{Sym}^*(H_{crys}^2(BG_{\bar{k}}/W_n))$, it would be enough to prove that $R\Gamma_{crys}(BG_{\bar{k}}/W_n) \simeq R\Gamma_{crys}(BG/W_n) \otimes_{W_n(k)} W_n(\bar{k})$. This reduces to the case when $n = 1$, so it is enough to show that $H_{crys}^i(BG_{\bar{k}}/W_1) = H_{crys}^i(BG/W_1) \otimes_k \bar{k}$. We note the following lemma.

Lemma 3.35. *Let M be a finite group scheme over a perfect field k . Then $H_{crys}^i(BM/W_1)$ is a finite dimensional vector space.*

Proof. By using descent along the syntomic map $\mathrm{Spec} k \rightarrow BM$ and [ABM19, Rmk. 2.4], it follows that $H_{crys}^i(BM/W_1) \simeq H_{dR}^i(BM)$. Here $H_{dR}^i(BM)$ denotes the de Rham cohomology of the smooth stack BM as in Construction 2.7 loc. cit. In order to show finiteness of $H_{dR}^i(BM)$, since BM is smooth, by using the Hodge–de Rham spectral sequence from Def. 3.1. (b) loc. cit. it is enough to show finiteness of the Hodge cohomology groups of BM . But this finiteness follows because the cotangent complex of BM and its exterior powers are perfect complexes on BM and M is a finite group scheme. cf. [Tot18, Thm. 3.1]. \square

Let us write G as the directed system $\{G_n\}$ of finite group schemes G_n over k . For a fixed n , we have $H_{crys}^i(B(G_n)_{\bar{k}}/W_1) = H_{crys}^i(BG_n/W_1) \otimes_k \bar{k}$. By the above lemma, we have

$$H_{crys}^i(BG_{\bar{k}}/W_1) = \varprojlim_n H_{crys}^i(B(G_n)_{\bar{k}}/W_1) = \varprojlim_n (H_{crys}^i(BG_n/W_1) \otimes_k \bar{k}).$$

We note that the pro-system $\{H_{crys}^i(BG_n/W_1)\}$ is essentially constant [SP, Tag 05PT]. Indeed, this can be seen by base changing to \bar{k} and noting that the pro-system $\{V_n\} := \{H_{crys}^i(BG_n/W_1) \otimes_k \bar{k}\}$ consists of finite dimensional \bar{k} -vector spaces and $\varprojlim_n V_n$ is finite dimensional as well. The first assertion follows from Lemma 3.35 and the second one follows from the statement of Proposition 3.34 for the algebraically closed field \bar{k} . Therefore,

$$H_{crys}^i(B(G_n)_{\bar{k}}/W_1) = \varprojlim_n (H_{crys}^i(BG_n/W_1) \otimes_k \bar{k}) = \varprojlim_n H_{crys}^i(BG_n/W_1) \otimes_k \bar{k} = H_{crys}^i(BG/W_1) \otimes_k \bar{k},$$

which finishes the proof. \square

Proof of Theorem 1.4. Follows from Proposition 3.34 and Proposition 3.23. \square

4 Prismatic cohomology

4.1 Prismatic cohomology for stacks

In this section, we start by recalling the notion of prismatic cohomology and then extend the notion of prismatic cohomology to stacks. The main reference for this section is [BS19] and [BMS19]. We will freely use the definitions and notations from their paper. Our somewhat terse exposition here is also loosely based on [ALB19]. We will sometimes use the notion of derived p -completion in the proofs; for its basic properties, we refer to [SP, Tag 091N]. More details on notions such as p -complete flatness can be found in [BMS19, §4].

Definition 4.1 ([BMS19, Def. 4.1]). Let $A \rightarrow B$ be a map of commutative rings. We will call this map p -completely (faithfully) flat if $B \otimes_A^L A/p \in D(A/p)$ is concentrated in degree zero, and is a (faithfully) flat A/p -algebra.

Definition 4.2 ([BMS19, Def. 4.1]). Let A be a commutative ring and M be an object in the derived category $D(A)$. Let $a, b \in \mathbb{Z} \sqcup \{\pm\infty\}$. We will say that M has p -complete Tor-amplitude in $[a, b]$ if $M \otimes_A^L A/p \in D(A/p)$ has Tor-amplitude in $[a, b]$. The latter condition means that if we set $M' := M \otimes_A^L A/p$, then $M' \otimes_{A/p}^L N \in D^{[a, b]}(A/p)$ for any A/p -module N .

Definition 4.3 ([BMS19, Def. 4.10]). A ring S is called *quasi-syntomic* if S is p -complete with bounded p^∞ -torsion and the cotangent complex $\mathbb{L}_{S/\mathbb{Z}_p}$ has p -complete Tor-amplitude in $[-1, 0]$. The category of all quasi-syntomic rings is denoted by QSyn . A map $S \rightarrow S'$ of p -complete rings with bounded p^∞ -torsion is a *quasi-syntomic morphism* if S' is p -completely flat over S and the cotangent complex $\mathbb{L}_{S'/S}$ has p -complete Tor-amplitude in $[-1, 0]$. A quasi-syntomic morphism is called a *quasi-syntomic cover* if the map $S \rightarrow S'$ is p -completely faithfully flat.

Remark 4.4. Note that since a quasi-syntomic ring S is assumed to have bounded p^∞ -torsion, the notion of derived p -completeness is equivalent to classical p -completeness in this case. *cf.* [SP, Tag 0923].

Definition 4.5 ([BMS19, Def. 4.18]). A ring S is called (integral) *perfectoid* if S is p -complete, such that $\pi^p = pu$ for some $\pi \in S$, $u \in S^\times$, the Frobenius is surjective on S/p and kernel of the map $\theta : A_{\text{inf}}(S) := W(S^b) \rightarrow S$ is principal.

Definition 4.6 ([BMS19, Def. 4.20]). A ring S is called quasi-regular semiperfectoid (QRSP) if $S \in \text{QSyn}$ and there exists a perfectoid ring R mapping surjectively onto S .

Definition 4.7. If R is any p -complete ring we will let $(R)_{\text{qsyn}}$ denote the (opposite) category of all p -complete rings over R which are quasi-syntomic. This category can be equipped with a Grothendieck topology generated by quasi-syntomic covers which turns this into a site.

Remark 4.8. By the results in [BMS19, §4.4], QRSP rings form a basis for the site $(R)_{\text{qsyn}}$. Therefore, in order to specify a sheaf on $(R)_{\text{qsyn}}$ amounts to assigning values to the QRSP rings satisfying the descent condition.

In [BS19] the authors define prismatic cohomology at first for p -complete smooth algebras and then extend it to all p -complete algebras [BS19, Construction 7.6] and call it the *derived prismatic cohomology*. An important fact [BS19, Prop. 7.10] about this construction is that for every QRSP ring S , the derived prismatic cohomology denoted as Δ_S lives only in degree 0, i.e., is discrete. Also, the functor $S \rightarrow \Delta_S$ forms a quasi-syntomic sheaf [BS19, Construction 7.6. (3)]. Using these facts the authors in [ALB19] construct a sheaf of rings $\mathcal{O}^{\text{pris}}$ on $(R)_{\text{qsyn}}$ such that for any QRSP ring S over R , one has

$$\Delta_S \simeq R\Gamma(S, \mathcal{O}^{\text{pris}}).$$

Further, we note that for any prism (A, I) , there is a decreasing filtration $\mathcal{N}^{\geq \bullet} A$ called the Nygaard filtration which is defined as

$$\mathcal{N}^{\geq i}(A) := \varphi^{-1}(I^i).$$

This equips the sheaf of rings $\mathcal{O}^{\text{pris}}$ with a decreasing filtration $\mathcal{N}^{\geq \bullet} \mathcal{O}^{\text{pris}}$ of sheaves.

Remark 4.9. We explain the construction of the Nygaard filtration in detail. Using [ALB19, Cor. 3.3.10], we have a morphism of topoi:

$$v : \mathrm{Shv}((R)_{\Delta}) \rightarrow \mathrm{Shv}((R)_{\mathrm{qsyn}}),$$

where $(R)_{\Delta}$ denotes the absolute prismatic site as in Def. 3.1.4 loc. cit. On $(R)_{\Delta}$ one defines a sheaf of rings \mathcal{O}_{Δ} by sending a prism $(A, I) \rightarrow A$. Using the notion of Nygaard filtration on a prism one can equip this sheaf of rings \mathcal{O}_{Δ} with the Nygaard filtration $\mathcal{N}^{\geq \bullet} \mathcal{O}_{\Delta}$ which sends a prism $(A, I) \rightarrow \mathcal{N}^{\geq \bullet} A \subseteq A$. In order to prove that the presheaf $(A, I) \rightarrow \mathcal{N}^{\geq \bullet} A$ is indeed a sheaf, we note that $\mathcal{N}^{\geq i}(A)$ is the kernel of a map of presheaf of rings $((A, I) \rightarrow A) \rightarrow ((A, I) \rightarrow A/I^i)$ obtained by composing $\varphi : A \rightarrow A$ with the projection map $A \rightarrow A/I^i$. Therefore it will be enough to show that the presheaf $(A, I) \rightarrow A/I^i$ is a sheaf. This follows from the proof of [BS19, Cor. 3.12] by noting that for a map $(A, I) \rightarrow (B, J)$ of prisms one has $I^i B = J^i$ by Lemma 3.5 loc. cit. One defines

$$\mathcal{N}^{\geq \bullet} \mathcal{O}^{\mathrm{pris}} := v_* \mathcal{N}^{\geq \bullet} \mathcal{O}_{\Delta}.$$

By [BS19, Prop. 7.2] and the above definition, it follows that $\mathcal{N}^{\geq \bullet} \mathcal{O}^{\mathrm{pris}}(S) \simeq \mathcal{N}^{\geq \bullet} \Delta_S$ for a QRSP ring $S \in (R)_{\mathrm{qsyn}}$. In fact, by Remark 4.8 this description can be used to define the sheaves $\mathcal{N}^{\geq \bullet} \mathcal{O}^{\mathrm{pris}}$ after one proves that it forms a sheaf on the basis objects. However, a priori it is not obvious, but, actually, more is true: the functor that sends a QRSP algebra $S \rightarrow \mathcal{N}^{\geq \bullet} \Delta_S$ forms a sheaf with vanishing higher cohomology, i.e., $H^i(S, \mathcal{N}^{\geq \bullet} \mathcal{O}^{\mathrm{pris}}) = 0$ for $i \geq 1$ and a QRSP algebra S . This fact follows from [BS19, Thm. 12.2] and [BMS19, Thm. 3.1].

Definition 4.10. We call \mathcal{Y} a quasi-syntomic stack over R if it is a stack with respect to the site $(R)_{\mathrm{qsyn}}$.

Remark 4.11. If $A \in (R)_{\mathrm{qsyn}}$ then by p -completely faithfully flat descent it follows that h_A is a sheaf with respect to $(R)_{\mathrm{qsyn}}$. If \mathcal{Y} is a quasi-syntomic stack, an arrow $h_A \rightarrow \mathcal{Y}$ will be denoted as $\mathrm{Spf} A \rightarrow \mathcal{Y}$.

Remark 4.12. Let X be a p -adic formal scheme over R . Then setting $\underline{X}(A) := \mathrm{Hom}_{\mathrm{formal\ sch}}(\mathrm{Spf} A, X)$ defines a quasi-syntomic sheaf on $(R)_{\mathrm{qsyn}}$ which we will denote by \underline{X} .

Definition 4.13. Let \mathcal{Y} be a quasi-syntomic stack. We define the *prismatic cohomology* of \mathcal{Y} to be

$$R\Gamma_{\Delta}(\mathcal{Y}) := R \varprojlim_{\mathrm{Spf} A \rightarrow \mathcal{Y}} \Delta_A,$$

where $A \in (R)_{\mathrm{qsyn}}$ and the derived limit is taken in the derived category of abelian groups.

Remark 4.14. Let \mathcal{Y} be a quasi-syntomic stack. We give an alternative description of $R\Gamma_{\Delta}(\mathcal{Y})$. One can define the quasi-syntomic site of \mathcal{Y} denoted as $(\mathcal{Y})_{\mathrm{qsyn}}$ to be the category whose objects are $\mathrm{Spf} A \rightarrow \mathcal{Y}$ for $A \in (R)_{\mathrm{qsyn}}$ equipped with the quasi-syntomic covers, i.e., we let $(\mathcal{Y})_{\mathrm{qsyn}} := (R)_{\mathrm{qsyn}/\mathcal{Y}}$. Then by the discussion preceding Definition 4.10, we obtain a sheaf of rings $\mathcal{O}^{\mathrm{pris}}$ on $(\mathcal{Y})_{\mathrm{qsyn}}$. Then it follows that $R\Gamma_{\Delta}(\mathcal{Y}) \simeq R\Gamma(\mathcal{Y}, \mathcal{O}^{\mathrm{pris}})$.

Definition 4.15. Let \mathcal{Y} be a quasi-syntomic stack. The *Nygaard filtration* on prismatic cohomology $R\Gamma_{\Delta}(\mathcal{Y})$ is defined to be

$$\mathcal{N}^{\geq \bullet} R\Gamma_{\Delta}(\mathcal{Y}) := R\Gamma(\mathcal{Y}, \mathcal{N}^{\geq \bullet} \mathcal{O}^{\mathrm{pris}}).$$

Definition 4.16. Now we consider an algebraic stack \mathcal{X} over R . We will define the p -adic completion of \mathcal{X} as a quasi-syntomic stack which will be denoted as $\widehat{\mathcal{X}}$. We define

$$\widehat{\mathcal{X}}(A) := \varprojlim \mathcal{X}(A/p^n),$$

where $A \in (R)_{\mathrm{qsyn}}$. This defines a sheaf of groupoids in the site $(R)_{\mathrm{qsyn}}$. Indeed, $\widehat{\mathcal{X}}$ by definition is an inverse limit of the sheaves on $(R)_{\mathrm{qsyn}}$ which sends $A \rightarrow \mathcal{X}(A/p^n)$ and therefore has to be a sheaf.

Definition 4.17. We define the prismatic cohomology of a stack \mathcal{X} to be

$$R\Gamma_{\Delta}(\mathcal{X}) := R\Gamma_{\Delta}(\widehat{\mathcal{X}}).$$

Proposition 4.18. *If X is a p -adic formal scheme over R , then $R\Gamma_{\Delta}(X) \simeq R\mathrm{Hom}_{D((R)_{\mathrm{qsyn}})}(\mathbb{Z}[\underline{X}], \mathcal{O}^{\mathrm{pris}})$, where \underline{X} is the quasi-syntomic sheaf associated to X on $(R)_{\mathrm{qsyn}}$.*

Proof. This follows from Remark 4.12 and Remark 4.14 by adjunction. \square

Remark 4.19. Instead of considering quasi-syntomic stacks, one could have also worked with “ p -adic formal stacks” which can be defined to be stacks with respect to the Grothendieck site on the category of p -complete and bounded p^{∞} -torsion rings equipped with p -completely faithfully flat covers. Any p -adic formal stack can also be regarded as a quasi-syntomic stack. The reason we work with the notion of quasi-syntomic stacks is because the notion of prismatic cohomology in Definition 4.13 would ultimately regard a p -adic formal stack as a quasi-syntomic stack. Hence one might as well define prismatic cohomology for quasi-syntomic stacks.

4.2 Application to prismatic Dieudonné theory

From now we will assume that the ring R is a QRSP ring. Let (Δ_R, I) be the prism associated to R by taking prismatic cohomology. We briefly recall some definitions from [ALB19] and review their theorem on classification of p -divisible groups in terms of filtered prismatic Dieudonné modules.

Definition 4.20 ([ALB19, Def. 4.1.10]). A filtered prismatic Dieudonné module over R is a collection $(M, \mathrm{Fil} M, \varphi_M)$ consisting of a finite locally free Δ_R -module M , a Δ_R -submodule $\mathrm{Fil} M$, and φ linear map $\varphi_M : M \rightarrow M$ satisfying

- (1) $\varphi_M(\mathrm{Fil} M) \subset IM$ and $\varphi_M(\mathrm{Fil} M)$ generates IM as a Δ_R -module
- (2) $(\mathcal{N}^{\geq 1} \Delta_R)M \subset \mathrm{Fil} M$ and $M/\mathrm{Fil} M$ is a finite locally free R -module.

Now let G be a p -divisible group. One makes the following definition.

Definition 4.21. The quasi-syntomic sheaf $\underline{G} \in (R)_{\mathrm{qsyn}}$ associated to a p -divisible group G is defined to be $\varinjlim \underline{G}_n$ where $\underline{G}_n \in (R)_{\mathrm{qsyn}}$ is the quasi-syntomic sheaf associated to $\mathrm{Spf} G_n$.

The authors in [ALB19] make the following definition which can be seen as a generalization of Theorem 3.8.

Definition 4.22 ([ALB19, Def. 4.2.8]). Let G be a p -divisible group over R . We define

$$M_{\Delta}(G) := \mathrm{Ext}_{(R)_{\mathrm{qsyn}}}^1(\underline{G}, \mathcal{O}^{\mathrm{pris}}),$$

$$\mathrm{Fil} M_{\Delta}(G) := \mathrm{Ext}_{(R)_{\mathrm{qsyn}}}^1(\underline{G}, \mathcal{N}^{\geq 1} \mathcal{O}^{\mathrm{pris}}),$$

and $\varphi_{M_{\Delta}(G)}$ as the endomorphism induced by φ on $\mathcal{O}^{\mathrm{pris}}$. Then

$$\underline{M}_{\Delta}(G) := (M_{\Delta}(G), \mathrm{Fil} M_{\Delta}(G), \varphi_{M_{\Delta}(G)})$$

is called the *filtered prismatic Dieudonné module* of G .

In this case, the main theorem of [ALB19] says the following.

Theorem 4.23. *The filtered prismatic Dieudonné module functor*

$$G \rightarrow \underline{M}_{\Delta}(G)$$

defines an anti-equivalence between the category p -divisible groups over R and the category of filtered prismatic Dieudonné modules over R .

Example 4.24. For the étale p -divisible group $\mathbb{Q}_p/\mathbb{Z}_p$ over R , we have $\underline{M}_{\Delta}(\mathbb{Q}_p/\mathbb{Z}_p) \simeq (\Delta_R, \mathcal{N}^{\geq 1} \Delta_R, \varphi)$. We refer to [ALB19, §4.7] and Rmk. 4.9.6 loc. cit. for more discussions.

Example 4.25. Let \hat{A} denote the p -adic completion of an abelian scheme A over R . Let us denote the p -divisible group associated to \hat{A} by $A[p^{\infty}]$ for simplicity. In this case, the prismatic Dieudonné module $M_{\Delta}(A[p^{\infty}])$ is locally free of rank $2 \cdot \dim A$ and is isomorphic to $H_{\Delta}^1(\hat{A})$ which identifies the natural Frobenius on both of these modules as well [ALB19, Thm 4.5.6]. Further, by using the proof of Prop. 4.5.9 loc. cit. one obtains that $\mathrm{Fil} M_{\Delta}(A[p^{\infty}]) \simeq \mathcal{N}^{\geq 1} H_{\Delta}^1(\hat{A})$, where $\mathcal{N}^{\geq 1} H_{\Delta}^1(\hat{A}) := H^1(\mathcal{N}^{\geq 1} R\Gamma_{\Delta}(\hat{A}))$. cf. Proposition 4.39.

Now we proceed towards proving Theorem 1.6. Let $G = \text{Spec } B$ be a finite flat group scheme over R . Since G is syntomic, it follows that $\text{Spf}(B)$ is quasi-syntomic. If BG denotes the associated quasi-syntomic stack, then $* \rightarrow BG$ is a G -torsor and is a quasi-syntomic cover. The Čech nerve of the map $* \rightarrow BG$ is given by the following simplicial quasi-syntomic sheaf

$$\cdots \underline{G} \times \underline{G} \times \underline{G} \rightrightarrows \underline{G} \times \underline{G} \rightrightarrows \underline{G} \rightrightarrows * .$$

The associated simplicial abelian group object is

$$\cdots \mathbb{Z}[\underline{G} \times \underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G}] \rightrightarrows \mathbb{Z} .$$

With this simplicial object we can associate an object of $D((R)_{\text{qsyn}})$ which we will denote simply by $\mathbb{Z}[BG]$.

Proposition 4.26. *Let G be a finite flat group scheme over R . Then $R\Gamma_{\Delta}(BG) \simeq R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG], \mathcal{O}^{\text{pris}})$.*

Proof. By Čech descent along the quasi-syntomic cover $* \rightarrow \underline{BG}$ we obtain that

$$R\Gamma_{\Delta}(BG) \simeq R\varprojlim (R\Gamma_{\Delta}(*) \rightrightarrows R\Gamma_{\Delta}(G) \rightrightarrows R\Gamma_{\Delta}(G \times G) \cdots).$$

Which by Proposition 4.18 is

$$\simeq R\varprojlim (R\text{Hom}_{D(R)}(\mathbb{Z}, \mathcal{O}^{\text{pris}}) \rightrightarrows R\text{Hom}_{D(R)}(\mathbb{Z}[\underline{G}], \mathcal{O}^{\text{pris}}) \rightrightarrows R\text{Hom}_{D(R)}(\mathbb{Z}[\underline{G} \times \underline{G}], \mathcal{O}^{\text{pris}}) \cdots).$$

We can take the $R\varprojlim$ inside as a homotopy colimit, which gives us that the above is $\simeq R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG], \mathcal{O}^{\text{pris}})$, as desired. \square

Definition 4.27. The classifying stack of a p -divisible group $G = \{G_n\}$ is defined to be $BG := \varinjlim BG_n$ where the filtered colimit is taken in the category of quasi-syntomic stacks with respect to the site $(\overline{R})_{\text{qsyn}}$.

Proposition 4.28. *For a p -divisible group $G = \{G_n\}$ we have $R\Gamma_{\Delta}(BG) \simeq R\varprojlim R\Gamma_{\Delta}(BG_n)$.*

Proof. This follows because BG is defined to be the filtered colimit of the quasi-syntomic stacks BG_n . \square

Since filtered colimits commute with finite limits, the Čech nerve of $* \rightarrow BG$ is given by the simplicial quasi-syntomic sheaf

$$\cdots \underline{G} \times \underline{G} \times \underline{G} \rightrightarrows \underline{G} \times \underline{G} \rightrightarrows \underline{G} \rightrightarrows * .$$

The associated simplicial abelian group object is

$$\cdots \mathbb{Z}[\underline{G} \times \underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G} \times \underline{G}] \rightrightarrows \mathbb{Z}[\underline{G}] \rightrightarrows \mathbb{Z} .$$

With this simplicial object, we can associate an object of $D((R)_{\text{qsyn}})$ which will be denoted by $\mathbb{Z}[BG]$. Since filtered colimits are exact in the category of abelian sheaves on $(R)_{\text{qsyn}}$, we have that $\varinjlim \mathbb{Z}[BG_n] \simeq \mathbb{Z}[BG]$.

Proposition 4.29. *Let G be a p -divisible group over R . Then $R\Gamma_{\Delta}(BG) \simeq R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG], \mathcal{O}^{\text{pris}})$.*

Proof. By Proposition 4.28 and Proposition 4.26 we have

$$R\Gamma_{\Delta}(BG) \simeq R\varprojlim R\Gamma_{\Delta}(BG_n) \simeq R\varprojlim R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG_n], \mathcal{O}^{\text{pris}}) .$$

Now

$$R\varprojlim R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG_n], \mathcal{O}^{\text{pris}}) \simeq R\text{Hom}_{D((R)_{\text{qsyn}})}(\varinjlim \mathbb{Z}[BG_n], \mathcal{O}^{\text{pris}}) .$$

Since $\varinjlim \mathbb{Z}[BG_n] \simeq \mathbb{Z}[BG]$, we obtain the required statement. \square

Remark 4.30. Alternatively, one could descend along the effective epimorphism $* \rightarrow BG$.

Proposition 4.31. *There is a spectral sequence with E_2 -page*

$$E_2^{i,j} = \text{Ext}_{(R)_{\text{qsyn}}}^i(H^{-j}(\mathbb{Z}[BG]), \mathcal{O}^{\text{pris}}) \implies H_{\Delta}^{i+j}(BG),$$

and another spectral sequence with E_1 -page

$$E_1^{i,j} = H_{\Delta}^j(G^i) \implies H_{\Delta}^{i+j}(BG),$$

where G^i denotes the i fold fibre product of $\text{Spf}G$ with itself. By convention, $G^0 = *$.

Proof. This follows in a way similar to Proposition 3.13. \square

Lemma 4.32. $H^0(\mathbb{Z}[BG]) \simeq \mathbb{Z}$, $H^{-1}(\mathbb{Z}[BG]) \simeq \underline{G}$, and $H^{-2}(\mathbb{Z}[BG]) \simeq \underline{G} \wedge \underline{G}$.

Proof. This follows exactly in the same ways as in the proof of Lemma 3.9 after further noting that for an abelian group G , the second group homology with integral coefficients $H_2(G, \mathbb{Z}) \simeq G \wedge G$. This isomorphism can be found in [CM52, Thm. 3]. \square

Proposition 4.33. *There is a natural isomorphism $H_{\Delta}^2(BG) \simeq \text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{G}, \mathcal{O}^{\text{pris}})$.*

Proof. By the E_2 spectral sequence from Proposition 4.31 and Lemma 4.32, this will follow once we prove that $\text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \wedge \underline{G}, \mathcal{O}^{\text{pris}}) = 0$ and $\text{Ext}_{(R)_{\text{qsyn}}}^i(\mathbb{Z}, \mathcal{O}^{\text{pris}}) = 0$ for $i \geq 1$. We begin by proving the first vanishing. We have a surjection $\underline{G} \otimes \underline{G} \rightarrow \underline{G} \wedge \underline{G} \rightarrow 0$ which gives us an injection

$$0 \rightarrow \text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \wedge \underline{G}, \mathcal{O}^{\text{pris}}) \rightarrow \text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \otimes \underline{G}, \mathcal{O}^{\text{pris}}).$$

Thus it is enough to show that $\text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \otimes \underline{G}, \mathcal{O}^{\text{pris}}) = 0$. Indeed,

$$\text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \otimes \underline{G}, \mathcal{O}^{\text{pris}}) \simeq \text{Hom}_{(R)_{\text{qsyn}}}(\underline{G}, \mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{G}, \mathcal{O}^{\text{pris}})),$$

and $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{G}, \mathcal{O}^{\text{pris}}) = 0$ since G is p -divisible and $\mathcal{O}^{\text{pris}}$ is derived p -complete. Therefore, we obtain the required statement. Now the second vanishing follows from the fact that for a QRSP ring S , its prismatic cohomology Δ_S is discrete, i.e., lives only in degree 0. This finishes the proof. \square

Proposition 4.34. *In the above situation, $\mathcal{N}^{\geq 1}H_{\Delta}^2(BG) := H^2(\mathcal{N}^{\geq 1}R\Gamma_{\Delta}(BG)) \subset H_{\Delta}^2(BG)$ and $\mathcal{N}^{\geq 1}H_{\Delta}^2(BG) \simeq \text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{G}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}})$.*

Proof. By Definition 4.15, we have $\mathcal{N}^{\geq 1}R\Gamma_{\Delta}(BG) = R\Gamma(BG, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}})$. Hence, analogous to Proposition 4.26, by descent along the effective epimorphism $* \rightarrow BG$, we obtain

$$\mathcal{N}^{\geq 1}R\Gamma_{\Delta}(BG) \simeq R\text{Hom}_{D((R)_{\text{qsyn}})}(\mathbb{Z}[BG], \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}}).$$

By the spectral sequence analogous to Proposition 4.31, we obtain that in order to prove $\mathcal{N}^{\geq 1}H_{\Delta}^2(BG) \simeq \text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{G}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}})$, by Lemma 4.32 it is enough to prove that $\text{Hom}_{(R)_{\text{qsyn}}}(\underline{G} \wedge \underline{G}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}}) = 0$ and $\text{Ext}_{(R)_{\text{qsyn}}}^i(\mathbb{Z}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}}) = 0$ for $i \geq 2$. By the proof of Proposition 4.33, in order to prove the first vanishing it is enough to prove that $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{G}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}}) = 0$. By the injection $0 \rightarrow \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}} \rightarrow \mathcal{O}^{\text{pris}}$ of sheaves, the required vanishing follows from the fact that $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{G}, \mathcal{O}^{\text{pris}}) = 0$ which was noted in proof of Proposition 4.33. Now for proving $\text{Ext}_{(R)_{\text{qsyn}}}^i(\mathbb{Z}, \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}}) = 0$ for $i \geq 2$, using the exact sequence

$$0 \rightarrow \mathcal{N}^{\geq 1}\mathcal{O}^{\text{pris}} \rightarrow \mathcal{O}^{\text{pris}} \rightarrow \mathcal{O} \rightarrow 0,$$

from [ALB19, Prop. 4.1.2], where \mathcal{O} denotes the structure sheaf on $(R)_{\text{qsyn}}$, it is enough to show that $\text{Ext}_{(R)_{\text{qsyn}}}^i(\mathbb{Z}, \mathcal{O}) = 0$ for $i \geq 1$ which follows from p -completely faithfully flat descent for bounded p^{∞} -torsion rings [BMS19, Remark. 4.9]. The inclusion $\mathcal{N}^{\geq 1}H_{\Delta}^2(BG) \subset H_{\Delta}^2(BG)$ now follows from the exact sequence above and the fact that $\text{Hom}(\underline{G}, \mathcal{O}) = 0$ since G is p -divisible and \mathcal{O} is derived p -complete. \square

Proof of Theorem 1.6. Follows from Proposition 4.33 and Proposition 4.34. \square

Remark 4.35. We note that $\text{Ext}^1(\mathbb{Z}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}) = 0$ as well. This follows from the exact sequence above and [ALB19, Thm. 3.4.6].

Remark 4.36. Similar to Proposition 3.14, it follows that for a p -divisible group G over R , we have $H_{\Delta}^1(BG) \simeq 0$.

Remark 4.37. One can define prismatic cohomology for higher quasi-syntomic stacks as well. In particular one can talk about prismatic cohomology of the n -stacks $K(G, n)$ for a p -divisible group G . Similar to Remark 3.17, one can prove that $H_{\Delta}^{n+1}(BG) \simeq M_{\Delta}(G)$ and $H_{\Delta}^n(BG) = 0$ for $1 \leq i \leq n$.

Now we look at the case where \widehat{A} is the p -adic completion of an abelian scheme A over R . In this situation one can consider the p -divisible group associated to \widehat{A} which will be written as $A[p^{\infty}]$. We let $B\widehat{A}$ denote the classifying stack of \widehat{A} which we define to be the p -adic completion of the stack BA . The quasi-syntomic sheaf represented by \widehat{A} will simply be written as \underline{A} . Then

Proposition 4.38. *We have $H_{\Delta}^2(B\widehat{A}) \simeq \text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{A}, \mathcal{O}^{\text{pris}})$ and $\mathcal{N}^{\geq 1} H_{\Delta}^2(B\widehat{A}) \simeq \text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{A}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}})$.*

Proof. This follows exactly in the same way as the proof of Proposition 4.33 after noting that $\text{Ext}_{(R)_{\text{qsyn}}}^i(\mathbb{Z}, \mathcal{O}^{\text{pris}}) = 0$ for $i \geq 1$ as before and $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{A}, \mathcal{O}^{\text{pris}}) = 0$ by [ALB19, Thm. 4.5.6]. The second part follows. \square

Proposition 4.39. *$H_{\Delta}^2(B\widehat{A})$ is naturally isomorphic to the prismatic Dieudonné module associated to the p -divisible group $A[p^{\infty}]$. Further, $\mathcal{N}^{\geq 1} H_{\Delta}^2(B\widehat{A}) \simeq \text{Fil } M_{\Delta}(A[p^{\infty}])$.*

Proof. Using Proposition 4.38, it is enough to prove that $\text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{A}, \mathcal{O}^{\text{pris}})$ is isomorphic to $\text{Ext}_{(R)_{\text{qsyn}}}^1(\underline{A[p^{\infty}]}, \mathcal{O}^{\text{pris}})$. Note that we have

$$\mathcal{E}xt_{(R)_{\text{qsyn}}}^1(\underline{A[p^{\infty}]}, \mathcal{O}^{\text{pris}}) \simeq \mathcal{E}xt_{(R)_{\text{qsyn}}}^1(\underline{A}, \mathcal{O}^{\text{pris}})$$

and

$$\mathcal{E}xt_{(R)_{\text{qsyn}}}^1(\underline{A[p^{\infty}]}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}) \simeq \mathcal{E}xt_{(R)_{\text{qsyn}}}^1(\underline{A}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}).$$

Indeed, to see the above isomorphisms, we note that $\underline{A}/\underline{A[p^{\infty}]} = \varinjlim_p \underline{A}$, and for any sheaf \mathcal{F} , we have

$$R\mathcal{H}om(\varinjlim_p \underline{A}, \mathcal{F}) \simeq R\varprojlim_p R\mathcal{H}om(\underline{A}, \mathcal{F}) \simeq R\mathcal{H}om(\underline{A}, R\varprojlim_p \mathcal{F}).$$

Now if \mathcal{F} is further assumed to be a sheaf of derived p -complete abelian groups with vanishing higher cohomology on a collection of basis objects, then $R\varprojlim_p \mathcal{F} \simeq 0$. To see this, it is enough to prove that for any such basis object V , one has $R\Gamma(V, R\varprojlim_p \mathcal{F}) \simeq R\varprojlim_p R\Gamma(V, \mathcal{F}) \simeq 0$. But this follows from our assumptions since $R\Gamma(V, \mathcal{F}) \simeq \mathcal{F}(V)$ and $\mathcal{F}(V)$ is derived p -complete [SP, Tag 091N]. Finally, by taking \mathcal{F} to be $\mathcal{O}^{\text{pris}}$ or $\mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}$ and using Remark 4.9 we obtain the required vanishings to yield the desired isomorphisms of $\mathcal{E}xt$ -groups above.

The proposition now follows by applying the global section functor and noting that $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{A[p^{\infty}]}, \mathcal{O}^{\text{pris}}) \simeq 0$, $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{A}, \mathcal{O}^{\text{pris}}) \simeq 0$ and $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{A[p^{\infty}]}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}) \simeq 0$, $\mathcal{H}om_{(R)_{\text{qsyn}}}(\underline{A}, \mathcal{N}^{\geq 1} \mathcal{O}^{\text{pris}}) \simeq 0$. These vanishings have been noted before in the proofs of Proposition 4.33, Proposition 4.34 and Proposition 4.38. This ends the proof. \square

As in Section 3.3, if \widehat{A} is the p -adic completion of an abelian scheme A over R , we can explicitly compute $H_{\Delta}^*(B(\widehat{A}))$.

Proposition 4.40. *We have a natural isomorphism $H_{\Delta}^{2*}(B\widehat{A}) \cong \text{Sym}^*(H_{\Delta}^1(\widehat{A}))$, and $H_{\Delta}^i(B(\widehat{A})) = 0$ for odd i .*

Proof. This follows exactly in the same way as in the proof in the crystalline case in Section 3.3 from the E_1 spectral sequence in Proposition 4.31, Künneth formula in prismatic cohomology [ALB19, Corollary 3.5.2] and the calculations of Prismatic cohomology of abelian varieties in Corollary 4.5.8 loc. cit. \square

References

- [ALB19] J. Anschütz and A. Le Bras, *Prismatic Dieudonné theory*, available at arXiv:1907.10525
- [ABM19] B. Antieau, B. Bhatt and A. Mathew, *Counterexamples to Hochschild-Kostant-Rosenberg in characteristic p* , available at arXiv:1909.11437
- [AGV72] M. Artin, A. Grothendieck, J. L. Verdier (eds.), *Théorie des Topos et Cohomologie Etale des Schémas - SGA 4. II*, LNM **270** Springer Heidelberg 1972.
- [Beh03] K. A. Behrend, *Derived l -adic categories for algebraic stacks*, Mem. Amer. Math. Soc., 163(774), 2003.
- [BBM82] P. Berthelot, L. Breen and W. Messing, *Théorie de Dieudonné cristalline. II*. Lecture Notes in Mathematics, Vol. 930. Springer-Verlag, Berlin, 1982.
- [BMS19] B. Bhatt, M. Morrow, and P. Scholze, *Topological Hochschild homology and integral p -adic Hodge theory*, Publications mathématiques de l’IHÉS 129 (2019), no. 1, 199310.
- [BS19] B. Bhatt and P. Scholze, *Prisms and prismatic cohomology*, available at arXiv:1905.08229
- [Bo53] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Annals of Mathematics, 57:115207, 1953
- [Dem72] M. Demazure, *Lectures on p -divisible groups*, Lecture Notes in Mathematics, Vol. 302. Springer-Verlag, Berlin-New York, 1972.
- [DP61] A. Dold and D. Puppe, *Homologie nicht-additiver Funktoren, Anwendungen*, Ann. Inst. Fourier, 11 (1961), 201-312.
- [Fo77] J.-M. Fontaine. *Groupes p -divisibles sur les corps locaux*, Astérisque, No. 47-48. Paris: Société Mathématique de France, 1977
- [FM87] J.-M. Fontaine and W. Messing, *p -adic periods and p -adic étale cohomology*, Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), Contemp. Math., vol. 67, Amer. Math. Soc., Providence, RI, 1987, pp. 179-207
- [Ill71] L. Illusie, *Complexe cotangent et déformations, I*. Lecture Notes in Mathematics 239, Springer (1971).
- [CM52] C. Miller, *The Second Homology Group of a Group; Relations among Commutators*. Proceedings of the American Mathematical Society 3, no. 4 (1952) 588-95.
- [Ols07] M. Olsson, *Crystalline cohomology of algebraic stacks and Hyodo-Kato cohomology*, ASTÉRIQUE **316** (2007) 412 pp.
- [Oor66] F. Oort, *Commutative group schemes*, Lectures Notes in Mathematics, Vol. 15. Springer-Verlag, Berlin, 1966.
- [Oor00] F. Oort, *Newton polygons and formal groups: conjectures by Manin and Grothendieck*, Ann. of Math. (2) **152** (2000), no. 1, 183-206.
- [SW14] P. Scholze and J. Weinstein, *Berkeley lectures on p -adic geometry*, available at <http://www.math.uni-bonn.de/people/scholze/Berkeley.pdf>
- [SP] The Stacks Project Authors. available at <https://stacks.math.columbia.edu/>
- [To06] B. Toën, *Higher and derived stacks: A global overview*, Algebraic Geometry-Seattle 2005, Proc. Sympos. Pure Math. 80. (2006)
- [Tot18] B. Totaro, *Hodge theory of classifying stacks*, Duke Mathematical Journal, **167** (2018), 1573-1621.